

Soil Ameliorant in Post Tin Mining Land of Bangka Island for Pepper Growth

Ismed Inonu

Department of Agrotechnology
University of Bangka Belitung
Pangkalpinang, Indonesia
ismedinonu@yahoo.co.id

Rion Apriyadi

Department of Agrotechnology
University of Bangka Belitung
Pangkalpinang, Indonesia
rion_apriyadi@yahoo.com

Ridwan Diaguna

Department of Agrotechnology
University of Bangka Belitung
Pangkalpinang, Indonesia
ridwandiaguna@gmail.com

Muhamad Rifal Rinaldi

Department of Agrotechnology
University of Bangka Belitung
Pangkalpinang, Indonesia
rifalrinaldi@yahoo.com

Abstract—Pepper is one of the spices plant and its production countinuous decrease caused by converting of the planting areas to tin-mining areas. One of the effort that could be done was the usage of sub-optimal land which has low fertility. The research was aimed to study the effect of soil ameliorant application for early growth of pepper plant in the sand tailing area and obtain the proper ameliorant. The research was carried from February to May 2017 in post tin mining land of PT Timah, Dwi Makmur Village, District of Merawang, Bangka Island. The research was arranged in a Randomized Block Design (RBD) single factor dan three replications. The factor used was soil ameliorant combination consisted of seven level i.e. NPK fertilizer, arbuscular mycorrhiza fungi (AMF), biofertilizer, NPK + biofertilizer, NPK + AMF, AMF + organic fertilizer, and NPK + AMF + organic fertilizer. The soil ameliorant applications showed significant effect to plant height, leaves number, segment number, stem diameter, leaf area and chlorophyll content. Complete combination (NPK + AMF + biofertilizer) was the best ameliorant to pepper growth in sand tailing area – post tin mining.

Keywords—ameliorant, AMF, NPK, biofertilizer, pepper, post tin mining land

I. INTRODUCTION

Production of white pepper has continued to decrease in the recent years. It's caused by the decrease of planting area that relate to the high of deforestation by mining activity. The planting area of white pepper decreases of 19.2% in last ten years, from 2004 to 2014, i.e. 201.484 ha in 2004 and 162.751 ha in 2014 [1]

In the future, ex-mining land is one of the biggest challenge of white pepper cultivation. According to [2], Bangka Belitung has post-tin mining land about 390.000 ha, and it's very potential to be utilized. Generally, post-tin mining land is sand tailing site which was dominated by sand fraction about 81% [3], 82% [4], 92% [5], 93% [6] and 96% [7]. Domination of sand tailing leads the low of soil water holding capacity about 5-10% [8], and its capacity just supply about 50-100 mm of the crops water requirement [4]. The soils has characteristics like these, describe the low of fertility, well on physical, chemical, and biology properties.

The utilization of post tin mining land for pepper cultivation requires of adding soil ameliorant. The soil ameliorant is required to improve the soil properties, viz physical, chemical, and biological, furthermore to supply the nutrition, directly. The soil ameliorant that could be used for improving the fertility of post-tin mining land among others; organic fertilizer [9, 10], inorganic fertilizer [11, 12], and mycorrhiza fungi [4].

Organic and biological fertilizers could increase the plant growth [13, 14], increased the soil biology activity [15], and could decreased the use of NPK fertilizer [16, 17, 18]. Furthermore, the adding of organic fertilizer increased the absorption of rainfall, slowed the run-off, decreased the evaporation rate and increased the plant production [19]. Biological agent fertilizers also could increase the physical and chemical soil properties [20].

In the post-tin mining land that has poor nutrient, inorganic fertilizer play a role to supply the nutrient for plant development, directly [21]. The use of Urea, SP-36, and KCl fertilizers could increase the growth and corn yield in the post-tin mining land [22]. Furthermore, application of NPK fertilizer increased the growth of shorgum [23], lettuce [11], and *Samanea saman* [15] which were cultivated in post-tin mining land.

Mycorrhiza is one of the soil fungi, and according to [24], could association with more than 80% types of crop, like forestry, agriculture, horticulture, and plantation. Application of mycorrhiza was required to rehabilitation of soil biology properties and mycorrhiza [25, 26]. Moreover, mycorrhiza could increases the ability of water uptake and nutrient [27]. The use of mycorrhiza as a soil ameliorant in post-tin mining land can improve the soil physical properties [28], chemical and biology properties [29]. Mycorrhiza could increase the rubber growth in post-tin mining land [4].

The adding of three materials as a soil ameliorant is expected could increase the fertility of post-tin mining land. The increasing of fertility will promote the white pepper growth in the land. To date, the type of soil ameliorant that could promote the white pepper growth in post-tin mining land hasn't been informed. The research was aimed to study

the soil ameliorant effect for early growing of white pepper in the sand tailing area and obtain the proper ameliorant.

II. METHODS

Schedule and site

The research was carried out from February to May 2017 in post-tin mining land of PT TIMAH Tbk which was located in Merawang, Bangka. The land used is 10 years old after post mining.

Experiment design

A Randomized Block Design (RBD) with single factor and three replications was applied in the research. The factor was soil ameliorant combination, i.e nitrogen-phosphor-kalium (NPK) fertilizer 'Mutiar', arbuscular mycorrhiza fungi (AMF) genus *Glomus* in zeolit carrier medium, and biofertilizer (*Nano Bio*). The combination consisted of seven levels i.e. B1 (60 g NPK plant⁻¹), B2 (20 g AMF plant⁻¹), B3 (2 cc biofertilizer l⁻¹ water), B4 (60 g NPK plant⁻¹ + 20 g AMF plant⁻¹), B5 (60 g NPK plant⁻¹ + 2 cc biofertilizer l⁻¹ water), B6 (20 g AMF plant⁻¹ + 2 cc biofertilizer l⁻¹ water), and B7 (60 g NPK plant⁻¹ + 20 g AMF plant⁻¹ + 2 cc biofertilizer l⁻¹ water).

Living climbing tree planting

Living tree for pepper climbing used *Pterocarpus indicus* which was obtained from Pangkalpinang City. The selection of living tree climbing was used based on plant height criteria, > 1 m and stem diameter about 3-5 cm. Living climbing tree was planted on the planting hole (40 cm x 40 cm x 40 cm), and about 10 - 15 cm from the planting hole of pepper plant. Living climbing tree was planted about 1 month before pepper was planted.

Pepper planting

Vegetative pepper seedlings (cv. Lampung Daun Lebar) three months old was obtained from Badan Pengelolaan, Pengembangan dan Pemasaran Lada (BP3L), Province of Bangka Belitung. Pepper seedlings was planted in planting hole (40 cm x 40 cm x 40 cm) and 2 m x 2m spacing, amount of 5 kgs of chicken manure and 7 kgs of top soil were applicated on hole planting. The seedlings was planted in soil until the neck seedlings using mixed of soil tailing, top soil, and chicken manure.

Ameliorant application

2 cc l⁻¹ water of biofertilizer was applicated by watered on planting hole four times, i.e. two weeks before pepper planting, one week after planting, two weeks after planting, and three weeks after planting. Mycorrhiza was applicated after planting, and the application of 20 g plant⁻¹ by sown around root area. Inorganic fertilizer (NPK) was applicated twice during planting season. The first application on two weeks after planting (20 g plant⁻¹), and the second application on three weeks after planting (40 g plant⁻¹).

Observation and data analysis

White pepper growth was observed by measuring the height, (cm), diameter of stem (cm), number of leaves, nodus number, and area of leaf (cm²), monthly. Chlorophyll content was observed on the end of observation and using chlorophyll meter. The main effect

was analyzed by the analysis of variance, and countinued by the Duncan's test (α 0.05) to compare among the treatment.

III. RESULTS AND DISCUSSION

Cultivation of pepper in post-tin mining land was very affected by the availability of nutrient for its growth. Post tin-mining land had reported has a poor nutrient, and poor physical and biology properties. This condition requires the fertilizer input to make the favourable environmental for pepper growth. The fertilizer input that can be used like inorganic fertilizer to supply the nutrient directly. Other that, the organic and biofertilizer input also can be added to contribute the nutrient supply by nitrogen fixation process from the air, nor the changing of nutrient available in the soil. Those effort is done to provide the nutrient for pepper growth in marginal land, like in post-tin mining land.

The effort to enhance the marginal land fertility of post-tin mining could be done by addition of soil ameliorant. In this research, the soil ameliorant used was combination of NPK, arbuscular mycorrhiza fungi (AMF), and biofertilizer. The soil ameliorant has significant effect to plant height, segment number, stem diameter, leaves number, chlorophyll content, and leaf area (Table 1). Those indicated that these combination could supply the nutrient and improve the soil physical and biology properties, so that affected the growth characters of pepper.

The differences of each combination was determined by contrast orthogonal test. This analysis showed that combination of NPK, mycorrhiza, and biofertilizer (B7) significantly different to other treatments on all of the characters observed, either to single soil ameliorant or two soil ameliorants. This was caused by directly nutrient could be absorbed by plant from NPK fertilizer, and the ability of mycorrhiza to supply the availability of P, or the its ability to provide the ecosytem to soil microbial which was added by fertilization of organic fertilizer (*Nano Bio*), so that could optimally to suply the nutrient.

Mycorrhiza fungi in ex mining land can play to improve the soil stability [30], by the ability of externals hyfa which plays to initiation the soil macro structure and micro agregation [31]. The formation of soil agregation could be done by the growing of hyfa in soil matrixs to structure shape which occupy the soil particle through physical bond. Hyfa and root create the physical and chemical environmental which would produce organic materials and amorf to tie up the soil particle, and the penetration of hyfa and micro-agregate of root tissue to micro-agregat structure, which increased the capacity and storage the C nutrient, and provide the micro-ecosystem for the soil microbial.

Other that, mycorrhiza colonization also could increase the nutrient uptake by decreasing the distance to the plant nutrient, increasing the nutrient uptake average and the concentration on the surface of absorption, and the change of nutrient shape for absorption to the plant root [32]. The large of mycellium colonization lead the increasing of water uptake by plant, and will promote the plant growth i.e. cell development and elongation by metabolism assimilate. The large of water uptake will lead the N, P, and K nutrient by

NPK fertilization, as well as, N fixed and P dissolved by the organism activity of Nano Bio could be absorbed in large quantity.

The single soil ameliorant application i.e. NPK, AMF, or Nano Bio showed didn't significant different to the combination of two soil ameliorants on the all of characters. Those could be shown by B1 vs B4&B5, B2 vs B4&B6, B3 vs B5&B6 (Table 2). The single soil ameliorant (B1 & B2 & B3) no significantly to the combination of two soil ameliorant on the all of the characters. Those was occurred cause the single soil ameliorant has a the role is not optimally. NPK fertilizer just could supply the nutrient in early fertilization, but not in along plant live, and the mycorrhiza fertilizer requires long time process for associated to root system and its process requires the availability nutrients. While, the organism of biofertilizer couldn't role optimally in unfavourable environmental for the growth of organism and host.

Biofertilizer is a liquid fertilizer and contains nine microbial consortium. Five of those microbial has function in N fixation, i.e. *Azospirillum sp* [33], *Rhizobium sp* [34], *Pseudomonas sp* [35], *Azotobacter sp* [36], and *Alcaligenes sp* [37]. Three of those microbial has function to phosphate solven, i.e. *Aspergillus niger* [38], *Lactobacillus sp* [39], *Bacillus sp* [40], and one of those microbial has role to decomposer, i.e. *Saccharomyces sp* [41].

The differences of among the treatments on the each characters was determined by using Duncan's test. The White pepper applied by ameliorant combination of NPK + mycorrhiza + organic fertilizer showed the best growth on the all of characters observed. This combination has the higher characters, i.e. plant height about 34.08 cm, leaves number about 18.75 blade, segment number about 11.50 segment, stem diameter about 2.45 cm, chlorophyll content about 14.63 mg l⁻¹, and leaf area about 226.48 cm²) than the others combination (Fig. 1).

This combination showed significant different to other combination on the plant height, leaves number, stem diameter, and leaf area characters. Its combination just didn't significant different to biofertilizer + mycorrhiza combination on the leaves number and chlorophyll content characters.

The combination of three soil ameliorant which supply the nutrient from the NPK fertilization will promote the plant growth, and then expand of the root. Those root was colonized by AMF, so that supply the P nutrient and provide the ecosystem for the microbial by improve the soil aggregation and water holding capacity. The increasing of water holding capacity would promote the nutrient absorption by mycorrhiza hyfa colonization, and furthermore these ecosystem would promote the optimally role of organism in N fixation and P solvent. The nutrient and water were absorbed by root, at same time.




Fig. 1. Increasing of plant height, leaves number, segment number, stem diameter, chlorophyll content, and leaf area on different ameliorant combination

TABLE 1. THE EFFECT OF AMELIORANT COMBINATION TO PEPPER GROWTH IN POST-TIN MINING LAND
*significant on F test at $\alpha=0.05$.

Character	F Value	CV (%)
Plant height (cm)	16.82**	13.71
Leaves number	4.32*	14.08
Segment number (segment)	11.65**	11.72
Stem diameter (cm)	11.71**	20.28
Chlorophyll content (cc/g)	3.95*	32.40
Leaf area (cm ²)	6.49*	25.04

The availability of nutrient will promote the plant metabolism activity and distribution of assimilate to the parts of plant. These is indicated by increasing of plant height, leaf number, segment number, stem diameter, chlorophyll content, and leaf area. This increasing is caused by the accumulation of assimilate which promote the cell differentiation and elongation. Mycorrhiza is a symbiotic mutualism between fungi and the rooting system [30]. In marginal land, mycorrhiza can increase the mostly plant growth, whereas not too positive impact on fertility land [42]. Mycorrhiza also increased contact area in soil, so that increased root absorption area till 47 level, and these would promote acces to nutrient in soil [28].

Mycorrhiza application to white pepper in post-tin mining land could increase plant height, leaves number, and branches [6], and could associated with root and shown by root infection about 36.67% [43]. Plant with mycorrhiza infection has higher N, P, K, and Mg absorption than plant without mycorrhiza infection [44]. Nano Bio application of 2 ml l⁻¹ water pursuant to [43] could increase leaves number, stem diameter, and chlorophyll content of pepper in post-tin mining land.

IV. CONCLUSION

1. Soil ameliorant affected to the early growing of pepper in post tin mining area.
2. NPK fertilizer could be substituted by mycorrhiza and organic fertilizer for the early growing of pepper in post tin mining area.
3. The proper ameliorant to the early growing of pepper in post tin mining land was the combination of 60 g NPK, 20 g mycorrhiza, and 2 cc l⁻¹ bio fertilizer.

REFERENCES

[1] Directorate General of Plantation Ministry of Agriculture, "Plantation Indonesian Statistics: Pepper Comodity 2014-2015, Jakarta: Directorate General of Plantation, 2015.

[2] Bappeda Provinsi Bangka Belitung, "Kondisi kerusakan lingkungan hidup di Provinsi Kepulauan Bangka Belitung," unpublished.

[3] Asmarhansyah, R. B. Badayos, P. B. Sanchez, P. C. Sta Cruz, and L. M. Florece, "Land suitability evaluation of abandoned tin-mining areas for Agricultural development in Bangka Island, Indonesia," *J. Degrade. Min. Land Manage*, vol 4, pp. 907-918, July 2017.

[4] Rusli, Y. Ferry, B. Hafif, and E. Wardiana, "The effectiveness of ameliorants, fertilizer, and mycorrhiza for rubber growth at post tin mining land," *J. TIDP*, vol 3, pp. 175-184, November 2016.

[5] I. Inonu, D. Budianta, M.U. Harum, Yakup, and A. Y. A. Wiralaga, "Ameliorasi Organik pada Tailing Pasir Pasca Penambangan Timah untuk Pertumbuhan Bibit Tanaman Karet. *J. Agrotopika*, vol 16, pp. 45-51, January 2011.

[6] Y. Ferry, J. Towaha, and K. D. Sasmita, "Plant water compost usage as a carrier of Mycorrhizal inoculant from Bushy Pepper cultivation in the post-tin mining soil," *J. Litri*, vol 19, pp. 15-23, March 2013.

[7] S. R. P. Sitorus, E. Kusumastuti, and N. Badri, "Post-mining land characteristics and rehabilitation technique in Bangka and Singkep Island," *J. Tanah dan Klim*, vol 27, pp. 57-73, July 2008.

[8] D. K. Majumdar, *Irrigation water management*, India: Prentice Hall of India, 2000, pp. 487.

[9] I. Inonu, N. S. Khodijah, and A. Supriadi, "Pakchoy (*Brassica rapa* L.) cultivation in sandy tailings of tin post-mining site with organic manure and NPK fertilizer ameliorant," *J. Lahan*

Suboptimal, vol 3, pp. 76-82, November 2016.

[10] Y. Ferry, J. Towaha, and K. D. Sasmita, "Perbaikan lahan bekas tambang timah: studi kasus; uji media tanah bekas tambang beberapa macam kompos untuk budidaya lada," *Bul. Ris. Tan. Rempah & Aneka Tan. Indust*, vol 1, pp. 296-308, July 2010.

[11] R. Diaguna, Royalaitani, I. Inonu, and E. Nurtjahya, "Growth of lettuce (*Lactuca sativa* L.) with NPK fertilization in tin post-mining and non distrubed land," *Proceeding of Suboptimal Land 2016*, p. 310-316, October 2016.

[12] Y. Ferry, and J. Towaha, "Effect of composition of N, P and K fertilizer on growth and production of Pepper in after tin m ining soil in Bangka," *Bul. RISTRI*, vol 2, pp. 305-310, October 2011.

[13] Z. F. Fawzy, A. M. El-Bassiony, L. Yunsheng, O. Zhu, and A. A. Ghoname, "Effect of mineral, organic, and bio-N fertilizers on growth, yield, and fruit quality of sweet pepper," *J. Appl. Sci. Res*, vol 8, pp. 3921-3933, August 2012.

[14] M. Zafar, N. Rahim, A. Shaheen, A. Khalik, T. Arjamand, M. Jamil, Z. Rehman, and T. Sultan, "Effect of combining poultry manure, inorganic phosphorous fertilizers, and phosphate solubilizing bacteria on growth, yield, protein content and P uptake in maize," *AAB Bioflux*, vol 3, pp. 47-58, January 2011.

[15] S. Darma, W. Kustiawan, D. Ruhiyat, and Sumaryono, "Organic fertilizers improves trembessi (*Samanea saman*) seedling growth, A case study of the implementation of post-mining land reclamation and revegetation within the forest cultivation zone," *IJSTR*, vol 6, pp. 393-399, October 2017.

[16] S. Widiati, Suliasih, and A. Muharam, "The effect of compost enriched with symbiotic nitrogen fixing and phosphate solubilizing bacteria on the growth of peas and the activity of phosphatase enzymes in the soil," *J. Hort*, vol 20, pp. 207-215, October 2010.

[17] Suliasih, Widawati, and A. Muharam, "The application of organic fertilizers and phosphate solubilizing bacteria to increase the growth of tomato and soil microbial activities," *J. Hort*, vol 20, pp. 241-246, October 2010.

[18] Suwandi, G. A. Sopha, M. P. Yufdi, "The effectiveness of organic fertilizer, NPK, and biofertilizer managements on growth and yields of shallots," *J. Hort*, vol 25, pp. 208-221, September 2015.

[19] O. Benhabib, A. Yazar, M. Qadir, E. Loureno, S. E. Jacobsen, "How can we improve mediterranean cropping systems," *J. Agron. Crop Sci*, vol 200, pp. 325-332, May 2014.

[20] N. Huang, W. Wang, Y. Yao, F. Zhu, W. Wang, and X. Chang, "The influence of different concentrations of bio-organic fertilizer on cucumber *Fusarium wilt* and soil microflora alterations, *PLoS One*, vol 12, pp. e0171490, February 2017.

[21] R. W. Cahyani, and A. K. Hardjana, "Trial of plant species with the treatment of planting media on post C quarry land in KHDTK Labanan, Berau District, East Kalimantan," *Pros. Sem. Nas. Masy. Biodiv. Indon*. vol 3, pp. 361-367, December 2017.

[22] D. Rusmawan, Muzammil, D. Y. Rinawati, Asmarhansyah, "Influence of fertilizing and variety on growth and yield of maize in ex-tin land central Bangka, Kepulauan Bangka Belitung". *Proceeding International Maize Conference*. pp. 209-213

[23] M. Nurcholis, A. Wijayani, and A. Widodo, "Clay and organic matter applications on the coarse quartz tailing material and the sorghum growth on the post mining at Bangka Island," *J. Degrade. Mining Land Manage*, vol 1, pp. 27-32, October 2013.

[24] H. Sukiman, "The use of mycorrhizae for improving the quality of seedling and land productivity in the city area," *Pros. Sem. Nas. Masy. Biodiv. Indon*, vol 1, pp. 2021-2026, September 2015.

[25] M. A. Abiala, O. O. Popoola, O. J. Olawuyi, J. O. Oyelude, A. O. Akanmu, A. S. Killani, O. Osonubi, and A. C. Odebode, "Harnessing the potentials of Vesicular Arbuscular Mycorrhizal (VAM) fungi to plant growth – A review," *Int. J. Pure Appl. Sci. Technol*, vol 14, pp. 61–79, December 2013.

[26] Y. Ferry, and Rusli, "The effects of Mychorryzal and NPK fertilization dosage for growth and production of *Robusta Coffe* among productive coconut plantation," *J. Pen. Tan. Indust*, vol 20, pp. 27–34, February 2014.

[27] Subaedah, "Pemanfaatan jamur mikoriza dalam meningkatkan ketersediaan hara fosfat dan pengaruhnya terhadap pertumbuhan bibit jarak pagar," *J. Agrivigor*, vol 6, pp. 174-177, September 2007.

[28] B. Hafif, S. Sabiham, I. Anas, A. Sutandi, and Suyamto, "Impact of brachiaria, arbuscular mycorrhiza, and potassium enriched rice

- straw compost on aluminum, potassium and stability of acid soil aggregates," *Indon. J. Agric. Sci.* vol 13, pp. 27–34, April 2012.
- [29] B. Bastari, The growth respons of single segment cutting of pepper to Mycorrhiza and organic materials application in some the growth medium of tailing compound, ID: Bangka Belitung University, 2010.
- [30] Suharno, and R. P. Sancayaningsih, "Arbuscular Mycorrhiza Fungi: The potential use of heavy metal mycorrhizo-remediation technology in mined field rehabilitation," *Bioteknologi*, vol 10, pp. 23-34, May 2013.
- [31] E. Orlowska, D. Orlowski, J. Mesjasz-Przybylowicz, and K. Turmau, "Role of mycorrhizal colonization in plant establishment on an alkaline gold mine tailing," *Int. J. Phytoremed.* vol 13, pp. 185-205, February 2011.
- [32] Masria, "Peranan mikoriza vesikular arbuskular (MVA) untuk meningkatkan resistensi tanaman terhadap cekaman kekeringan dan ketersediaan P pada lahan kering," *Partner*, vol 15, pp. 48-56, February 2015.
- [33] S. Widiawati, and A. Muharam, "The laboratory test of *Azospirillum* sp. isolated from several ecosystems," *J. Hort.* vol 22, pp. 250-267, January 2012.
- [34] Novriani, "Peranan *Rhizobium* dalam meningkatkan ketersediaan nitrogen bagi tanaman kedelai," *AgronobiS*, vol 3, pp. 35-42, March 2011.
- [35] H. K. Sejati, M. Astiningrum, and Tujiyanta, "Pengaruh macam pupuk kandang dan konsentrasi *Pseudomonas fluorescens* pada hasil tanaman bawang merah (*Allium cepa* fa. *Ascalonicum* L.) varietas rok kuning," *VIGOR J. Ilmu Pertanian Tropika dan Subtropika*, vol 2, pp. 55-59, March 2017.
- [36] I. Widiyawati, Sugiyanta, A. Junaedi, and R. Widyastuti, "The role of nitrogen-fixing bacteria to reduce the rate of inorganic nitrogen fertilizer on lowland rice," *J. Agron. Indonesia*, vol 42, pp. 96-102, January 2014.
- [37] S. Otte, J. Schalk, J. G. Kuenen, and M. S. Jetten, "Hydroxylamine oxidation and subsequent nitrous oxide production by the heterotrophic ammonia oxidizer *Alcaligenes faecalis*," *Applied Microbiology and Biotechnology*, vol 51, pp. 255-261, February 1999.
- [38] I. J. Artha, H. Guchi, and P. Marbun, "The effectivity of *Aspergillus niger* and *Penicillium* sp to increase the phosphate availability and the growth of corn in andisol soil," *J. Online Agroekoteknologi*, vol 1, pp. 1277-1287, September 2013.
- [39] D. Sudiarti, "The effectiveness of biofertilizer on plant growth soybean "Edamame" (*Glycine max*)," *J. SainHealth*, vol 1, pp. 46-55, September 2017.
- [40] N. Djaenuddin, and A. Muis, Karakteristik bakteri antagonis *Bacillus subtilis* dan potensinya sebagai agens pengendali hayati penyakit tanaman," *Proceeding of National Seminar in Serelia*, pp. 489-494, September 2015
- [41] Y. Rahmah, S. Bahri, and Chairul, "Fermentasi nira nipah menjadi bioetanol menggunakan *Saccharomyces cerevisiae* dengan penambahan urea sebagai sumber nitrogen," *JOM FTEKNIK*, vol 2, pp. 1-5, October 2015.
- [42] S. E. Smith, and D. Read, *Mycorrhizal symbiosis*, 3rd ed., New York: Academic Press – Elsevier. 2008, pp. 800.
- [43] Sumalia, The influence of mychorryza and biofertilizer dosage for Pepper growth in sand tailing medium, ID: Bangka Belitung University, 2017.
- [44] I. Sasli, and A. Ruliansyah, "Pemanfaatan mikoriza arbuskular spesifik lokasi untuk efisiensi pemupukan pada tanaman jagung di lahan gambut tropis," *AGROVIGOR*, vol 5, pp. 65-74, September 2012.

PROSIDING/PROSIDING/Soil Ameliorant in Post Tin Mining Land of Bangka.pdf

By Ismed Inonu

Soil Ameliorant in Post Tin Mining Land of Bangka Island for Pepper Growth

Ismed Inonu
Department of Agrotechnology
University of Bangka Belitung
Pangkalpinang, Indonesia
ismedinonu@yahoo.co.id

Rion Apriyadi
Department of Agrotechnology
University of Bangka Belitung
Pangkalpinang, Indonesia
rion_apriyadi@yahoo.com

Ridwan Diaguna
Department of Agrotechnology
University of Bangka Belitung
Pangkalpinang, Indonesia
ridwandiaguna@gmail.com

Muhamad Rifal Rinaldi
Department of Agrotechnology
University of Bangka Belitung
Pangkalpinang, Indonesia
rifalrinaldi@yahoo.com

Abstract—Pepper is one of the spices plant and its production continuous decrease caused by converting of the planting areas to tin-mining areas. One of the effort that could be done was the usage of sub-optimal land which has low fertility. The research was aimed to study the effect of soil ameliorant application for early growth of pepper plant in the sand tailing area and obtain the proper ameliorant. The research was carried from February to May 2017 in post tin mining land of PT Timah, Dwi Makmur Village, District of Merawang, Bangka Island. The research was arranged in a Randomized Block Design (RBD) single factor dan three replications. The factor used was soil ameliorant combination consisted of seven level i.e. NPK fertilizer, arbuscular mycorrhiza fungi (AMF), biofertilizer, NPK + biofertilizer, NPK + AMF, AMF + organic fertilizer, and NPK + AMF + organic fertilizer. The soil ameliorant applications showed significant effect to plant height, leaves number, segment number, stem diameter, leaf area and chlorophyll content. Complete combination (NPK + AMF + biofertilizer) was the best ameliorant to pepper growth in sand tailing area – post tin mining.

Keywords—ameliorant, AMF, NPK, biofertilizer, pepper, post tin mining land

I. INTRODUCTION

Production of white pepper has continued to decrease in the recent years. It's caused by the decrease of planting area that relate to the high of deforestation by mining activity. The planting area of white pepper decreases of 19.2% in last ten years, from 2004 to 2014, i.e. 201.484 ha in 2004 and 162.751 ha in 2014 [1]

In the future, ex-mining land is one of the biggest challenge of white pepper cultivation. According to [2], Bangka Belitung has post-tin mining land about 390.000 ha, and it's very potential to be utilized. Generally, post-tin mining land is sand tailing site which was dominated by sand fraction about 81% [3], 82% [4], 92% [5], 93% [6] and 96% [7]. Domination of sand tailing leads the low of soil water holding capacity about 5-10% [8], and its capacity just supply about 50-100 mm of the crops water requirement [4]. The soils has characteristics like these, describe the low of fertility, well on physical, chemical, and biology properties.

The utilization of post tin mining land for pepper cultivation requires of adding soil ameliorant. The soil ameliorant is required to improve the soil properties, viz physical, chemical, and biological, furthermore to supply the nutrition, directly. The soil ameliorant that could be used for improving the fertility of post-tin mining land among others; organic fertilizer [9, 10], inorganic fertilizer [11, 12], and mycorrhiza fungi [4].

Organic and biological fertilizers could increase the plant growth [13, 14], increased the soil biology activity [15], and could decreased the use of NPK fertilizer [16, 17, 18]. Furthermore, the adding of organic fertilizer increased the absorption of rainfall, slowed the run-off, decreased the evaporation rate and increased the plant production [19]. Biological agent fertilizers also could increase the physical and chemical soil properties [20].

In the post-tin mining land that has poor nutrient, inorganic fertilizer play a role to supply the nutrient for plant development, directly [21]. The use of Urea, SP-36, and KCl fertilizers could increase the growth and corn yield in the post-tin mining land [22]. Furthermore, application of NPK fertilizer increased the growth of shorgum [23], lettuce [11], and *Samanea saman* [15] which were cultivated in post-tin mining land.

Mycorrhiza is one of the soil fungi, and according to [24], could association with more than 80% types of crop, like forestry, agriculture, horticulture, and plantation. Application of mycorrhiza was required to rehabilitation of soil biology properties and mycorrhiza [25, 26]. Moreover, mycorrhiza could increases the ability of water uptake and nutrient [27]. The use of mycorrhiza as a soil ameliorant in post-tin mining land can improve the soil physical properties [28], chemical and biology properties [29]. Mycorrhiza could increase the rubber growth in post-tin mining land [4].

The adding of three materials as soil ameliorant is expected could increase the fertility of post-tin mining land. The increasing of fertility will promote the white pepper growth in the land. To date, the type of soil ameliorant that could promote the white pepper growth in post-tin mining land hasn't been informed. The research was aimed to study

the soil ameliorant effect for early growing of white pepper in the sand tailing area and obtain the proper ameliorant.

II. METHODS

Schedule and site

The research was carried out from February to May 2017 in post-tin mining land of PT TIMAH Tbk which was located in Merawang, Bangka. The land used is 10 years old after post mining.

Experiment design

A Randomized Block Design (RBD) with single factor and three replications was applied in the research. The factor was soil ameliorant combination, i.e nitrogen-phosphor-kalium (NPK) fertilizer 'Mutiarā', arbuscular mycorrhiza fungi (AMF) genus *Glomus* in zeolit carrier medium, and biofertilizer (*Nano Bio*). The combination consisted of seven levels i.e. B1 (60 g NPK plant⁻¹), B2 (20 g AMF plant⁻¹), B3 (2 cc biofertilizer l⁻¹ water), B4 (60 g NPK plant⁻¹ + 20 g AMF plant⁻¹), B5 (60 g NPK plant⁻¹ + 2 cc biofertilizer l⁻¹ water), B6 (20 g AMF plant⁻¹ + 2 cc biofertilizer l⁻¹ water), and B7 (60 g NPK plant⁻¹ + 20 g AMF plant⁻¹ + 2 cc biofertilizer l⁻¹ water).

Living climbing tree planting

Living tree for pepper climbing used *Pterocarpus indicus* which was obtained from Pangkalpinang City. The selection of living tree climbing was used based on plant height criteria, > 1 m and stem diameter about 3-5 cm. Living climbing tree was planted on the planting hole (40 cm x 40 cm x 40 cm), and about 10 - 15 cm from the planting hole of pepper plant. Living climbing tree was planted about 1 month before pepper was planted.

Pepper planting

Vegetative pepper seedlings (cv. Lampung Daun Lebar) three months old was obtained from Badan Pengelolaan, Pengembangan dan Pemasaran Lada (BP3L), Province of Bangka Belitung. Pepper seedlings was planted in planting hole (40 cm x 40 cm x 40 cm) and 2 m x 2m spacing, amount of 5 kgs of chicken manure and 7 kgs of top soil were applied on hole planting. The seedlings was planted in soil until the neck seedlings using mixed of soil tailing, top soil, and chicken manure.

Ameliorant application

2 cc l⁻¹ water of biofertilizer was applied by watered on planting hole four times, i.e. two weeks before pepper planting, one week after planting, two weeks after planting, and three weeks after planting. Mycorrhiza was applied after planting, and the application of 20 g plant⁻¹ by sown around root area. Inorganic fertilizer (NPK) was applied twice during planting season. The first application on two weeks after planting (20 g plant⁻¹), and the second application on three weeks after planting (40 g plant⁻¹).

Observation and data analysis

White pepper growth was observed by measuring the height, (cm), diameter of stem (cm), number of leaves, nodus number, and area of leaf (cm²), monthly. Chlorophyll content was observed on the end of observation and using chlorophyll meter. The main effect

was analyzed by the analysis of variance, and continued by the Duncan's test (α 0.05) to compare among the treatment.

III. RESULTS AND DISCUSSION

Cultivation of pepper in post-tin mining land was very affected by the availability of nutrient for its growth. Post tin-mining land had reported has a poor nutrient, and poor physical and biology properties. This condition requires the fertilizer input to make the favourable environmental for pepper growth. The fertilizer input that can be used like inorganic fertilizer to supply the nutrient directly. Other that, the organic and biofertilizer input also can be added to contribute the nutrient supply by nitrogen fixation process from the air, nor the changing of nutrient available in the soil. The effort is done to provide the nutrient for pepper growth in marginal land, like in post-tin mining land.

The effort to enhance the marginal land fertility of post-tin mining could be done by addition of soil ameliorant. In this research, the soil ameliorant used was combination of NPK, arbuscular mycorrhiza fungi (AMF), and biofertilizer. The soil ameliorant has significant effect to plant height, segment number, stem diameter, leaves number, chlorophyll content, and leaf area (Table 1). Those indicated that these combination could supply the nutrient and improve the soil physical and biology properties, so that affected the growth characters of pepper.

The differences of each combination was determined by contrast orthogonal test. This analysis showed that combination of NPK, mycorrhiza, and biofertilizer (B7) significantly different to other treatments on all of the characters observed, either to single soil ameliorant or two soil ameliorants. This was caused by directly nutrient could be absorbed by plant from NPK fertilizer, and the ability of mycorrhiza to supply the availability of P, or the its ability to provide the ecosystem to soil microbial which was added by fertilization of organic fertilizer (*Nano Bio*), so that could optimally to supply the nutrient.

Mycorrhiza fungi in ex mining land can play to improve the soil stability [30], by the ability of external hypha which plays to initiation the soil macro structure and micro aggregation [31]. The formation of soil aggregation could be done by the growing of hypha in soil matrix to structure shape which occupy the soil particle through physical bond. Hypha and root create the physical and chemical environmental which would produce organic materials and amorf to tie up the soil particle, and the penetration of hypha and micro-aggregate of root tissue to micro-agregat structure, which increased the capacity and storage the C nutrient, and provide the mycro-ecosystem for the soil microbial.

Other that, mycorrhiza colonization also could increase the nutrient uptake by decreasing the distance to the plant nutrient, increasing the nutrient uptake average and the concentration on the surface of absorption, and the change of nutrient shape for absorption to the plant root [32]. The large of mycellium colonization lead the increasing of water uptake by plant, and will promote the plant growth i.e. cell development and elongation by metabolism assimilate. The large of water uptake will lead the N, P, and K nutrient by

NPK fertilization, as well as, N fixed and P dissolved by the organism activity of Nano Bio could be absorbed in large quantity.

The single soil ameliorant application i.e. NPK, AMF, or Nano Bio showed didn't significant different to the combination of two soil ameliorants on the all of characters. Those could be shown by B1 vs B4&B5, B2 vs B4&B6, B3 vs B5&B6 (Table 2). The single soil ameliorant (B1 & B2 & B3) no significantly to the combination of two soil ameliorant on the all of the characters. Those was occurred cause the single soil ameliorant has a the role is not optimally. NPK fertilizer just could supply the nutrient in early fertilization, but not in along plant live, and the mycorrhiza fertilizer requires long time process for associated to root system and its process requires the availability nutrients. While, the organism of biofertilizer couldn't role optimally in unfavourable environmental for the growth of organism and host.

Biofertilizer is a liquid fertilizer and contains nine microbial consortium. Five of those microbial has function in N fixation, i.e. *Azospirillum sp* [33], *Rhizobium sp* [34], *Pseudomonas sp* [35], *Azotobacter sp* [36], and *Alcaligenes sp* [37]. Three of those microbial has function to phosphate solven, i.e. *Aspergillus niger* [38], *Lactobacillus sp* [39], *Bacillus sp* [40], and one of those microbial has role to decomposer, i.e. *Saccharomyces sp* [41].

The differences of among the treatments on the each characters was determined by using Duncan's test. The White pepper applied by ameliorant combination of NPK + mycorrhiza + organic fertilizer showed the best growth on the all of characters observed. This combination has the higher characters, i.e. plant height about 34.08 cm, leaves number about 18.75 blade, segment number about 11.50 segment, stem diameter about 2.45 cm, chlorophyll content about 14.63 mg l⁻¹, and leaf area about 226.48 cm²) than the others combination (Fig. 1).

This combination showed significant different to other combination on the plant height, leaves number, stem diameter, and leaf area characters. Its combination just didn't significant different to biofertilizer + mycorrhiza combination on the leaves number and chlorophyll content characters.

The combination of three soil ameliorant which supply the nutrient from the NPK fertilization will promote the plant growth, and then expand of the root. Those root was colonized by AMF, so that supply the P nutrient and provide the ecosystem for the microbial by improve the soil agregation and water holding capacity. The increasing of water holding capacity would promote the nutrient absorption by mycorrhiza hyfa colonization, and furthermore these ecosystem would promote the optimally role of organism in N fixation and P solvent. The nutrient and water were absorbed by root, at same time.




Fig. 1. Increasing of plant height, leaves number, segment number, stem diameter, chlorophyll content, and leaf area on different ameliorant combination

TABLE 1. THE EFFECT OF AMELIORANT COMBINATION TO PEPPER GROWTH IN POST-TIN MINING LAND
*significant on F test at $\alpha=0.05$.

Character	F Value	CV (%)
Plant height (cm)	16.82**	13.71
Leaves number	4.32*	14.08
Segment number (segment)	11.65**	11.72
Stem diameter (cm)	11.71**	20.28
Chlorophyll content (cc/g)	3.95*	32.40
Leaf area (cm ²)	6.49*	25.04

The availability of nutrient will promote the plant metabolism activity and distribution of assimilate to the parts of plant. These is indicated by increasing of plant height, leaf number, segment number, stem diameter, chlorophyll content, and leaf area. This increasing is caused by the accumulation of assimilate which promote the cell differentiation and elongation. Mycorrhiza is a symbiotic mutualism between fungi and the rooting system [30]. In marginal land, mycorrhiza can increase the mostly plant growth, whereas not too positive impact on fertility land [42]. Mycorrhiza also increased contact area in soil, so that increased root absorption area till 47 level, and these would promote access to nutrient in soil [28].

Mycorrhiza application to white pepper in post-tin mining land could increase plant height, leaves number, and branches [6], and could associated with root and shown by root infection about 36.67% [43]. Plant with mycorrhiza infection has higher N, P, K, and Mg absorption than plant without mycorrhiza infection [44]. Nano Bio application of 2 ml l⁻¹ water pursuant to [43] could increase leaves number, stem diameter, and chlorophyll content of pepper in post-tin mining land.

IV. CONCLUSION

1. Soil ameliorant affected to the early growing of pepper in post tin mining area.
2. NPK fertilizer could be substituted by mycorrhiza and organic fertilizer for the early growing of pepper in post tin mining area.
3. The proper ameliorant to the early growing of pepper in post tin mining land was the combination of 60 g NPK, 20 g mycorrhiza, and 2 cc l⁻¹ bio fertilizer.

REFERENCES

- [1] Directorate General of Plantation Ministry of Agriculture, "Plantation Indonesian Statistics: Pepper Comodity 2014-2015, Jakarta: Directorate General of Plantation, 2015.
- [2] Bappeda Provinsi Bangka Belitung, Kondisi kerusakan lingkungan hidup di Provinsi Kepulauan Bangka Belitung," unpublished.
- [3] Asmarhansyah, R. B. Badayos, P. B. Sanchez, P. C. Sta Cruz, and L. M. Florece, "Land suitability evaluation of abandoned tin-mining areas for Agricultural development in Bangka Island, Indonesia," *J. Degrade. Min. Land Manage*, vol 4, pp. 907-918, July 2017.
- [4] Rusli, Y. Ferry, B. Hafif, and E. Wardiana, "The effectiveness of ameliorants, fertilizer, and mycorrhiza for rubber growth at post tin mining land," *J. TIDP*, vol 3, pp. 175-184, November 2016.
- [5] I. Inonu, D. Budianta, M.U. Harum, Yakup, and A. Y. A. Wiralaga, "Ameliorasi Organik pada Tailing Pasir Pasca Penambangan Timah untuk Pertumbuhan Bibit Tanaman Karet. *J. Agrotopika*, vol 16, pp. 45-51, January 2011.
- [6] Y. Ferry, J. Towaha, and K. D. Sasmita, "Plant water compost usage as a carrier of Mycorrhizal inoculant from Bushy Pepper cultivation in the post-tin mining soil," *J. Litri*, vol 19, pp. 15-23, March 2013.
- [7] S. R. P. Sitorus, E. Kusumastuti, and N. Badri, "Post-mining land characteristics and rehabilitation technique in Bangka and Singkep Island," *J. Tanah dan Iklim*, vol 27, pp. 57-73, July 2008.
- [8] D. K. Majumdar, Irrigation water management, India: Prentice Hall of India, 2000, pp. 487.
- [9] I. Inonu, N. S. Khodijah, and A. Supriadi, "Pakchoy (*Brassica rapa* L.) cultivation in sandy tailings of tin post-mining site with organic manure and NPK fertilizer ameliorant," *J. Lahan*

- [10] *Suboptimal*, vol 3, pp. 76-82, November 2016.
- [10] Y. Ferry, J. Towaha, and K. D. Sasmita, "Perbaikan lahan bekas tambang timah: studi kasus; uji media tanah bekas tambang dengan beberapa macam kompos untuk budidaya lada," *Bul. Ris. Tan. Rempah & Aneka Tan. Indust*, vol 1, pp. 296-308, July 2010.
- [11] R. Diaguna, Royalaitani, I. Inonu, and E. Nurtjahya, "Growth of lettuce (*Lactuca sativa* L.) with NPK fertilization in tin post-mining and non disturbed land," *Proceeding of Suboptimal Land 2016*, p. 310-316, October 2016.
- [12] Y. Ferry, and J. Towaha, "Effect of composition of N, P and K fertilizer on growth and production of Pepper in after tin mining soil in Bangka," *Bul. RISTRI*, vol 2, pp. 305-310, October 2011.
- [13] Z. F. Fawzy, A. M. El-Bassiony, L. Yunsheng, O. Zhu, and A. A. Ghoname, "Effect of mineral, organic, and bio-N fertilizers on growth, yield, and fruit quality of sweet pepper," *J. Appl. Sci. Res*, vol 8, pp. 3921-3933, August 2012.
- [14] M. Zafar, N. Rahim, A. Shaheen, A. Khalik, T. Arjamand, M. Jamil, Z. Rehman, and T. Sultan, "Effect of combining poultry manure, inorganic phosphorous fertilizers, and phosphate solubilizing bacteria on growth, yield, protein content and P uptake in maize," *AAB Bioflux*, vol 3, pp. 47-58, January 2011.
- [15] S. Darma, W. Kustiawan, D. Ruhayat, and Sumaryono, "Organic fertilizers improves trembessi (*Samanea saman*) seedling growth. A case study of the implementation of post-mining land reclamation and revegetation within the forest cultivation zone," *IJSTR*, vol 6, pp. 393-399, October 2017.
- [16] S. Widiati, Suliasih, and A. Muharam, "The effect of compost enriched with symbiotic nitrogen fixing and phosphate solubilizing bacteria on the growth of peas and the activity of phosphatase enzymes in the soil," *J. Hort*, vol 20, pp. 207-215, October 2010.
- [17] Suliasih, Widawati, and A. Muharam, "The application of organic fertilizers and phosphate solubilizing bacteria to increase the growth of tomato and soil microbial activities," *J. Hort*, vol 20, pp. 241-246, October 2010.
- [18] Suwandi, G. A. Sopha, M. P. Yufli, "The effectiveness of organic fertilizer, NPK, and biofertilizer managements on growth and yields of shallots," *J. Hort*, vol 25, pp. 208-221, September 2015.
- [19] O. Benhabib, A. Yazar, M. Qadir, E. Loureno, S. E. Jacobsen, "How can we improve mediterranean cropping systems," *J. Agron. Crop Sci*, vol 200, pp. 325-332, May 2014.
- [20] N. Huang, W. Wang, Y. Yao, F. Zhu, W. Wang, and X. Chang, "The influence of different concentrations of bio-organic fertilizer on cucumber *Fusarium wilt* and soil microflora alterations," *PLoS One*, vol 12, pp. e0171490, February 2017.
- [21] R. W. Cahyani, and A. K. Hardjana, "Trial of plant species with the treatment of planting media on post C quarry land in KHDTK Labanan, Berau District, East Kalimantan," *Pros. Sem. Nas. Masy. Biodiv. Indon*, vol 3, pp. 361-367, December 2017.
- [22] D. Rusmawan, Muzammil, D. Y. Rinawati, Asmarhansyah, "Influence of fertilizing and variety on growth and yield of maize in ex-tin land central Bangka, Kepulauan Bangka Belitung". *Proceeding International Maize Conference*. pp. 209-213
- [23] M. Nurcholis, A. Wijayani, and A. Widodo, "Clay and organic matter applications on the coarse quartz tailing material and the sorghum growth on the post mining at Bangka Island," *J. Degrade. Mining Land Manage*, vol 1, pp. 27-32, October 2013.
- [24] H. Sukiman, "The use of mycorrhizae for improving the quality of seedling and land productivity in the city area," *Pros. Sem. Nas. Masy. Biodiv. Indon*, vol 1, pp. 2021-2026, September 2015.
- [25] M. A. Abiala, O. O. Popoola, O. J. Olawuyi, J. O. Oyelude, A. O. Akanmu, A. S. Killani, O. Osonubi, and A. C. Odebode, "Harnessing the potentials of Vesicular Arbuscular Mycorrhizal (VAM) fungi to plant growth – A review," *Int. J. Pure Appl. Sci. Technol*, vol 14, pp. 61-79, December 2013.
- [26] Y. Ferry, and Rusli, "The effects of Mychorryzal and NPK fertilization dosage for growth and production of *Robusta Coffe* among productive coconut plantation," *J. Pen. Tan. Indust*, vol 20, pp. 27-34, February 2014.
- [27] Subaedah, "Pemanfaatan jamur mikoriza dalam meningkatkan ketersediaan hara fosfat dan pengaruhnya terhadap pertumbuhan bibit jarak pagar," *J. Agrivigor*, vol 6, pp. 174-177, September 2007.
- [28] B. Hafif, S. Sabiham, I. Anas, A. Sutandi, and Suyamto, "Impact of brachiaria, arbuscular mycorrhiza, and potassium enriched rice

- straw compost on aluminum, potassium and stability of acid soil aggregates," *Indon. J. Agric. Sci.* vol 13, pp. 27–34, April 2012.
- [29] B. Bastari, The growth respons of single segment cutting of pepper to Mycorrhiza and organic materials application in some the growth medium of tailing compound, ID: Bangka Belitung University, 2010.
- [30] Suharno, and R. P. Sancayaningsih, "Arbuscular Mycorrhiza Fungi: The potential use of heavy metal mycorrhizo-remediation technology in mined field rehabilitation," *Bioteknologi*, vol 10, pp. 23-34, May 2013.
- [31] E. Orłowska, D. Orłowski, J. Mesjasz-Przybyłowicz, and K. Tuma, "Role of mycorrhizal colonization in plant establishment on an alkaline gold mine tailing," *Int. J. Phytoremed.* vol 13, pp. 185-205, February 2011.
- [32] Masria, "Peranan mikoriza vesikular arbuskular (MVA) untuk meningkatkan resistensi tanaman terhadap cekaman kekeringan dan ketersediaan P pada lahan kering," *Partner*, vol 15, pp. 48-56, February 2015.
- [33] S. Widiawati, and A. Muharam, "The laboratory test of *Azospirillum* sp. isolated from several ecosystems," *J. Hort.* vol 22, pp. 250-267, January 2012.
- [34] Novriani, "Peranan *Rhizobium* dalam meningkatkan ketersediaan nitrogen bagi tanaman kedelai," *AgronomiS*, vol 3, pp. 35-42, March 2011.
- [35] H. K. Sejati, M. Astiningrum, and Tujiyanta, "Pengaruh macam pupuk kandang dan konsentrasi *Pseudomonas fluorescens* pada hasil tanaman bawang merah (*Allium cepa* fa. *Ascalonicum* L.) varietas rok kuning," *VIGOR J. Ilmu Pertanian Tropika dan Subtropika*, vol 2, pp. 55-59, March 2017.
- [36] I. Widiawati, Sugiyanta, A. Junaedi, and R. Widyastuti, "The role of nitrogen-fixing bacteria to reduce the rate of inorganic nitrogen fertilizer on lowland rice," *J. Agron. Indonesia*, vol 42, pp. 96-102, January 2014.
- [37] S. Otte, J. Schalk, J. G. Kuenen, and M. S. Jetten, "Hydroxylamine oxidation and subsequent nitrous oxide production by the heterotrophic ammonia oxidizer *Alcaligenes faecalis*," *Applied Microbiology and Biotechnology*, vol 51, pp. 255-261, February 1999.
- [38] I. J. Artha, H. Guchi, and P. Marbun, "The effectivity of *Aspergillus niger* and *Penicillium sp* to increase the phosphate availability and the growth of corn in andisol soil," *J. Online Agroekoteknologi*, vol 1, pp. 1277-1287, September 2013.
- [39] D. Sudiarti, "The effectiveness of biofertilizer on plant growth soybean "Edamame" (*Glycine max*)," *J. SainHealth*, vol 1, pp. 46-55, September 2017.
- [40] N. Djaenuddin, and A. Muis, Karakteristik bakteri antagonis *Bacillus subtilis* dan potensinya sebagai agens pengendali hayati penyakit tanaman," *Proceeding of National Seminar in Serelia*, pp. 489-494, September 2015
- [41] Y. Rahmah, S. Bahri, and Chairul, "Fermentasi nira nipah menjadi bioetanol menggunakan *Saccharomyces cerevisiae* dengan penambahan urea sebagai sumber nitrogen," *JOM FTEKNIK*, vol 2, pp. 1-5, October 2015.
- [42] S. E. Smith, and D. Read, *Mycorrhizal symbiosis*, 3rd ed., New York: Academic Press – Elsevier. 2008, pp. 800.
- [43] Sumalia, The influence of mychorryza and biofertilizer dosage for Pepper growth in sand tailing medium, ID: Bangka Belitung University, 2017.
- [44] I. Sasli, and A. Ruliansyah, "Pemanfaatan mikoriza arbuskular spesifik lokasi untuk efisiensi pemupukan pada tanaman jagung di lahan gambut tropis," *AGROVIGOR*, vol 5, pp. 65-74, September 2012.

PROSIDING/PROSIDING/Soil Ameliorant in Post Tin Mining Land of Bangka.pdf

ORIGINALITY REPORT

6%

SIMILARITY INDEX

PRIMARY SOURCES

- 1** T Lestari, R Apriyadi, Hartina. " Optimization of maize (L) cultivation in post tin mining land ", IOP Conference Series: Earth and Environmental Science, 2020 154 words — **6%**
Crossref

EXCLUDE QUOTES ON

EXCLUDE MATCHES < 3%

EXCLUDE BIBLIOGRAPHY ON

**LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEERREVIEW
KARYA ILMIAH: PROSIDING***

Judul Jurnal Ilmiah (Artikel) : Soil Ameliorant in Post Tin Mining Land of Bangka Island for Pepper Growth

Jumlah Penulis : 4

Status Pengusul : Penulis Pertama

Identitas Jurnal Ilmiah : 8. Nama Prosiding : International Conference on Maritime and Archipelago (ICoMA 2018)

9. Nomor E-ISSN : 2352-5401

10. Volume, nomor, bulan, tahun : Vol. 167, Oktober 2019

11. Penerbit : Atlantis Press.

12. DOI artikel (jika ada) : <https://doi.org/10.2991/icoma-18.2019.23>

13. Alamat web Prosiding : <https://www.atlantis-press.com>

14. Indeks : Google Scholar

Kategori Publikasi Prosiding (berdasarkan kategori yang tepat) :

	Internasional terindeksasi Scimagojr dan Scopus
	Internasional terindeksasi Scopus, IEEE Explore, SPIE
	Internasional

II. Hasil Penilaian Validasi

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	Similarity index= 6%
2.	Linearitas	Isi tulisan sesuai dengan bidang ilmu pengusul

II. Hasil Penilaian PeerReview:

Komponen Yang Dinilai5)	Nilai Maksimal Jurnal Ilmiah.....6)			Nilai Akhir Yang Diperoleh7)
	Internasional terindeksasi Scimagojr dan Scopus	Internasional terindeksasi Scopus, IEEE Explore, SPIE	Internasional	
e. Kelengkapan unsure isi Prosiding (10%)	<input style="width: 30px; height: 15px;" type="text"/>	<input style="width: 30px; height: 15px;" type="text"/>	1,5	1,5
h. Ruang lingkup dan kedalaman pembahasan (30%)			4,5	4,5
i. Kecukupan dan kemutahiran data/informasi dan metodologi (30%)			4,5	4,5
j. Kelengkapan unsure dan kualitas penerbit (30%)			4,5	4,5
Total=(100%)			15	15
Nilai Pengusul= 60% x 15= 9				9

KOMENTAR/ULASAN PEERREVIEW	
Kelengkapan unsur isi Prosiding	Kelengkapan unsur dari proseding ini Lengkap, memiliki ISSN dan memiliki yang versi online sehingga dapat ditelusuri , penulis artikel di proseding ini berasal dari banyak negara
Ruang lingkup dan kedalaman pembahasan	Ruang lingkup sesuai dengan metode yang dilakukan dengan pembahasan yang mendalam
Kecukupan dan Kemutakhiran data/informasi dan metodologi	Data dan informasi dalam artikel ini lengkap dan mutakhir yang didukung oleh 85% pustaka terbitan 10 tahun terakhir dan metodologi yang dilakukan dapat memecahkan permasalahan dalam penelitian ini.
Kelengkapan unsur dan kualitas penerbit	Proseding ini lengkap dan kualitas penerbit baik. Proseding internasional ini terindex pada Google Scholar .

Baturaja, 30 Januari 2023
Reviewer 1



Prof. Dr. Ir. Gribaldi, M.Si.
196404151990031002
GB pada Prodi Agroteknologi Fakultas Pertanian Universitas Baturaja

*dinilai oleh dua Reviewer secara terpisah

**coret yang tidak perlu

**LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : PROSIDING***

Judul Jurnal Ilmiah (Artikel) : Soil Ameliorant in Post Tin Mining Land of Bangka Island for Pepper Growth

Jumlah Penulis : 4

Status Pengusul : Penulis Pertama, bukan penulis korespondensi

Identitas Jurnal Ilmiah : 8. Nama Prosiding : International Conference on Maritime and Archipelago (ICoMA 2018) ✓

9. Nomor E-ISSN : 2352-5401

10. Volume, nomor, bulan, tahun : Vol. 167, Oktober 2019

11. Penerbit : Atlantis Press.

12. DOI artikel (jika ada) : <https://doi.org/10.2991/icoma-18.2019.23>

13. Alamat web Prosiding : <https://www.atlantis-press.com>

14. Indeks : Google Scholar ✓

Kategori Publikasi Prosiding (beri pada kategori yang tepat) :

Internasional terindeksasi Scimagojr dan Scopus

Internasional terindeksasi Scopus, IEEE Explore, SPIE

Internasional

II. Hasil Penilaian Validasi

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	Similarity index = 6% ✓
2.	Linearitas	Isi tulisan sesuai dengan bidang ilmu pengusul


II. Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai5)	Nilai Maksimal Jurnal Ilmiah 6)			Nilai Akhir Yang Diperoleh 7)
	Internasional terindeksasi Scimagojr dan Scopus	Internasional terindeksasi Scopus, IEEE Explore, SPIE	Internasional	
	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
a. Kelengkapan unsur isi Prosiding (10%)			1,5	1,5
b. Ruang lingkup dan kedalaman pembahasan (30%)			4,5	4
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)			4,5	4
d. Kelengkapan unsur dan kualitas penerbit (30%)			4,5	4,5

Total = (100%)			15	14
Nilai Pengusul = 60% x =	→	60% x 14 =	8,4	

KOMENTAR/ULASAN PEER REVIEW	
Kelengkapan unsur isi Prosiding	Prosiding sudah memenuhi kriteria, akhir buku
Ruang lingkup dan kedalaman pembahasan	Pembahasan kurang dalam karena tidak menunjukkan gambar tata letak mikurizer
Kecukupan dan kemutakhiran data/informasi dan metodologi	Metodologi sudah sesuai
Kelengkapan unsur dan kualitas penerbit	Kualitas penerbit sudah sesuai

Balunujuk, 2 Februari 2023
Reviewer 2



Dr. Tri Lestari S.P., M.Si.
197607162021212004

Lektor Kepala pada Prodi Agroteknologi Universitas Bangka Belitung

* dinilai oleh dua Reviewer secara terpisah

** coret yang tidak perlu