Ethnozoology of Lanun community towards biodiversity conservation at Tebat Rasau ancient lake geosite in Simpang Renggiang District, East Belitung Regency


by Dosen Akuakultur

Submission date: 08-Dec-2023 01:18PM (UTC+0700)

Submission ID: 2227738284

File name: 2023.2931-2938 artikel.pdf (307.1K)

Word count: 4350 Character count: 24639



Ethnozoology of Lanun community towards biodiversity conservation at Tebat Rasau ancient lake geosite in Simpang Renggiang District, East Belitung Regency

¹Endang Bidayani, ²Devi Valeriani, ¹Fitri S. Valen

Aquaculture Department, Faculty of Agriculture Fisheries and Biology, Universitas Bangka Belitung, Balunijuk, Bangka Belitung, Indonesia; Department of Economics, Faculty of Economics, Universitas Bangka Belitung, Bangka, 33172, Indonesia. Corresponding author: E. Bidayani, endangbidayani@gmail.com.

Abstract. Tebat Rasau ancient lake geosite in Simpang Renggiang District, East Belitung Regency is part of the Belitong UNESCO Global Geopark in 2021. Tebat Rasau Ancient Lake has high biodiversity, and is specifically home to protect primitive species namely ornamental arowana fish (*Scleropages formosus*), ampong fish (*Channa marulius*) and green-spotted puffer fish (*Dichotomyctere nigroviridis*). This research examined the ethnozoology of the Lanun community as an effort to conserve lake resources. The objective of this research is to study the ethnozoology of Lanun community towards conservation efforts of Tebat Rasau ancient lake resources. The research method was a survey with FGD technique with participants of all 13 members of Lanun community, Lintang Village Government, Renggiang Sub-District Head, and East Belitung Regency Research and Developm 1 Agency. The research was conducted in Simpang Renggiang District, East Belitung Regency. Data collection methods in this research included primary data and secondary data. In supporting data collection to calculate the community value of the area, this research used economic resource valuation techniques with questionnaires, interviews, observation and documentation. Secon 2 by data collection was supported by relevant literature studies. This research also applied explorative descriptive data analysis method. The results of the research showed that the Lanun community's efforts to preserve the Tebat Rasau geosite are preserving the local culture of environmentally friendly fishing known as nirok nanggok and bebanjor. This local tradition is expected to prevent the extinction of fish resources in the future.

Key Words: economic value, freshwater fishes, Lintang Village, local culture.

Introduction. The Belitong UNESCO Global Geopark is a part of Bangka Belitung provints that has prominent geological elements, surrounded by Karimata Strait in the north and Gaspar Strait in the west. Its geographical boundaries are 107°13'00"-108°51'00" East longitude and 02°29'30"-03°25'00" South latitude. The Belitong Geopark has a total area of about 4,800 km² of land and 13,000 km² of sea and is surrounded by 241 small islands, including Mendanau, Kalimambang, Gresik and Selu islands (Pranit & Musthofa 2021).

Belitong Global Geopark, designated by UNESCO in 2021, is a home for more than 288,000 people from diverse cultures including the Sawang Tribe, as well as home for plant and animal species endemic to Belitung Island. The unique Tor granite landscape, meteorite impact remnants, mining heritage, and diverse local culture make it a site of significant tourism potential. At the same time, however, its complex ecology requires careful and integrated management to ensure long-term sustainability (en.unesco.org).

Belitong UNESCO Global Geopark tourism is included in the green tourism category. According to Arismayanti (2015), Jumadi & Aditya (2021), and Alvianna et al (2022), green tourism is a form of ecotourism that focuses on sustainable tourism, meaning tourism activities that do not damage tourist sites (environmentally friendly). In addition, Parmawati et al (2020) and Megawati et al (2023) stated that green tourism development strategies include high community involvement, and high participation of

tourists in protecting the environment. According to Zukhri et al (2021), Belitong Geopark is an environment-based tourism.

One of the geosites of the Belitong UNESCO Global Geopark is Tebat Rasau ancient lake (Figure 1). The lake has high biodiversity, and is particularly home to ornamental arowana (*Scleropages formosus*), ampong fish (*Channa marulius*) and greenspotted pufferfish (*Dichotomyctere nigroviridis*) (belitonggeopark.net). According to Djapani et al (2021), freshwater and forest ecosystems in Tebat Rasau, apart from having an important role for the community, can also be a natural laboratory. For generations, local people have had the knowledge to utilize natural resources there, such as how to catch fish and use plants as spices. The Geopark concept helps add the scientific reference that the Lenggang River is the habitat of the primitive species Asian arowana (*S. formosus*) which is an endangered species and is included in the IUCN Red List.



Figure 1. Tebat Rasau ancient lake (Source: babelinsight.id).

The investigation of the utilization of biodiversity, particularly indigenous lake fish, to meet the requirements of local communities has not yet been explored in scholarly research. Hence, the accurate recording of used data has been lacking. It is crucial to ensure the maintenance of protected primitive fish species in order to sustain their conservation (Ndobe et al 2022; Valen et al 2023). Djapani et al (2021) said that the Lanun Tebat Rasau community refers to the Tebat Rasau Geosite Management community, which was established in the early part of 2018. The primary objective of this community is to conserve the Tebat Rasau Rawa and the Lenggang River, which serve as vital sources of freshwater for the local communities residing in the vicinity. The community regularly traverses the Lenggang River and the surrounding Kerangas forest on a daily basis. Subsequently, in the year 2019, the organization transitioned into the Tourism Awareness Group (Pokdarwis) and initiated conservation initiatives aimed at preserving indigenous and native fish species.

This research aims to examine the ethnozoology of the Lanun Community as an effort to conserve lake resources. According to Hasan et al (2022), ethnozoology discovers the knowledge of local people about the use of animals around them. Ethnozoology studies the utilization and management of animal species diversity that is closely related to the culture of society and to an ethnic group (Hassan et al 2022). Moreover, ethnozoology examines the relationship from the past to the present between people and the animals around them (Hunn & Brown 2011).

Sustainable management of lake resources should also consider economic, ecological and social aspects (Hasan et al 2023a). The economic dimension entails augmenting the local community's revenue in the vicinity of Tebat Rasau by means of educational tourism centered on ethnozoology. The ecological dimension involves safeguarding and conserving aquatic biota and the surrounding environment. Lastly, the social dimension encompasses the community's involvement in endeavors aimed at

preserving the resources of Tebat Rasau. Therefore, this research aims to examine the ethnozoology of the Lanun community towards conservation efforts of Tebat Rasau Ancient Lake resources.

This research is expected to give some benefits to: 1) the local government, policy brief as input in the preparation of management policies for the Tebat Rasau ancient lake geosite; 2) the community, providing direction for lake resource conservation efforts; and 3) researchers, as a contribution to sustainable lake resource management.

Material and Method. The research was conducted in Simpang Renggiang District, East Belitung Regency (Figure 1) from April to June 2023. The location selection technique was carried out with purposively sampling, with the consideration that Tebat Rasau ancient lake has protected primitive species, as a resource conservation effort. Belitung Island is approximately 200 km to the east of the nearest locality in Sumatra, and 250 km to the west of the nearest locality in Borneo (Hasan et al 2023a; Hasan et al 2023b).




Figure 1. Location of research in Tebat Rasau, East Belitung Regency (red triangle).

The method used in this research was the survey method, in which the main sources of data and information were obtained from respondents as research samples (Bidayani et al 2023). Data collection was done through focus group discussion (FGD) with participants, namely all the 13 members of the Lanun community, Lintang Village Government, Renggiang Sub-District Head, and the Regional Research and Development Agency of East Belitung Regez cy.

In this research, the primary data collection employed questionnaires, interviews, observation and decumentation. Secondary data collection was supported by relevant literature studies. The data analysis method used is the exploratory descriptive method, which describes the situation that occurs systematically and factually with the aim of describing and solving the research problems.

Results. From the research, it was shown that the types of fish with economic value produced from Tebat Rasau Lake included *Wallago* sp., *Channa marulioides*, *Scleropages formosus* and others. The utilization of fish was for consumption and as ornamental fish. In detail, the types of fish sampled from Tebat Rasau lake are presented in Table 1.

Table 1 Fish species of Tebat Rasau Lake

Tapah (Wallago sp.) 50.000-75.000 kg¹	No	Types of fish	Price (Ph)	
2 Ampong (Channa marulioides) 3 Tupok (Channa lucius) 4 Tengkelesak (Scleropages formosus) 5 Baong (Hemibagrus nemurus) 6 Mengkawak (Channa striata) 7 Mentutu (Oxyeleotris marmorata) 8 Kaca (Gymnochanda filamentosa) 9 Cincang nangka (Eirmotus sp.) 10 Tempalak (Betta sp.) 11 Tilan (Mastacembelus sp.) 12 Temelayar tanggok (Sphaerichtys sp.) 13 Gersik (Nemacheilus sp.) 14 Lelais (Kryptopterus sp.) 15 Bantak (Osteochilus) 16 Keperas (Cyclocheilichthys apogon) 17 Seluang (Bebidis) 18 Ban (Barbodes sp.) 19 Buntal Air Tawar (Pao palembangensis) 10 Kejinding (Pristolepis sp.) 20 Cempedik (Osteochilus spilurus) 21 Kepunyu (Belontia sp.) 22 Cempedik (Osteochilus spilurus) 23 Semupuk (Nandus sp.) 25 Temerucong (Luciocephalus pulcher) 26 Temeliyong (Chaca bankanensis) 27 Lenjing (Macrognathus sp.) 28 Senggarong (Mastacembelus notophthalmus) 29 Kemuring (Striuntius sp.) 35.000 kg⁻¹ 35.000 kg⁻¹ 35.000 head⁻¹ 45.000-50.000 head⁻¹ 45.000-50.000 head⁻¹ 35.000 head⁻¹	No	Types of fish	Price (4p)	
3 Tupok (Channa lucius) 40.000 kg¹¹ 4 Tengkelesak (Scleropages formosus) ≥100.000 kg¹¹ 5 Baong (Hemibagrus nemurus) 50.000 kg¹¹ 6 Mengkawak (Channa striata) 40.000-50.000 kg¹¹ 7 Mentutu (Oxyeleotris marmorata) 70.000 kg¹¹ 8 Kaca (Gymnochanda filamentosa) 2.000 head¹¹ 9 Cincang nangka (Eirmotus sp.) 2.000 head¹¹ 10 Tempalak (Betta sp.) 5.000 head¹¹ 11 Tilan (Mastacembelus sp.) 100.000 head¹¹ 12 Temelayar tanggok (Sphaerichtys sp.) 4.000-5.000 head¹¹ 13 Gersik (Nemacheilus sp.) 1.000-5.000 head¹¹ 14 Lelais (Kryptopterus sp.) 1.500 head¹¹ 15 Bantak (Osteochilus) 50.000 *70.000 kg¹¹ 16 Keperas (Cyclocheilichthys apogon) 35.000 kg¹¹ 17 Seluang (Bebidis) 50.000 kg¹¹ 18 Ban (Barbodes sp.) 35.000 kg¹¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head¹¹ 20 Kelik (Clarias sp) 50.000 kg¹¹ 21 Kepinding (Pristolepis sp.) 50				
4 Tengkelesak (<i>Scleropages formosus</i>) 5 Baong (<i>Hemibagrus nemurus</i>) 6 Mengkawak (<i>Channa striata</i>) 7 Mentutu (<i>Oxyeleotris marmorata</i>) 8 Kaca (<i>Gymnochanda filamentosa</i>) 9 Cincang nangka (<i>Eirmotus</i> sp.) 10 Tempalak (<i>Betta</i> sp.) 11 Tilan (<i>Mastacembelus</i> sp.) 12 Temelayar tanggok (<i>Sphaerichtys</i> sp.) 13 Gersik (<i>Nemacheilus</i> sp.) 14 Lelais (<i>Kryptopterus</i> sp.) 15 Bantak (<i>Osteochilus</i>) 16 Keperas (<i>Cyclocheilichthys apogon</i>) 17 Seluang (<i>Bebidis</i>) 18 Ban (<i>Barbodes</i> sp.) 19 Buntal Air Tawar (<i>Pao palembangensis</i>) 20 Kelik (<i>Clarias</i> sp) 21 Kepinding (<i>Pristolepis</i> sp.) 22 Cempedik (<i>Osteochilus spilurus</i>) 23 Semupuk (<i>Nandus</i> sp.) 24 Kepuyu (<i>Belontia</i> sp.) 25 Temerucong (<i>Luciocephalus pulcher</i>) 26 Temeliyong (<i>Chaca bankanensis</i>) 27 Lenjing (<i>Macrognathus</i> sp.) 28 Senggarong (<i>Mastacembelus notophthalmus</i>) 29 Kemuring (<i>Striuntius</i> sp.) 35.000 kg⁻¹ 35.000 kg⁻¹ 45.000-50.000 kg⁻¹ 35.000 head⁻¹ 35.000 kg⁻¹ 35.000 kg⁻¹ 35.000 kg⁻¹ 35.000 kg⁻¹ 35.000-50.000 kg⁻¹ 35.000 kg⁻¹ 35.000-50.000 kg⁻¹				
5 Baong (Hemibagrus nemurus) 50.000 kg ⁻¹ 6 Mengkawak (Channa striata) 40.000-50.000 kg ⁻¹ 7 Mentutu (Oxyeleotris marmorata) 70.000 kg ⁻¹ 8 Kaca (Gymnochanda filamentosa) 2.000 head ⁻¹ 9 Cincang nangka (Eirmotus sp.) 2.000 head ⁻¹ 10 Tempalak (Betta sp.) 5.000 head ⁻¹ 11 Tilan (Mastacembelus sp.) 100.000 head ⁻¹ 12 Temelayar tanggok (Sphaerichtys sp.) 4.000-5.000 head ⁻¹ 13 Gersik (Nemacheilus sp.) 1.000-5.000 head ⁻¹ 14 Lelais (Kryptopterus sp.) 1.500 head ⁻¹ 15 Bantak (Osteochilus) 50.000-70.000 kg ⁻¹ 16 Keperas (Cyclocheilichthys apogon) 35.000 kg ⁻¹ 17 Seluang (Bebidis) 50.000 kg ⁻¹ 18 Ban (Barbodes sp.) 35.000 kg ⁻¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 kg ⁻¹ 20 Kelik (Clarias sp) 50.000 kg ⁻¹ 21 Kepinding (Pristolepis sp.) 50.000-75.000 kg ⁻¹ 22 Cempedik (Osteochilu				
6 Mengkawak (Channa striata) 40.000-50.000 kg ⁻¹ 7 Mentutu (Oxyeleotris marmorata) 70.000 kg ⁻¹ 8 Kaca (Gymnochanda filamentosa) 2.000 head ⁻¹ 9 Cincang nangka (Eirmotus sp.) 2.000 head ⁻¹ 10 Tempalak (Betta sp.) 5.000 head ⁻¹ 11 Tilan (Mastacembelus sp.) 100.000 head ⁻¹ 12 Temelayar tanggok (Sphaerichtys sp.) 4.000-5.000 head ⁻¹ 13 Gersik (Nemacheilus sp.) 1.500 head ⁻¹ 14 Lelais (Kryptopterus sp.) 1.500 head ⁻¹ 15 Bantak (Osteochilus) 50.000-70.000 kg ⁻¹ 16 Keperas (Cyclocheilichthys apogon) 35.000 kg ⁻¹ 17 Seluang (Bebidis) 50.000 kg ⁻¹ 18 Ban (Barbodes sp.) 35.000 kg ⁻¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head ⁻¹ 20 Kelik (Clarias sp) 50.000 kg ⁻¹ 21 Kepinding (Pristolepis sp.) 40.000-50.000 kg ⁻¹ 22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 <t< td=""><td></td><td></td><td colspan="2"></td></t<>				
7 Mentutu (Oxyeleotris marmorata) 70.000 kg¹¹ 8 Kaca (Gymnochanda filamentosa) 2.000 head¹¹ 9 Cincang nangka (Eirmotus sp.) 2.000 head¹¹ 10 Tempalak (Betta sp.) 5.000 head¹¹ 11 Tilan (Mastacembelus sp.) 100.000 head¹¹ 12 Temelayar tanggok (Sphaerichtys sp.) 4.000-5.000 head¹¹ 13 Gersik (Nemacheilus sp.) 1.000-5.000 head¹¹ 14 Lelais (Kryptopterus sp.) 1.500 head¹¹ 15 Bantak (Osteochilus) 50.000-70.000 kg¹¹ 16 Keperas (Cyclocheilichthys apogon) 35.000 kg¹¹ 17 Seluang (Bebidis) 50.000 kg¹¹ 18 Ban (Barbodes sp.) 35.000 kg¹¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head¹¹ 20 Kelik (Clarias sp) 50.000 kg¹¹ 21 Kepinding (Pristolepis sp.) 40.000 kg¹¹ 22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg¹¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg¹¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg¹¹ 25 Temerucong (Luciocephalus pulcher)				
8 Kaca (Gymnochanda filamentosa) 2.000 head ⁻¹ 9 Cincang nangka (Eirmotus sp.) 2.000 head ⁻¹ 10 Tempalak (Betta sp.) 5.000 head ⁻¹ 11 Tilan (Mastacembelus sp.) 100.000 head ⁻¹ 12 Temelayar tanggok (Sphaerichtys sp.) 4.000-5.000 head ⁻¹ 13 Gersik (Nemacheilus sp.) 1.000-5.000 head ⁻¹ 14 Lelais (Kryptopterus sp.) 1.500 head ⁻¹ 15 Bantak (Osteochilus) 50.000-70.000 kg ⁻¹ 16 Keperas (Cyclocheilichthys apogon) 35.000 kg ⁻¹ 17 Seluang (Bebidis) 50.000 kg ⁻¹ 18 Ban (Barbodes sp.) 35.000 kg ⁻¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head ⁻¹ 20 Kelik (Clarias sp) 50.000 kg ⁻¹ 21 Kepinding (Pristolepis sp.) 40.000 kg ⁻¹ 22 Cempedik (Osteochilus spilurus) 50.000-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (Luciocephalus pulcher) 15.000-50.000 head ⁻¹ 26		,		
9		. ,		
10 Tempalak (Betta sp.) 5.000 head 1 11 Tilan (Mastacembelus sp.) 100.000 head 1 12 Temelayar tanggok (Sphaerichtys sp.) 4.000-5.000 head 1 13 Gersik (Nemacheilus sp.) 1.000-5.000 head 1 14 Lelais (Kryptopterus sp.) 1.500 head 4 15 Bantak (Osteochilus) 50.000-70.000 kg 1 16 Keperas (Cyclocheilichthys apogon) 35.000 kg 1 17 Seluang (Bebidis) 50.000 kg 1 18 Ban (Barbodes sp.) 35.000 kg 1 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head 1 20 Kelik (Clarias sp) 50.000 kg 1 21 Kepinding (Pristolepis sp.) 40.000 kg 1 22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg 1 23 Semupuk (Nandus sp.) 35.000-40.000 kg 1 24 Kepuyu (Belontia sp.) 40.000-50.000 kg 1 25 Temerucong (Luciocephalus pulcher) 15.000 head 1 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head 1 27 Lenjing (Macrognathus sp.) 35.000-50.000 head 1 28 Senggarong (Mastac		Kaca (<i>Gymnochanda filamentosa</i>)		
11 Tilan (Mastacembelus sp.) 100.000 head ¹ 12 Temelayar tanggok (Sphaerichtys sp.) 4.000-5.000 head ¹ 13 Gersik (Nemacheilus sp.) 1.000-5.000 head ¹ 14 Lelais (Kryptopterus sp.) 1.500 head ¹ 15 Bantak (Osteochilus) 50.000-70.000 kg ¹ 16 Keperas (Cyclocheilichthys apogon) 35.000 kg ¹ 17 Seluang (Bebidis) 50.000 kg ¹ 18 Ban (Barbodes sp.) 35.000 kg ¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head ¹ 20 Kelik (Clarias sp) 50.000 kg ¹ 21 Kepinding (Pristolepis sp.) 40.000 kg ¹ 22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg ¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ¹ 25 Temerucong (Luciocephalus pulcher) 15.000 head ¹ 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head ¹ 27 Lenjing (Macrognathus sp.) 15.000-20.000 head ¹ 28 Senggarong (Mastacembelus notophthalmus) 35.000-50.000 kg ¹ 29	9	Cincang nangka (Eirmotus sp.)	2.000 head^{-1}	
12 Temelayar tanggok (Sphaerichtys sp.) 4.000-5.000 head-1 13 Gersik (Nemacheilus sp.) 1.000-5.000 head-1 14 Lelais (Kryptopterus sp.) 1.500 head-2 15 Bantak (Osteochilus) 50.000-70.000 kg-1 16 Keperas (Cyclocheilichthys apogon) 35.000 kg-1 17 Seluang (Bebidis) 50.000 kg-1 18 Ban (Barbodes sp.) 35.000 kg-1 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head-1 20 Kelik (Clarias sp) 50.000 kg-1 21 Kepinding (Pristolepis sp.) 40.000 kg-1 22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg-1 23 Semupuk (Nandus sp.) 35.000-40.000 kg-1 24 Kepuyu (Belontia sp.) 40.000-50.000 kg-1 25 Temerucong (Luciocephalus pulcher) 15.000-50.000 head-1 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head-1 27 Lenjing (Macrognathus sp.) 15.000-20.000 head-1 28 Senggarong (Mastacembelus notophthalmus) 35.000-50.000 kg-1 29 Kemuring (Striuntius sp.) 35.000-50.000 kg-1 <td>10</td> <td>Tempalak (<i>Betta</i> sp.)</td> <td colspan="2">5.000 head⁻¹</td>	10	Tempalak (<i>Betta</i> sp.)	5.000 head ⁻¹	
13	11	Tilan (Mastacembelus sp.)		
14 Lelais (Kryptopterus sp.) 1.500 head [4] 15 Bantak (Osteochilus) 50.000-70.000 kg-1 16 Keperas (Cyclocheilichthys apogon) 35.000 kg-1 17 Seluang (Bebidis) 50.000 kg-1 18 Ban (Barbodes sp.) 35.000 kg-1 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head-1 20 Kelik (Clarias sp) 50.000 kg-1 21 Kepinding (Pristolepis sp.) 40.000 kg-1 22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg-1 23 Semupuk (Nandus sp.) 35.000-40.000 kg-1 24 Kepuyu (Belontia sp.) 40.000-50.000 kg-1 25 Temerucong (Luciocephalus pulcher) 15.000-50.000 head-1 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head-1 27 Lenjing (Macrognathus sp.) 15.000-20.000 head-1 28 Senggarong (Mastacembelus notophthalmus) 35.000-50.000 kg-1 29 Kemuring (Striuntius sp.) 35.000-50.000 kg-1	12	Temelayar tanggok (Sphaerichtys sp.)	4.000-5.000 head ⁻¹	
15 Bantak (Osteochilus) 50.000-70.000 kg ⁻¹ 16 Keperas (Cyclocheilichthys apogon) 35.000 kg ⁻¹ 17 Seluang (Bebidis) 50.000 kg ⁻¹ 18 Ban (Barbodes sp.) 35.000 kg ⁻¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head ⁻¹ 20 Kelik (Clarias sp) 50.000 kg ⁻¹ 21 Kepinding (Pristolepis sp.) 40.000 kg ⁻¹ 22 Cempedik (Osteochilus spilurus) 50.40-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (Luciocephalus pulcher) 15.000-50.000 head ⁻¹ 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head ⁻¹ 27 Lenjing (Macrognathus sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (Mastacembelus notophthalmus) 35.000-50.000 kg ⁻¹ 29 Kemuring (Striuntius sp.) 35.000-50.000 kg ⁻¹	13	Gersik (<i>Nemacheilus</i> sp.)	1.000-5.000 head ⁻¹	
16 Keperas (Cyclocheilichthys apogon) 35.000 kg ⁻¹ 17 Seluang (Bebidis) 50.000 kg ⁻¹ 18 Ban (Barbodes sp.) 35.000 kg ⁻¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head ⁻¹ 20 Kelik (Clarias sp) 50.000 kg ⁻¹ 21 Kepinding (Pristolepis sp.) 40.000 kg ⁻¹ 22 Cempedik (Osteochilus spilurus) 50.40-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (Luciocephalus pulcher) 15.000-50.000 head ⁻¹ 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head ⁻¹ 27 Lenjing (Macrognathus sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (Mastacembelus notophthalmus) 35.000-50.000 kg ⁻¹ 29 Kemuring (Striuntius sp.) 35.000-50.000 kg ⁻¹	14	Lelais (Kryptopterus sp.)	1.500 head 🔽	
17 Seluang (Bebidis) 50.000 kg ⁻¹ 18 Ban (Barbodes sp.) 35.000 kg ⁻¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head ⁻¹ 20 Kelik (Clarias sp) 50.000 kg ⁻¹ 21 Kepinding (Pristolepis sp.) 40.000 kg ⁻¹ 22 Cempedik (Osteochilus spilurus) 50.40-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (Luciocephalus pulcher) 15.000 head ⁻¹ 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head ⁻¹ 27 Lenjing (Macrognathus sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (Mastacembelus notophthalmus) 35.000-50.000 kg ⁻¹ 29 Kemuring (Striuntius sp.) 35.000-50.000 kg ⁻¹	15	Bantak (Osteochilus)	50.000-70.000 kg ⁻¹	
18 Ban (Barbodes sp.) 35.000 kg ⁻¹ 19 Buntal Air Tawar (Pao palembangensis) 20.000-25.000 head ⁻¹ 20 Kelik (Clarias sp) 50.000 kg ⁻¹ 21 Kepinding (Pristolepis sp.) 40.000 kg ⁻¹ 22 Cempedik (Osteochilus spilurus) 50.40-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (Luciocephalus pulcher) 15.000 head ⁻¹ 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head ⁻¹ 27 Lenjing (Macrognathus sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (Mastacembelus notophthalmus) 35.000 head ⁻¹ 29 Kemuring (Striuntius sp.) 35.000-50.000 kg ⁻¹	16	Keperas (Cyclocheilichthys apogon)	35.000 kg ⁻¹	
19 Buntal Air Tawar (<i>Pao palembangensis</i>) 20 Kelik (<i>Clarias</i> sp) 21 Kepinding (<i>Pristolepis</i> sp.) 22 Cempedik (<i>Osteochilus spilurus</i>) 23 Semupuk (<i>Nandus</i> sp.) 24 Kepuyu (<i>Belontia</i> sp.) 25 Temerucong (<i>Luciocephalus pulcher</i>) 26 Temeliyong (<i>Chaca bankanensis</i>) 27 Lenjing (<i>Macrognathus</i> sp.) 28 Senggarong (<i>Mastacembelus notophthalmus</i>) 29 Kemuring (<i>Striuntius</i> sp.) 20.000-25.000 kg ⁻¹ 40.000-55.000 kg ⁻¹ 40.000-50.000 kg ⁻¹ 50.000-50.000 head ⁻¹ 50.000-50.000 kg ⁻¹	17	Seluang (Bebidis)	50.000 kg ⁻¹	
20 Kelik (Clarias sp) 50.000 kg ⁻¹ 21 Kepinding (Pristolepis sp.) 40.000 kg ⁻¹ 22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (Luciocephalus pulcher) 15.000 head ⁻¹ 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head ⁻¹ 27 Lenjing (Macrognathus sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (Mastacembelus notophthalmus) 35.000 head ⁻¹ 29 Kemuring (Striuntius sp.) 35.000-50.000 kg ⁻¹	18	Ban (Barbodes sp.)	35.000 kg ⁻¹	
21 Kepinding (Pristolepis sp.) 40.000 kg ⁻¹ 22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (Luciocephalus pulcher) 15.000 head ⁻¹ 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head ⁻¹ 27 Lenjing (Macrognathus sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (Mastacembelus notophthalmus) 35.000 head ⁻¹ 29 Kemuring (Striuntius sp.) 35.000-50.000 kg ⁻¹	19	Buntal Air Tawar (Pao palembangensis)	20.000-25.000 head ⁻¹	
22 Cempedik (Osteochilus spilurus) 50.400-75.000 kg ⁻¹ 23 Semupuk (Nandus sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (Belontia sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (Luciocephalus pulcher) 15.000 head ⁻¹ 26 Temeliyong (Chaca bankanensis) 45.000-50.000 head ⁻¹ 27 Lenjing (Macrognathus sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (Mastacembelus notophthalmus) 35.000 head ⁻¹ 29 Kemuring (Striuntius sp.) 35.000-50.000 kg ⁻¹	20	Kelik (Clarias sp)	50.000 kg ⁻¹	
23 Semupuk (<i>Nandus</i> sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (<i>Belontia</i> sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (<i>Luciocephalus pulcher</i>) 15.000 head ⁻¹ 26 Temeliyong (<i>Chaca bankanensis</i>) 45.000-50.000 head ⁻¹ 27 Lenjing (<i>Macrognathus</i> sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (<i>Mastacembelus notophthalmus</i>) 35.000 head ⁻¹ 29 Kemuring (<i>Striuntius</i> sp.) 35.000-50.000 kg ⁻¹	21	Kepinding (Pristolepis sp.)	40.000 kg ⁻¹	
23 Semupuk (<i>Nandus</i> sp.) 35.000-40.000 kg ⁻¹ 24 Kepuyu (<i>Belontia</i> sp.) 40.000-50.000 kg ⁻¹ 25 Temerucong (<i>Luciocephalus pulcher</i>) 15.000 head ⁻¹ 26 Temeliyong (<i>Chaca bankanensis</i>) 45.000-50.000 head ⁻¹ 27 Lenjing (<i>Macrognathus</i> sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (<i>Mastacembelus notophthalmus</i>) 35.000 head ⁻¹ 29 Kemuring (<i>Striuntius</i> sp.) 35.000-50.000 kg ⁻¹	22	Cempedik (Osteochilus spilurus)	50.400-75.000 kg ⁻¹	
25 Temerucong (<i>Luciocephalus pulcher</i>) 15.000 head ⁻¹ 26 Temeliyong (<i>Chaca bankanensis</i>) 45.000-50.000 head ⁻¹ 27 Lenjing (<i>Macrognathus</i> sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (<i>Mastacembelus notophthalmus</i>) 35.000 head ⁻¹ 29 Kemuring (<i>Striuntius</i> sp.) 35.000-50.000 kg ⁻¹	23	Semupuk (Nandus sp.)	35.000-40.000 kg ⁻¹	
25 Temerucong (<i>Luciocephalus pulcher</i>) 15.000 head ⁻¹ 26 Temeliyong (<i>Chaca bankanensis</i>) 45.000-50.000 head ⁻¹ 27 Lenjing (<i>Macrognathus</i> sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (<i>Mastacembelus notophthalmus</i>) 35.000 head ⁻¹ 29 Kemuring (<i>Striuntius</i> sp.) 35.000-50.000 kg ⁻¹	24			
26 Temeliyong (<i>Chaca bankanensis</i>) 45.000-50.000 head ⁻¹ 27 Lenjing (<i>Macrognathus</i> sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (<i>Mastacembelus notophthalmus</i>) 35.000 head ⁻¹ 29 Kemuring (<i>Striuntius</i> sp.) 35.000-50.000 kg ⁻¹	25		15.000 head ⁻¹	
27 Lenjing (<i>Macrognathus</i> sp.) 15.000-20.000 head ⁻¹ 28 Senggarong (<i>Mastacembelus notophthalmus</i>) 35.000 head ⁻¹ 29 Kemuring (<i>Striuntius</i> sp.) 35.000-50.000 kg ⁻¹	26	Temeliyong (Chaca bankanensis)	45.000-50.000 head ⁻¹	
28 Senggarong (<i>Mastacembelus notophthalmus</i>) 35.000 head ⁻¹ 29 Kemuring (<i>Striuntius</i> sp.) 35.000-50.000 kg ⁻¹	27	, • ,		
29 Kemuring (<i>Striuntius</i> sp.) 35.000-50.000 kg ⁻¹	28			
		3 ()		

Source: Primary data processed (2023).

Jackfruit minced fish is also the raw material for making crackers, which are sold at Rp 5,000/pack of 50 grams, and sambel ingkung (Fish Floss) at Rp 25,000/pack of 100 grams. Besides fish, Tebat Rasau lake also produces a variety of non-fish food products with economic value, including kelulut honey, sepang tea, gelam leaf tea, pelawan leaf tea, berebat tea, and yellow root tea. Food ingredients and product selling prices are presented in Table 2.

Table 2 Food ingredients and product selling price

No.	Туре	Price (Rp)
1	Kelulut honey (Trigona sp.)	50,000/bottle of 100 mL
2	Sepang tea (Caesalpinia sappan)	10,000/pack of 10 grams
3	Gelam leaf tea (Melaleuca cajuputi)	10,000/pack of 10 grams
4	Contrarian leaf tea (Tristaniopsis merguensis)	10,000/pack of 10 grams
5	Tea berebat (Rubus moluccanus)	10,000/pack of 10 grams
6	Yellow root tea (Xanthorhiza simplicissima)	10,000/pack of 10 grams

Source: Primary data processed (2023).

Tebat Rasau Lake is home to a number of native and endemic fish species, making it an ideal location for educational tourism. The Tebat Rasau geosite offers tourists the opportunity to engage in cultural tourism activities, including the traditional practices of catching the fishes (nanggok, nyirok, and gasing). Additionally, they can indulge in culinary tourism by participating in communal dining experiences known as bedulang. Furthermore, the geosite offers educational tourism opportunities, allowing visitors to learn about the endemic and native fish species that inhabit Tebat Rasau Lake. In addition, tourists originate from several other nations, including America, England, Germany, and France, as well as from the local and domestic population, and they have unrestricted access without any associated fees.

Tebat Rasau Lake serves as a habitat for indigenous fish species, providing a source of sustenance for local communities and contributing to their economic well-being. Additionally, the lake serves as a recreational destination for fishing enthusiasts, while also playing a crucial role in the preservation of cultural traditions and local knowledge. Moreover, Tebat Rasau Lake attracts tourists due to its unique characteristics and offerings. According to Djapani et al (2021), the name Rasau is taken from a type of pandanus plant that grows in swamps and rivers. The Kerangas forest ecosystem around the lake also holds unique flora and herbal medicine potential. This geological heritage and biodiversity shape the culture of the local community, which is evident in the way they capture and manage natural resources in the Tebat Rasau area.

Lanun community was established on January 17, 2018. This community has a chairman, vice chairman, treasurer and members with a total of 13 people. Community members come from the people of Lintang Village. Lanun community has a vision and mission. The vision of the Lanun community is to become an organization that participates in developing the potential of members and community involvement and other stakeholders in identifying developing issues and implementing various new approaches in efforts to preserve regional culture, nature conservation, education and entrepreneurship. In addition, the community's mission is: 1) to be active in the preservation of regional culture; 2) preservation of nature, forest ecosystems and wetlands of the Lenggang River; 3) being active in providing education; and 4) entrepreneurship. The tasks and roles of community members include: guiding educational tours to introduce types of fish, providing culinary tours, and others.

According to Utina (2012), traditional values, attitudes and behaviors with ecological insight in the life order of local communities form the ecological intelligence of a community. This local value, for example, applies to coastal communities, which is quite effective in managing natural resources and efforts to preserve their ecosystems.

The Lanun community's effort to preserve Tebat Rasau geosite is to preserve the local culture of environmentally friendly fishing known as nirok nanggok, and bebanjor. Nirok nanggok is a traditional way for Belitung people to catch freshwater fish and it is usually done in group in the river. This activity that is usually carried out by rural communities, has become a routine after the dry season arrives. This routine is still carried out every year during the long dry season between August and September. The areas which still preserve this culture include Belantu Village, Kembiri Village, Membalong Village, and the southern part of Belitung Island. During the dry season, a lot of rivers and swamps recede, the water cannot flow, and the stagnant river water holds many types of freshwater fish.

The name "nirok nanggok" comes from the name of the tools used for fishing, namely tirok and tanggok. Tirok is a kind of sharp wooden stick, approximately 1 cm in diameter, the length of the stick varies from about 2 to 4 meters, at the base of the stick is mounted a spearhead of metal, usually white iron that is pointed and sharp. While tanggok is a kind of net-like tool made of rattan, small in size and with a curved handle, this tool is used for catching/netting fish. Tanggok also has another form, which uses a net mounted on a rectangular rattan frame.

Nirok nanggok tradition is led by a local water shaman and witnessed by village leaders and the local community. This tradition has the function of fostering cohesiveness and strengthening community compliance because residents only take fish for the needs of serving meals together and also do not carry out large-scale exploitation. Furthermore,

this activity serves as a manifestation of the indigenous knowledge and cultural practices within the Belitung community.

In addition to nirok nanggok, a regional tradition that is preserved to maintain the sustainability of Tebat Rasau geosite is bebanjor (fishing), which is looking for freshwater fish using fishing rods along the Lenggang and Tebat Rasau rivers. Fishermen usually stay in huts around the river that is made from sago leaves as a place to rest while waiting for the banjor (hook) to be eaten by the fish.

According to Djapani et al (2021), the Lanun community is a community group that offers tourism services with the main treats being educational and gastronomy tourism. Tourists can go on an excursion while increasing knowledge of geology, biodiversity and local wisdom. The bedulang meal menu, sepang tea spice drinks, pelawan tea and tanggar coffee are very interesting treats. In addition to tour activities, Tebat Rasau also organizes a Nature School that aims to introduce local children to the richness of nature so that the preservation of Tebat Rasau's nature continues from generation to generation.

Discussion. The local wisdom of the community in utilizing fish resources aims to preserve the environment for future generations. The nirok nanggok and bebanjor traditions that are carried out by the Lanun community are conservation efforts to maintain fish resources in Tebat Rasau lake so that they are sustainable. This is in accordance with the opinion of Saputri & Setiawan (2020), that the self-defense process in the form of local wisdom is a form of manifestation and the role of the community in efforts to protect areas that are useful for the lives of rural communities. According to Juniarta et al (2013), the community's local wisdom-based management model has several cultures that have the potential as local wisdom which can later be used as a basis for community empowerment in environmental management.

According to Pinto (2015), coastal and marine resources are currently experiencing degradation as a result of utilization behavior that is not environmentally friendly. Utilization tends to be destructive and damaging and does not consider aspects of conservation and sustainability of resources (Jerikho et al 2023; Hasan et al 2023c). The community plays an important role; therefore management is based on local resource empowerment. Tradition and customary law are related to and beneficial to coastal and marine resource management efforts in Lembata Regency, East Nusa Tenggara Province (Pinto 2015).

Local wisdom is life values which are passed down from generation to generation in the form of religion, culture or customs, generally in oral form in a social system of a society (Witasari 2022). Sulaiman (2010) defines local wisdom in more detail as knowledge built by the community from generation to generation, related to relationships with nature and natural resources. The existence of local wisdom traditions as a form of marine resource conservation includes control tools, not using dangerous tools, a commitment not to catch and kill sea turtles and a commitment not to throw garbage into the sea as well as seeding and planting mangroves (Zamzawi et al 2017).

Some regions in Indonesia have already owned a lot of local wisdom, as example in environmental management such as PhanglimaLaot, Awig-awig, and Mane'e. The wisdom applied almost fell by the local community, such as phanglimalaot in Aceh and awigawig in Buleleng, Bali. Its existence greatly affects the sustainability of the local environment (Zamzawi et al 2017).

Conclusions. Tebat Rasau ancient lake has high biodiversity. The types of fish with economic value produced from Tebat Rasau Lake include *Wallago* sp., *Channa marulioides*, *Scleropages formosus* and others. The utilization of fish was for consumption and as ornamental fish. The Lanun community's effort to preserve Tebat Rasau geosite is to preserve the local culture of environmentally friendly fishing known as nirok nanggok, and bebanjor. This local tradition is expected to prevent the extinction of fish resources in the future.

Acknowledgements. We gratefully acknowledge to Lanun community, the Rector of Universitas Bangka Belitung and the Institute for Research and Community Service of Universitas Bangka Belitung for their support in this research.

Conflict of interest. The authors declare that there is no conflict of interest.

References

- Alvianna S., Hidayatullah S., Windhyastiti I., Khouroh U., 2022 [The role of green tourism perception, environmental concern and intention of participation in green tourism on environmentally responsible tourism behavior]. Jurnal Manajemen dan Kewirausahaan 10(1):79-87. [in Indonesian]
- Arismayanti N. K., 2015 Pariwisata Hijau Sebagai Alternatif Pengembangan Desa Wisata di Indonesia. Jurnal Analisis Pariwisata 15(2):1-15. [in Indonesian]
- Bidayani E., Reniati R., Valen F. S., 2023 Local wisdom of coastal communities in management of fishery resources in conflict areas of unconventional tin mining in Central Bangka Regency. AACL Bioflux 16(3):1277-1283.
- Djapani Y., Sulaksana N., Muljana B., 2021 Peranan Komunitas Lokal Dalam Perencanaan Pengembangan Geosite Di Kawasan Geopark Belitong. Academia Praja: Jurnal Ilmu Politik, Pemerintahan dan Administrasi Publik 4(1):64-88. [in Indonesian]
- Hassan M., Haq S. M., Ahmad R., Majeed M., Sahito H. A., Shirani M., Mubeen I., Aziz M. A., Pieroni A., Bussmann R. W., Alataway A., Dewidar A. Z., Al-Yafrsi M., Elansary H. O., Yessoufou K., 2022 Traditional use of wild and domestic fauna among different ethnic groups in the western Himalayas a cross cultural analysis. Animals 12(17):2276.
- Hasan V., South J., Valen F. S., Andriyono S., 2023a Endangered black marsh turtle, Siebenrockiella crassicollis (Gray, 1831) (Reptilia, Testudines, Geoemydidae): distribution extension and first record from Belitung Island, Indonesia. Check List 19 (4):505-508.
- Hasan V., Vieira L. O., South J., Ottoni F. P., 2023b First record of *Mastacembelus notophthalmus* (Actinopterygii: Synbranchiformes: Mastacembelidae) for Belitung Island, Indonesia. Acta Ichthyologica et Piscatoria 53:123-127.
- Hasan V., Andraini N. E., Isroni W., Sari L. A., Nafisyah A. L., Dewi N. N., Putri D. N. A., Prasasti T. A. B., Ramadhani A. A., Daniel K., South J., Vieira L. O., Ottoni F. P., Maftuch, Faqih A. R., Wirabuana P. Y. A. P., Tamam M. B., Valen F. S., 2023c Fish diversity of the Bengawan Solo River estuary, East Java, Indonesia. Biodiversitas 24(4):2207-2216.
- Hunn E. S., Brown C. H., 2011 Linguistic ethnobiology. In: Ethnobiology. Anderson E. N., Pearsall D., Hunn E., Turner N. (eds), John Wiley & Sons, Inc., pp. 319-333.
- Jerikho R., Akmal S. G., Hasan V., Yonvitner, Novák J., Magalhães A. L. B., Maceda-Veiga A., Tlusty M. F., Rhyne A. L., Slavík O., Patoka J., 2023 Foreign stingers: South American freshwater river stingrays *Potamotrygon* spp. established in Indonesia. Scientific Reports 13(1):7255.
- Jumadi, Aditya A., 2021 Pariwisata Hijau dan Pemasaran Pariwisata Hijau Sebagai Upaya Mewujudkan Pariwisata Berkelanjutan Di Era Covid-19. Prosiding Seminar Nasional dan Call For Papers, Fakultas Ekonomi Universitas Kristen Immanuel: "Transformasi Indonesia 2021: Inovasi & Strategi Pemulihan Ekonomi Pascapandemi Covid-19", pp. 98-103. [in Indonesian]
- Juniarta H. P., Susilo E., Primyastanto M., 2013 Kajian profil kearifan lokal masyarakat pesisir pulau gili kecamatan Sumberasih kabupaten Probolinggo Jawa Timur. ECSOFiM (Economic and Social of Fisheries and Marine Journal) 1(1):11-25. [in Indonesian]
- Megawati, Tahir T., Hasan M., Dinar M., 2023 Tourism development strategy in improving the community economy at Tanjung Bira Beach, Bonto Bahari District, Bulukumba Regency. Journal of Entrepreneurship Riwew 1(1):1-15
- Ndobe S., Gani A., Bakri A. A., Andriyani D. T., Wuniarto E., Khartiono L. D., Herjayanto M., Hasan V., Moore A. M., 2022 First and recurrent records of *Stiphodon surrufus* Watson & Kottelat, 1995 (Gobiiformes, Gobiidae, Sicydiinae), a naturally rare amphidromous goby, in Sulawesi, Indonesia. Check List 18(2):253-260

- Parmawati R., Pangestuti E., Wike, Hardyansah R., 2020 Development and sustainable tourism strategies in Red Islands beach, Banyuwangi Regency. Journal of Indonesian Tourism and Development Studies 8(3):174-180.
- Pinto Z., 2015 Kajian Perilaku Masyarakat Pesisir yang Mengakibatkan Kerusakan Lingkungan (Studi Kasus di Pantai Kuwaru, Desa Poncosari, Kecamatan Srandakan, Kabupaten Bantul, Provinsi DIY). Jurnal Wilayah dan lingkungan 3(3):163-174. [in Indonesian]
- Pranit D., Musthofa B. M., 2021 Pemberdayaan dan Pengembangan Agrowisata Lebah Madu Trigona Dalam Mendukung Pengembangan Geopark Belitung. Jurnal Vokasi Indonesia 9(2):39-46. [in Indonesian]
- Saputri W. E., Setiawan A. R., 2020 Potret Pelestarian Lingkungan Alam Berdasarkan Kearifan Lokal Masyarakat Colo. SocArXiv f9umc, Center for Open Science. [in Indonesian]
- Sulaiman, 2010 Kebijakan Pengelolaan Perikanan Berbasis Kearifan Lokal di Aceh. KANUN: Jurnal Ilmu Hukun 12(1):176-195. [in Indonesian]
- Utina R., 2012 Kecerdasan Ekologis Dalam Kearifan Lokal Masyarakat Bajo Desa Torosiaje Provinsi Gorontalo. Prosiding Konferensi dan Seminar Nasional Pusat Studi Lingkungan Hidup Indonesia KE 21, 13-15 September 2012 di Mataram, pp. 14-20. [in Indonesian]
- Valen F. S., Notonegoro H., Pamungkas A., Swarlanda, Hasan V., 2023 Revolutionary breakthrough: unveiling the first DNA barcoding of the endemic wild *Betta burdigala* (Kottelat and Ng 1994) (Anabantiformes: Osphronemidae): a critically endangered wid *Betta* from Bangka Island, Indonesia. International Journal of Agriculture and Biology 30(4):269-275.
- Witasari N., 2022 Whispers from the forest: local wisdom in forest conservation and utilization. Paramita: Historical Studies Journal 33(1):23-32
- Zamzawi L., Nursyirwan E., Syahrizal, Ermayanti, 2017 [The local wisdom in marine resource conservation in Indonesia: a case study of newcomers in Pariaman, West Sumatra]. Advances in Social Science, Education and Humanities Research, volume 136. 2nd International Conference on Social and Political Development (ICOSOP 2017), Atlantis Press, pp. 391-400. [in Indonesian]
- Zukhri N., Rosalina E., Christianingrum C., 2021 Geopark Belitong: environment based tourism branding in Belitung Island. IOP Conference Series: Earth and Environmental Science 926(1):012075.
- *** https://indonesia.un.org/en/128882-introducing-belitong-island-indonesia%E2%80% 99s-newest-unesco-global-geopark.
- *** https://www.babelinsight.id/content/read/239/danau-purba-tebat-rasau-bagian-dari-geopark-belitong/babelinsight.id.

Received: 16 August 2023. Accepted: 15 October 2023. Published online: 17 November 2023.

Endang Bidayani, Aquaculture Department, Faculty of Agriculture Fisheries and Biology, Universitas Bangka Belitung, ampus terpadu UBB street, Balunijuk, Bangka Belitung 33127, Indonesia, e-mail: endangbidayani@gmail.com

Devi Valeriani, Department of Economics, Faculty of Economics, Universitas Bangka Belitung, Bangka, 33172, Indonesia, e-mail: deviyono92@gmail.com

Fitri Sil Valen, Aquaculture Department, Faculty of Agriculture Fisheries and Biology, Universitas Bangka Belitung, Kampus terpadu UBB street, Balunijuk, Bangka Belitung 33127, Indonesia, e-mail: fitrisilvalen@ubb.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Bidayani E., Valeriani D., Valen F. S., 2023 Ethnozoology of Lanun community towards biodiversity conservation at Tebat Rasau ancient lake geosite in Simpang Renggiang District, East Belitung Regency. AACL Bioflux 16(6):2931-2938.

Ethnozoology of Lanun community towards biodiversity conservation at Tebat Rasau ancient lake geosite in Simpang Renggiang District, East Belitung Regency

ORIGIN	ALITY REPORT			
1 SIMIL	2% ARITY INDEX	14% INTERNET SOURCES	8% PUBLICATIONS	5% STUDENT PAPERS
PRIMAF	RY SOURCES			
1	www.bi	oflux.com.ro		4%
2	www.bi	3%		
3	Submitted to Flagler College Student Paper			2%
4	WWW.Wyssen.com Internet Source			2%
5	WWW.U r Internet Sour	nesco.org		2%

Exclude quotes On Exclude bibliography On

Exclude matches

< 2%