

Korespondensi Artikel

1. Submitted: July 20, 2025

Manuscript ID: EJABF-2510-6839

Manuscript Title: Growth Performance of Nile Fish and Catfish Fry Cultivated with Bio-Enzyme Supplementation and Its Agribusiness Perspective

Authors: Andri Kurniawan, Muntoro Muntoro, Ardiansyah Kurniawan, UFI AYU WULANDARI, Didin Erlan, Venisa Pratama, Syafiq Muhammad Syakir, Danella Aisah Maheswari, Safitri Fadia, Cahya Fitri

Dear Dr. Andri Kurniawan

I wish to acknowledge receiving the of the above mentioned manuscript.

It should be noted that the manuscript will be reviewed for possible publication in the Scientific Journals Management System.

Please be sure that the submitted manuscript has not been published previously and will not be submitted elsewhere prior to our decision.

Our editorial decision will be brought to your attention once the paper has been reviewed due the referees consideration.

I wish to take this opportunity to thank you for sharing your work with us.

Truly yours,

Executive managing Editor of Egyptian Journal of Aquatic Biology and Fisheries

2. Review form Reviewer: Aug 2, 2025

Reviewer comments of article are attached

3. Revision by Author: Aug 20, 2025

Dear Managing Editor of EJABF,

I am writing to confirm that I have submitted the revised version of our manuscript, prepared in accordance with the reviewer's comments. I have also uploaded the proof of payment for the article processing charge.

Manuscript details

Manuscript ID: *EJABF-2510-6839

*

Title: Growth Performance of Nile Fish and Catfish Fry Cultivated with Bio-Enzyme Supplementation and Its Agribusiness Perspective

Kindly confirm receipt of the revised files and the payment proof, and please let me know if any further information or actions are required from my side.

Thank you for your attention and consideration.

Best regards,


Dr. Andri Kurniawan

Department of Biology, Faculty of Science and Engineering

Bangka Belitung University

 andri_pangkal@yahoo.co.id

4. Accepted: Sep. 20, 2025
5. LoA Confirmation: Oct. 28, 2025



Oct. 28, 2025

Acceptance

Dear Muntoro, Andri Kurniawan, Ardiansyah Kurniawan, Ufi Ayu Wulandari, Didin Erlan, Venisa Pratama³, Syafiq Muhammad Syakir, Danella Aisah Maheswari, Safitri Fadia, Cahya Fitri

I am pleased to inform you that your paper entitled "Growth Performance of Tilapia and Catfish Fry Cultivated with Bio-Enzyme Supplementation and Its Agribusiness Perspective" (Manuscript ID: EJABF- 2510-6839) has been accepted for publication in the Egyptian Journal of Aquatic Biology and Fisheries (EJABF).

Congratulations on your successful submission, and we look forward to featuring your valuable research in our journal.

Regards,
Prof. Magdy T. Khalil
Co-Editor-In Chief of EJABF Journal,
Faculty of Science,
Ain Shams University,
Cairo, Egypt.

magdy khalil



6. Published: 29 Oktober 2025

Dear Author,

I am delighted to inform you that your manuscript has been successfully published in our journal, EJABF. Congratulations on this significant achievement!

You can find your article published in Volume 29 Issue 5; article 174

We are proud to have your work featured in our journal and look forward to any future contributions you may have.

Once again, congratulations

Best regards,

Mohamed Elnaggar

Administrative Director
EJABF



Growth Performance of Nile Fish and Catfish Fry Cultivated with Bio-Enzyme Supplementation and Its Agribusiness Perspective

Muntoro¹, Andri Kurniawan^{2*}, Ardiansyah Kurniawan³, Ufi Ayu Wulandari³, Didin Erlan³, Venisa Pratama³, Syafiq Muhammad Syakir³, Danella Aisah Maheswari², Safitri Fadiah², Cahya Fitri²

¹Department of Agribusiness, Faculty of Agriculture, Fisheries and Marine, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

²Department of Biology, Faculty of Science and Engineering, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

³Department of Aquaculture, Faculty of Agriculture, Fisheries and Marine, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

*Corresponding Author: andri_pangkal@yahoo.co.id

ARTICLE INFO

Article History:

Received: June 1, 2025

Accepted: July 17, 2025

Online:

Keywords:

Aquaculture,
Bio-Enzyme;
Effectiveness; Growth
Rate; Fish Fry

ABSTRACT

Nile fish and catfish are freshwater fish commodities that are widely cultivated by the community because of their relatively easy maintenance, low cost, and short time. The quality of the cultivation water environment is an important factor and influences the growth performance and life of these commodities. This study aimed to analyze the growth performance of Nile fish and catfish fry cultivated with the addition or supplementation of bio-enzymes with different concentrations in the maintenance water. This study used an experimental method with three treatments: control (without bio-enzymes), 500 ml/900 L or 0.6 ml/L bio-enzymes, and 1000 ml/900 L or 1.1 ml/L bio-enzymes. The results illustrate that bio-enzyme supplementation into Nile fish fry ponds compared to the control treatment without bio-enzymes showed a higher contribution to weight growth than length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation into catfish fry ponds compared to the control without bio-enzymes showed a higher contribution to length growth than weight growth. Supplementation with 500 ml of bio-enzyme resulted in the highest absolute weight growth of Nile fish fry, which was 16.95 g with a daily weight growth rate of 0.42 g/day and a specific weight growth rate of 3.35% and 100% survival rate. Supplementation with 500 ml of bio-enzyme resulted in the highest absolute length growth of catfish fry, which was 9.20 cm with a daily length growth rate of 0.23 cm/day and a specific length growth rate of 2.23% and 100% survival rate. Further research is needed to determine the effectiveness of supplementation dosage as well as the time and frequency of bio-enzyme supplementation on improving water quality as a culture medium and the growth performance of cultivated fish fry. In agribusiness perspective, using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming.

INTRODUCTION

Fish farming, known as aquaculture, is one of the leading fish food sources with the fastest growth in productivity worldwide (**Sumaila *et al.*, 2022**). Freshwater fish farming has been widely developed by communities, both on a household and large scale. Among the freshwater fish commonly cultivated by the community are Nile fish and catfish (**Nalle *et al.*, 2021**).

The development of fish farming productivity is greatly influenced by water quality, which significantly supports fish growth performance (**Tumwesigye *et al.*, 2022**; **Ajayi & Okoh, 2014**). Water quality, manifested in the form of physical, chemical, and biological parameters, affects the entire life cycle, including survival, growth, and reproduction (**Siegers *et al.*, 2019**; **Koniyo, 2020**; **Kulla *et al.*, 2020**).

Aquaculture water quality is generally contaminated by organic matter from leftover feed and metabolic waste in the form of fish feces. One effort to maintain and improve water quality in fish farming is the addition of enzymes to the culture environment.

Bio-enzymes or Bioz are fermented products made from fruits or vegetables that are widely used for various purposes due to their probiotic microorganism content and bioactive compounds (**Ducatelle *et al.*, 2018**). It is produced in 3 months fermentation process with ratio 1 : 3 : 10 for sugar : waste : water (**Rasit *et al.*, 2019**; **Galintin *et al.*, 2021**; **Kurniawan *et al.*, 2024**). Bio-enzymes are also known as ecoenzymes, garbage enzymes, terrazymes, fruit enzymes, flower enzymes, and so on (**Neupane & Khadka, 2019**; **Singh *et al.*, 2019**). It is a fermentation product.

One use of bio-enzymes is to accelerate the waste degradation process and improve environmental quality (**Ardianita *et al.*, 2023**). However, several studies have shown that the use of bio-enzymes (ecoenzymes) in fish farming environments does not show a different effect on reducing ammonia, nitrate, and nitrite and does not result in good fish growth (**Kusumawati *et al.*, 2018**; **Bambang *et al.*, 2024**). Based on the above description, this study was conducted to confirm previous studies by analyzing the growth performance of Nile fish and catfish fry cultivated with the addition or supplementation of bio-enzymes with different concentrations in the maintenance water.

MATERIALS AND METHODS

This research was conducted at the Hatchery, Bangka Belitung University, from June to August 2025. The samples used in this study were 100 Nile fish fry measuring 6-8 cm and catfish fry measuring 5-7 cm in each pond. Both types of samples were cultivated in concrete ponds measuring 2 m long, 1.5 m wide, and 0.6 m high, filled with water to a depth of 30 cm.

The research treatment consisted of two, namely cultivation media without added bio-enzymes and cultivation media added bio-enzymes. Bio-enzyme supplementation into the cultivation media was carried out before fish maintenance with doses of 500 ml and 1000 ml in each pond containing 900 L of water or treatments of 0.6 ml/L and 1.1 ml/L. The test sample fish fry were maintained in the research pond for 40 days by being fed pellets *ad satiation*, namely feeding until the fish were full and stopped eating.

Fish sample measurements were conducted on 10% of the population in each treatment. The research parameters observed in fish were absolute weight growth (g), absolute length growth (cm), daily weight growth rate (g/day), daily length growth rate (cm/day), specific weight growth rate (%), specific length growth rate (%), and survival (%). The water quality parameters observed were pH, temperature (°C), dissolved oxygen (DO) (ppm), total dissolved solids (TDS) (ppm), and electrical conductivity (EC) (mS/cm). The research data were analyzed descriptively and presented in tables and graphs.

RESULTS AND DISCUSSION

The results of the study showed that bio-enzyme supplementation into the pond compared to the control treatment without bio-enzyme can have an effect or no effect on the growth performance of fish fry. Bio-enzyme supplementation into Nile fish fry ponds contributed more to higher weight growth than length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation into catfish fry ponds resulted in higher length growth than weight growth. Supplementation of 500 ml of bio-enzyme resulted in the highest absolute weight growth of Nile fish fry, namely 16.95 g with a daily weight growth rate of 0.42 g/day and a specific weight growth rate of 3.35% and 100% survival. Supplementation of 500 ml of bio-enzyme resulted in the highest absolute length growth of catfish fry, namely 9.20 cm with a daily length growth rate of 0.23 cm/day and a specific length growth rate of 2.23% and 100% survival.

Growth of Nile fish

Supplementation of bio-enzymes into the cultivation medium, namely the maintenance water in the Nile fish fry pond resulted in higher weight growth of the fry at the end of the observation compared to the control treatment without bio-enzymes. However, the length growth of the Nile fish fry reared in the pond with bio-enzyme supplementation was lower than the control treatment without bio-enzymes (Fig. 1). The absolute growth, daily growth rate, and specific growth rate of the Nile fish fry were shown in Table 1. The research data also did not show any symptoms of disease in the fry reared, but there were 3 fish that died during maintenance in the 1000 ml bio-enzyme treatment, while in the other treatments no dead fry were found during maintenance (Table 1).




Fig. 1. Growth in weight and length of Nile fish fry

Table 1. Calculation of growth and survival of Nile fish fry

Parameter	Nile fish fry		
	Control	Bio-enzyme	
		500 ml	1000 ml
Absolute weight gain (g)	11.94	16.95	16.47
Absolute length growth (cm)	17.75	12.98	10.20
Daily weight gain rate (g/day)	0.30	0.42	0.41
Daily length growth rate (cm/day)	0.44	0.32	0.26
Specific weight growth rate (%)	1.88	3.35	3.13
Specific length growth rate (%)	2.98	2.74	2.23
Survival rate (%)	100	100	97

Growth of Catfish

Supplementation of bio-enzymes into the culture medium, namely the maintenance water in the catfish fry pond, resulted in lower weight growth of the fry at the end of the observation compared to the control treatment without bio-enzymes. However, the length growth of the catfish fry reared in the pond with bio-enzyme supplementation was higher than the control treatment without bio-enzymes (Fig. 2). The absolute growth, daily growth rate, and specific growth rate of the catfish fry were shown in Table 1. The research data also showed no symptoms of disease and no dead fry were found during the maintenance (Table 2).




Fig. 2. Growth in weight and length of catfish fry

Table 2. Calculation of growth and survival of catfish fry

Parameter	Catfish fry		
	Control	Bio-enzyme	
		500 ml	1000 ml
Absolute weight gain (g)	15.48	15.40	13.28
Absolute length growth (cm)	7.55	9.20	7.85
Daily weight gain rate (g/day)	0.39	0.39	0.33
Daily length growth rate (cm/day)	0.19	0.23	0.20
Specific weight growth rate (%)	5.65	5.69	5.23
Specific length growth rate (%)	1.94	2.23	1.99
Survival rate (%)	100	100	100

Water quality as a medium for cultivating Nile fish and catfish fry

The results of water quality measurements used as a culture medium for Nile fish and catfish fry—pH, temperature, DO, TDS, and EC—showed that all parameters were dynamic. This indicates that water quality fluctuates during cultivation and does not reflect a relationship between bio-enzyme supplementation in the culture medium and water quality.

Water quality in the Nile fish fry rearing ponds showed a pH value that tended to be higher than that in the catfish fry ponds. The water temperature in the Nile fish fry rearing ponds was relatively lower than that in the catfish fry ponds. Dissolved oxygen in the water in the Nile fish fry ponds was relatively better than that in the catfish fry ponds. Total dissolved solids and electrical conductivity in the Nile fish and catfish fry control ponds were relatively higher at certain times and relatively lower at other times compared to the TDS in the bio-enzyme treatment ponds.

The water quality in the Nile fish fry pond had the highest pH value in the 500 ml bio-enzyme treatment, the highest temperature in the 1000 ml bio-enzyme treatment, the highest DO in the 500 ml bio-enzyme treatment, and the highest TDS and EC in the control treatment. The water quality in the catfish fry pond had the highest pH value in the 500 ml bio-enzyme treatment, the highest temperature in the control treatment, and the highest DO, TDS, and EC in the control treatment. The water quality conditions and comparison between the water quality in the Nile fish and catfish fry ponds were shown in Fig. 3 and Fig. 4.




Fig. 3. Water quality as a medium for Nile fish cultivation




Fig. 4. Water quality as a medium for catfish cultivation

Sikku *et al.* (2023) conducted a study on the application of 5-15 ml/L of ecoenzyme (bio-enzyme) in Nile fish fry (3 cm in length) cultivation pond maintained for 45 days. The absolute weight produced ranged from 0.98-1.76 g with a specific weight growth rate ranging from 2.33-4.19%. The water pH value during maintenance ranged from 7.6-9.1. When compared with the specific weight growth rate in this study which ranged from 3.13-3.35% (Table 1), then supplementation with a high concentration of bio-enzyme (15 ml/L) resulted in a better specific weight growth rate than the low concentration (0.6-1.1 ml/L) in this study. However, at a concentration of 5-10 ml/L in the study by **Sikku *et al.* (2023)** showed a lower value (2.33-2.98%) compared to the treatment in this study. Thus, there is no linear pattern that shows that the higher the concentration of bio-enzyme given, the higher the growth of Nile fish fry.

Bio-enzyme research has also been conducted on catfish fry cultivation. The results of **Kusumawati *et al.* (2018)** research show that the addition of bio-enzymes to catfish fry ponds maintained for 4 weeks (approximately 28 days) has a length of 5.9-8.0 cm with a weight of 2.01-4.32 g from a length of 5.1-5.8 cm and a weight of 1.12-1.22 g on day 0. Meanwhile, the results of measurements on the 26th day of this study have shown that the weight and length data were 8.76 g and 9.22 cm respectively in the control treatment; 8.14 g and 10.12 cm in the 500 ml bio-enzyme treatment (0.6 ml/L); and 7.90 g and 9.82 cm in the 1000 ml bio-enzyme treatment (1.1 ml/L) (Figure 4). This difference may be caused by the contribution of bio-enzymes in the research of **Kusumawati *et al.* (2018)** found lower pH and DO values, ranging from 5.85 to 6.89 and DO from 2.95 to 3.71 ppm in the fourth week of culture. This study, however, found pH values ranging from 8.33 to 9.08 and DO from 4.8 to 6.8 ppm. Based on SNI 6484.4:2014, water quality for catfish fry culture has a pH of 6.5 to 8 and DO > 3 ppm.

The description shows that specifically, the contribution of bio-enzyme supplementation to the growth performance of Nile fish fry was better than that of catfish fry (Table 1 and Table 2). This was in line with the statement by **Herliwati *et al.* (2024)** that the addition of bio-enzymes can improve water quality, which can impact the survival of catfish fry. However, the addition of bio-enzymes to the cultivation water medium only affected the growth of Nile fish and not catfish. Similar conditions were also found in the results of research by **Mado *et al.* (2023)** and **Natsir *et al.* (2024)** who found increased growth of Nile fish with the application of bio-enzymes to the cultivation water medium. However, according to **Kusumawati *et al.* (2018)** and **Bambang *et al.* (2024)**, catfish growth did not change with the same application. This is possible because catfish have a tolerance to less than ideal waters (**Hidayati *et al.*, 2021**), so that when water quality improves, its metabolism does not affect. Meanwhile, Nile fish experience increased metabolism when quality improves, resulting in better growth. Nile fish fry experience increased growth and survival when oxygen and ammonia levels are more appropriate (**Islami *et al.*, 2017**). When compared to the pH, DO, and TDS conditions in Fig. 3 and Fig. 4, it can be seen that supplementing bio-enzymes into the culture medium results in better water quality for Nile fish fry cultivation than for catfish fry. Furthermore, Fig. 3 and Fig. 4 also show that the total dissolved solids in catfish fry ponds were higher than in Nile fish fry ponds. High dissolved solids can hinder the growth performance of fish fry.

Rahman *et al.* (2020) explained that total dissolved solids are organic and inorganic substances dissolved in water that can be a factor causing water turbidity. According to **Damayanti (2014)**, increased TDS concentrations in water can originate from feed waste and accumulated metabolic waste such as fish feces. According to **Al Zamzami *et al.* (2019)**, the length of maintenance time can contribute to an increase in dissolved organic matter, one of which comes from feed waste and fish feces. Furthermore, various studies have revealed that the presence of nitrogen waste can produce ammonia, nitrate, and nitrite

compounds that can cause oxygen depletion, disease, and even death in fish (**Munguti *et al.*, 2021; Li *et al.*, 2023; Mustafa *et al.*, 2024; Kurniawan *et al.*, 2025**).

Relatively low bio-enzyme concentrations were considered ineffective in reducing waste contamination, particularly in catfish fry ponds. This is visually confirmed by the cloudier and more odorous water in catfish fry ponds compared to Nile fish fry ponds. This is thought to inhibit the growth of catfish fry in rearing ponds. **Kusumawati *et al.* (2018)** explained that administering bio-enzymes at low concentrations does not significantly reduce ammonia and nitrite concentrations. This contributed to the slow growth of catfish fry.

Higher bio-enzyme concentrations, duration, and frequency of bio-enzyme administration need to be tested for their effectiveness in improving the quality of aquaculture water and its effect on optimal growth for fish fry. This is in line with **Putri *et al.* (2024)** who stated that the longer the time and the higher the eco-enzyme dosage, the higher the effectiveness in reducing waste contamination as indicated by the biological oxygen demand (COD) and total suspended solids (TSS) in the environment. **Yatma *et al.* (2024)** also revealed that increasing the eco-enzyme dosage used was able to significantly reduce the dissolved solids (TDS) content and improve water quality. Furthermore, water quality and nutrient contribute to fish morphological and its growth (**Serdiati *et al.*, 2024; Mustikasari *et al.*, 2020**).

Discussion about Agribusiness Perspective of Bio-Enzyme Supplementation for Aquaculture

From an agribusiness perspective, bio-enzyme supplementation in aquaculture presents a high-potential growth sector that aligns with key trends: sustainable food production, environmental regulation, and consumer preference for clean-label seafood. Strategic investment in research and development, production, and farmer outreach can unlock long-term profitability and ecological benefits.

Perspective in agribusiness opportunities, including market potential, value addition, product development and diversification, challenges from an agribusiness viewpoint, and investment and agribusiness models. In Indonesia (and many other countries) Nile fish and catfish are major freshwater/pond-aquaculture species. Grower demand is strong for both local consumption and export. A value proposition; higher growth/lower feed cost/improved survival is very attractive to fish farmers, in particular Nile fish and catfish. Use the enzyme supplementation in aquaculture can improve output and profit. In other hand, enzyme supplementation can reduce feed costs (better feed conversion ratio or FCR), improves animal health and survival rates — leading to higher yields and profits, reduces environmental impact, enhancing the farm's market reputation and compliance with green regulations (Varshini & Gayathri, 2023).

Using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming. The benefits using bio-enzymes in aquaculture from an agribusiness perspective:

Improved Productivity and Profitability

Enhanced Feed Conversion Ratio (FCR) where bioenzymes help fish and shrimp digest feed more efficiently, reducing feed costs — a major operational expense. Faster growth rates where enzymes promote better nutrient absorption, accelerating growth and reducing time to market. Lower mortality rates where enzymes improve gut health and immunity, reducing disease outbreaks.

Reduction in Input Costs

Less dependency on antibiotics and chemicals where enzymes can reduce disease pressure naturally. Better water quality management where bioenzymes help degrade organic waste and ammonia, reducing the need for chemical water treatments.

Sustainable Branding & Market Access

Eco-friendly production where enzyme use aligns with growing consumer demand for sustainable and antibiotic-free aquaculture. Certification ready where farms using bioenzymes can more easily meet organic or eco-certification standards.

Waste Reduction and Circular Economy

Utilization of agricultural byproducts where some bioenzymes are produced using agri-waste (e.g., fruit peels, crop residues), supporting circular bioeconomy models. Efficient waste breakdown where help manage sludge and organic waste, reducing environmental impact.

New Revenue Streams

Local production of bioenzymes where farmers or agri-entrepreneurs can engage in low-cost production of bioenzymes using fermentation, creating micro-enterprises. Diversification for agro-industries where companies in biotech, agriculture, or food waste management can expand into aquaculture support services.

Regulatory and Environmental Compliance

Supports green policies where use of bioenzymes aligns with government and international sustainability goals in aquaculture. Reduces pollution where better water quality means less eutrophication and contamination of surrounding ecosystems.

CONCLUSION

The addition or supplementation of bio-enzymes to Nile fish and catfish fry ponds produced different contributions. Bio-enzyme supplementation to Nile fish fry ponds compared to the control treatment without bio-enzymes showed a higher contribution to weight growth than to length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation to catfish fry ponds compared to the control without bio-enzymes showed a higher contribution to length growth than to weight growth. The results of water quality measurements as a medium for cultivating Nile fish and catfish fry, namely pH, temperature, DO, TDS, and EC showed that all parameters were dynamic. This means that water quality fluctuates during cultivation and does not reflect the relationship between bio-enzyme supplementation to the cultivation medium and water quality. In agribusiness perspective, using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming.

ACKNOWLEDGMENT

The authors would like to thank Bangka Belitung University for supporting and funding this research and publication through Grant of Penelitian Unggulan 2025 with grant number 1116/UN50/M/PP/2025.

CONFLICTS OF INTEREST

All the authors declare no conflict of interest.

FUNDING

This research and publication were supported and funded through the Grant of Penelitian Unggulan 2025 grant number: 1116/UN50/M/PP/2025.

REFERENCES

- Ajayi, A. O. and Okoh, A. I.** (2014). Bacteriological study of pond water for aquaculture purposes. *Journal of Food Agriculture and Environment*, 12(2): 1260-1265.
- Al Zamzami, I. M.; Sari, A. H. W. and Perwira, I. Y.** (2019). Fluktuasi bahan organik dan residu terlarut dalam budidaya ikan koi di Tulungagung Jawa Timur. *Current Trends in Aquatic Science*, 2(1): 79-84.
- Ardianita, N.; Andriani, R. and Maya, N. P.** (2023). Efektivitas ekoenzim terhadap kelulushidupan ikan lele (*Clarias* sp.) pada media yang tercemar limbah batik. *Biology Natural Resources Journal*, 2(1): 24-27.

- Bambang, B.; Umar, N. A. and Indrawati, E.** (2024). Pengaruh ekoenzim dosis berbeda terhadap perbaikan kualitas air budidaya dalam pembesaran ikan lele mutiara *Clarias* sp. *Journal of Aquaculture and Environment*, 6(2): 84-89.
- Damayanti, H. O.** (2014). Tinjauan kualitas dan dampak ekonomi konsentrasi total dissolved solid (TDS) air di area pertambakan Desa Bulumanis Kidul. *Jurnal Litbang: Media Informasi Penelitian, Pengembangan dan IPTEK*, 10(2): 103-113.
- Ducatelle, R.; Goossens, E.; De Meyer, F.; Eeckhaut, V.; Antonissen, G.; Haesebrouck, F. and Van Immerseel, F.** (2018). Biomarkers for monitoring intestinal health in poultry: Present status and future perspectives. *Veterinary Research*, 49: 43.
- Galintin, O.; Rasit, N. and Hamzah, S.** (2021). Production and characterization of eco enzyme produced from fruit and vegetable wastes and its influence on the aquaculture sludge. *Biointerface Research in Applied Chemistry*, 11(3): 10205-10214.
- Herliwati, H.; Rahman, M.; Ahmadi, A. and Tony, F.** (2024). Variations in the addition of eco enzyme in feed on the survival of siamese catfish (*Pangasius hypophthalmus*) seeds raised in aquariums. *Fish Scientiae*, 14(2): 145-154.
- Hidayati, S. N.; Laili, S. and Santoso, H.** (2021). Pengaruh kualitas air kolam terpal terhadap pertumbuhan ikan lele dumbo (*Clarias gariepinus*). *Jurnal Ilmiah Biosaintropis*, 6(2): 19-25.
- Islami, A. N.; Hasan, Z. and Anna, Z.** (2017). Pengaruh perbedaan siphonisasi dan aerasi terhadap kualitas air, pertumbuhan, dan kelangsungan hidup pada budidaya ikan nila (*Oreochromis niloticus*) stadia benih. *Jurnal Perikanan Kelautan*, 8(1): 73-82.
- Koniyo, Y.** (2020). Analisis kualitas air pada lokasi budidaya ikan air tawar di Kecamatan Suwawa Tengah. *Jurnal Technopreneur*, 8(1): 52-58.
- Kulla, O. L. S.; Yuliana, E. and Supriyono, E.** (2020). Analisis kualitas air dan kualitas lingkungan untuk budidaya ikan di Danau Laimadat, Nusa Tenggara Timur. *Pelagicus*, 1(3): 135-144.
- Kurniawan, A.; Mustikasari, D.; Kurniawan, A.; Muntoro, M.; Setiadi, J. and Kholishah, N. A.** (2024). The preliminary study about physico-chemical property of bio-enzyme produced from orange fruits waste treated with different concentrations of probiotic. *Jurnal Ilmu Lingkungan*, 22(4), 861-867.
- Kurniawan, A.; Mustikasari, D.; Muntoro.; Setiadi, J.; Wulandari, U. A.; Pratama, V.; Fadia, S.; Maheswari, D. A.; Fitri, C. and Iqbal, D. H.** (2025). Effectiveness

of decaying fruits as a carbon source on fish productivity and water quality improvement in aquaculture: A narrative review. *Egyptian Journal of Aquatic Biology & Fisheries*, 29(4): 2459-2481.

Kusumawati, A. A.; Suprpto, D. and Haeruddin, H. (2018). Pengaruh ekoenzim terhadap kualitas air dalam pembesaran ikan lele (*Clarias gariepinus*), 7(4): 307-314.

Li, H.; Cui, Z.; Cui, H.; Bai, Y.; Yin, Z. and Qu, K. (2023). Hazardous substances and removal in review. *Aquaculture*, 569: 739399.

Mado, A. M.; Umar, N. A. and Aqmal, A. (2023). Pengaruh eco-enzym terhadap kualitas air pada sistem budidaya akuaponik ikan nila *Oreochromis niloticus* dan tanaman sawi *Brassica juncea*. *Journal of Aquaculture and Environment*, 5(2): 66-72.

Munguti, J. M.; Sendi, J. G.; Obiero, K. O.; Ogello, E. O.; Kyule, D. N.; Liti, D. M. and Musalia, L. M. (2021). Aqua-feed wastes: Impact on natural systems and practical mitigations-A review. *Journal of Agricultural Science*, 13(1): 111-121.

Mustafa, S. A.; Al-Rudainy, A. J. and Salman, N. M. (2024). Effect of environmental pollutants on fish health: An overview. *Egyptian Journal of Aquatic Research*, 50(2): 225-233.

Mustikasari, D.; Suryaningsih, S. and Nuryanto, A. (2020). Morphological variation of blue panchax (*Aplocheilichthys panchax*) lives in different habitat assessed using truss morphometric. *Biosaintifika: Journal of Biology & Biology Education*, 12(3): 399-407.

Nalle, M. M. D.; Kamlasi, Y. and Edo, S. I. (2021). Budidaya ikan lele dan ikan nila oleh Kelompok Tani Tunfeu Desa Ponain Kecamatan Amarasi Kabupaten Kupang di masa Covid-19. *Jurnal Pengabdian Perikanan Indonesia*, 1(2): 151-156.

Natsir, N. I.; Jayadi, J. and Hadijah, S. (2024). Aplikasi eco-enzyme terhadap pertumbuhan dan tingkat kelangsungan hidup ikan nila (*Oreochromis niloticus*). *Jurnal Akuakultur Nusantara*, 1(2): 96-106.

Neupane, K. and Khadka, R. (2019). Production of garbage enzyme from different fruit and vegetable wastes and evaluation of its enzymatic and antimicrobial efficacy. *Tribhuvan University Journal of Microbiology*, 6: 113-118.

Putri, M. C.; Widiarini, R. and Ramadanintyas, K. N. (2024). Effectiveness of eco enzymes in reducing chemical parameters of tofu factory liquid waste in Klumutan Village, Saradan District. *Journal of Social Research*, 3(9): 1-10.

- Rahman, M. F. I.; Wibisana, H. and Zainab, S. (2020).** Analisa dan pemetaan total padatan terlarut di pesisir Pantai Pasuruan dengan citra satelit Terra Modis. *Journal of Civil Engineering*, 16(2): 144-156.
- Rasit, N.; Fern, L. H. and Ghani, W. A. W. A. K. (2019).** Production and characterization of eco enzyme produced from tomato and orange wastes and its influence on the aquaculture sludge. *International Journal of Civil Engineering and Technology*, 10(3): 967-980.
- Serdiati, N.; Islamy, R. A.; Mamat, N. B.; Hasan, V. and Valen, F. S. (2024).** Nutritional value of alligator weed (*Alternanthera philoxeroides*) and its application for herbivorous aquaculture feed. *International Journal of Agriculture and Biosciences*, 13(3): 318-324.
- Siegers, W. H.; Prayitno, Y. and Sari, A. (2019).** Pengaruh kualitas air terhadap pertumbuhan ikan nila nirwana (*Oreochromis* sp.) pada tambak payau. *The Journal of Fisheries Development*, 3(2): 95-104.
- Sikku, E. S.; Umar, N. A. and Indrawati, E. (2023).** Penggunaan eco-enzim dengan dosis berbeda pada teknologi akuaponik sederhana untuk optimalisasi pertumbuhan ikan nila *Oreochromis niloticus*. *Journal of Aquaculture and Environment*, 5(2): 60-65.
- Singh, H.; Prasad, A. and Nal, K. V. (2019).** Study and analysis of bio enzymes based natural products. *International Journal of Engineering Applied Sciences and Technology*, 4(8): 205-207.
- Tumwesigye, Z.; Tumwesigye, W.; Opio, F.; Kemigabo, C. and Mujuni, B. (2022).** The effect of water quality on aquaculture productivity in Ibanda District, Uganda. *Aquaculture Journal*, 2(1): 23-36.
- Varshini, B. and Gayathrif, V. (2023).** Role of eco-enzymes in sustainable development. *Nature Environment and Pollution Technology*, 22(3): 1299-1310.
- Yatma, M. A.; Hamzani, S. and Zubaidah, T. (2024).** The effect of eco enzyme on improving river water quality based on turbidity, temperature, pH, and TDS parameters. *Global Health & Environmental Perspectives*, 1(3): 164-168.



Growth Performance of Nile Fish and Catfish Fry Cultivated with Bio-Enzyme Supplementation and Its Agribusiness Perspective

Muntoro¹, Andri Kurniawan^{2*}, Ardiansyah Kurniawan³, Ufi Ayu Wulandari³, Didin Erlan³, Venisa Pratama³, Syafiq Muhammad Syakir³, Danella Aisah Maheswari², Safitri Fadiah², Cahya Fitri²

¹Department of Agribusiness, Faculty of Agriculture, Fisheries and Marine, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

²Department of Biology, Faculty of Science and Engineering, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

³Department of Aquaculture, Faculty of Agriculture, Fisheries and Marine, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

*Corresponding Author: andri_pangkal@yahoo.co.id

ARTICLE INFO

Article History:

Received: June 1, 2025

Accepted: July 17, 2025

Online:

Keywords:

Aquaculture,
Bio-Enzyme;
Effectiveness; Growth
Rate; Fish Fry

ABSTRACT

Nile fish and catfish are freshwater fish commodities that are widely cultivated by the community because of their relatively easy maintenance, low cost, and short time. The quality of the cultivation water environment is an important factor and influences the growth performance and life of these commodities. This study aimed to analyze the growth performance of Nile fish and catfish fry cultivated with the addition or supplementation of bio-enzymes with different concentrations in the maintenance water. This study used an experimental method with three treatments: control (without bio-enzymes), 500 ml/900 L or 0.6 ml/L bio-enzymes, and 1000 ml/900 L or 1.1 ml/L bio-enzymes. The results illustrate that bio-enzyme supplementation into Nile fish fry ponds compared to the control treatment without bio-enzymes showed a higher contribution to weight growth than length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation into catfish fry ponds compared to the control without bio-enzymes showed a higher contribution to length growth than weight growth. Supplementation with 500 ml of bio-enzyme resulted in the highest absolute weight growth of Nile fish fry, which was 16.95 g with a daily weight growth rate of 0.42 g/day and a specific weight growth rate of 3.35% and 100% survival rate. Supplementation with 500 ml of bio-enzyme resulted in the highest absolute length growth of catfish fry, which was 9.20 cm with a daily length growth rate of 0.23 cm/day and a specific length growth rate of 2.23% and 100% survival rate. Further research is needed to determine the effectiveness of supplementation dosage as well as the time and frequency of bio-enzyme supplementation on improving water quality as a culture medium and the growth performance of cultivated fish fry. In agribusiness perspective, using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming.

Commented [H1]: Tilapia

Replace all

Commented [H2]: make analysis for agribusiness perspective

INTRODUCTION

Fish farming, known as aquaculture, is one of the leading fish food sources with the fastest growth in productivity worldwide (Sumaila *et al.*, 2022). Freshwater fish farming has been widely developed by communities, both on a household and large scale. Among the freshwater fish commonly cultivated by the community are Nile fish and catfish (Nalle *et al.*, 2021).

Commented [H3]: add more reference

The development of fish farming productivity is greatly influenced by water quality, which significantly supports fish growth performance (Tumwesigye *et al.*, 2022; Ajayi & Okoh, 2014). Water quality, manifested in the form of physical, chemical, and biological parameters, affects the entire life cycle, including survival, growth, and reproduction (Siegers *et al.*, 2019; Koniyo, 2020; Kulla *et al.*, 2020).

Commented [H4]: pay attentions

Aquaculture water quality is generally contaminated by organic matter from leftover feed and metabolic waste in the form of fish feces. One effort to maintain and improve water quality in fish farming is the addition of enzymes to the culture environment.

Bio-enzymes or Bioz are fermented products made from fruits or vegetables that are widely used for various purposes due to their probiotic microorganism content and bioactive compounds (Ducatelle *et al.*, 2018). It is produced in 3 months fermentation process with ratio 1 : 3: 10 for sugar : waste : water (Rasit *et al.*, 2019; Galintin *et al.*, 2021; Kurniawan *et al.*, 2024). Bio-enzymes are also known as ecoenzymes, garbage enzymes, terrazymes, fruit enzymes, flower enzymes, and so on (Neupane & Khadka, 2019; Singh *et al.*, 2019). It is a fermentation product.

One use of bio-enzymes is to accelerate the waste degradation process and improve environmental quality (Ardianita *et al.*, 2023). However, several studies have shown that the use of bio-enzymes (ecoenzymes) in fish farming environments does not show a different effect on reducing ammonia, nitrate, and nitrite and does not result in good fish growth (Kusumawati *et al.*, 2018; Bambang *et al.*, 2024). Based on the above description, this study was conducted to confirm previous studies by analyzing the growth performance of Nile fish and catfish fry cultivated with the addition or supplementation of bio-enzymes with different concentrations in the maintenance water.

MATERIALS AND METHODS

This research was conducted at the Hatchery, Bangka Belitung University, from June to August 2025. The samples used in this study were 100 Nile fish fry measuring 6-8 cm and catfish fry measuring 5-7 cm in each pond. Both types of samples were cultivated in concrete ponds measuring 2 m long, 1.5 m wide, and 0.6 m high, filled with water to a depth of 30 cm.

The research treatment consisted of two, namely cultivation media without added bio-enzymes and cultivation media added bio-enzymes. Bio-enzyme supplementation into the cultivation media was carried out before fish maintenance with doses of 500 ml and 1000 ml in each pond containing 900 L of water or treatments of 0.6 ml/L and 1.1 ml/L. The test sample fish fry were maintained in the research pond for 40 days by being fed pellets *ad satiation*, namely feeding until the fish were full and stopped eating.

Commented [H5]: is this pre treatment of reference?

Fish sample measurements were conducted on 10% of the population in each treatment. The research parameters observed in fish were absolute weight growth (g), absolute length growth (cm), daily weight growth rate (g/day), daily length growth rate (cm/day), specific weight growth rate (%), specific length growth rate (%), and survival (%). The water quality parameters observed were pH, temperature (°C), dissolved oxygen (DO) (ppm), total dissolved solids (TDS) (ppm), and electrical conductivity (EC) (mS/cm). The research data were analyzed descriptively and presented in tables and graphs.

RESULTS AND DISCUSSION

The results of the study showed that bio-enzyme supplementation into the pond compared to the control treatment without bio-enzyme can have an effect or no effect on the growth performance of fish fry. Bio-enzyme supplementation into Nile fish fry ponds contributed more to higher weight growth than length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation into catfish fry ponds resulted in higher length growth than weight growth. Supplementation of 500 ml of bio-enzyme resulted in the highest absolute weight growth of Nile fish fry, namely 16.95 g with a daily weight growth rate of 0.42 g/day and a specific weight growth rate of 3.35% and 100% survival. Supplementation of 500 ml of bio-enzyme resulted in the highest absolute length growth of catfish fry, namely 9.20 cm with a daily length growth rate of 0.23 cm/day and a specific length growth rate of 2.23% and 100% survival.

Growth of Nile fish

Supplementation of bio-enzymes into the cultivation medium, namely the maintenance water in the Nile fish fry pond resulted in higher weight growth of the fry at the end of the observation compared to the control treatment without bio-enzymes. However, the length growth of the Nile fish fry reared in the pond with bio-enzyme supplementation was lower than the control treatment without bio-enzymes (Fig. 1). The absolute growth, daily growth rate, and specific growth rate of the Nile fish fry were shown in Table 1. The research data also did not show any symptoms of disease in the fry reared, but there were 3 fish that died during maintenance in the 1000 ml bio-enzyme treatment, while in the other treatments no dead fry were found during maintenance (Table 1).




Fig. 1. Growth in weight and length of Nile fish fry

Commented [H6]: Replace figure and word of Nile fish to tilapia

Table 1. Calculation of growth and survival of Nile fish fry

Parameter	Nile fish fry		
	Control	Bio-enzyme	
		500 ml	1000 ml
Absolute weight gain (g)	11.94	16.95	16.47
Absolute length growth (cm)	17.75	12.98	10.20
Daily weight gain rate (g/day)	0.30	0.42	0.41
Daily length growth rate (cm/day)	0.44	0.32	0.26
Specific weight growth rate (%)	1.88	3.35	3.13
Specific length growth rate (%)	2.98	2.74	2.23
Survival rate (%)	100	100	97

Growth of Catfish

Supplementation of bio-enzymes into the culture medium, namely the maintenance water in the catfish fry pond, resulted in lower weight growth of the fry at the end of the observation compared to the control treatment without bio-enzymes. However, the length growth of the catfish fry reared in the pond with bio-enzyme supplementation was higher than the control treatment without bio-enzymes (Fig. 2). The absolute growth, daily growth rate, and specific growth rate of the catfish fry were shown in Table 1. The research data also showed no symptoms of disease and no dead fry were found during the maintenance (Table 2).




Fig. 2. Growth in weight and length of catfish fry

Table 2. Calculation of growth and survival of catfish fry

Parameter	Catfish fry		
	Control	Bio-enzyme	
		500 ml	1000 ml
Absolute weight gain (g)	15.48	15.40	13.28
Absolute length growth (cm)	7.55	9.20	7.85
Daily weight gain rate (g/day)	0.39	0.39	0.33
Daily length growth rate (cm/day)	0.19	0.23	0.20
Specific weight growth rate (%)	5.65	5.69	5.23
Specific length growth rate (%)	1.94	2.23	1.99
Survival rate (%)	100	100	100

Water quality as a medium for cultivating Nile fish and catfish fry

The results of water quality measurements used as a culture medium for Nile fish and catfish fry—pH, temperature, DO, TDS, and EC—showed that all parameters were dynamic. This indicates that water quality fluctuates during cultivation and does not reflect a relationship between bio-enzyme supplementation in the culture medium and water quality.

Water quality in the Nile fish fry rearing ponds showed a pH value that tended to be higher than that in the catfish fry ponds. The water temperature in the Nile fish fry rearing ponds was relatively lower than that in the catfish fry ponds. Dissolved oxygen in the water in the Nile fish fry ponds was relatively better than that in the catfish fry ponds. Total dissolved solids and electrical conductivity in the Nile fish and catfish fry control ponds were relatively higher at certain times and relatively lower at other times compared to the TDS in the bio-enzyme treatment ponds.

The water quality in the Nile fish fry pond had the highest pH value in the 500 ml bio-enzyme treatment, the highest temperature in the 1000 ml bio-enzyme treatment, the highest DO in the 500 ml bio-enzyme treatment, and the highest TDS and EC in the control treatment. The water quality in the catfish fry pond had the highest pH value in the 500 ml bio-enzyme treatment, the highest temperature in the control treatment, and the highest DO, TDS, and EC in the control treatment. The water quality conditions and comparison between the water quality in the Nile fish and catfish fry ponds were shown in Fig. 3 and Fig. 4.




Fig. 3. Water quality as a medium for Nile fish cultivation




Fig. 4. Water quality as a medium for catfish cultivation

Sikku *et al.* (2023) conducted a study on the application of 5-15 ml/L of ecoenzyme (bio-enzyme) in Nile fish fry (3 cm in length) cultivation pond maintained for 45 days. The absolute weight produced ranged from 0.98-1.76 g with a specific weight growth rate ranging from 2.33-4.19%. The water pH value during maintenance ranged from 7.6-9.1. When compared with the specific weight growth rate in this study which ranged from 3.13-3.35% (Table 1), then supplementation with a high concentration of bio-enzyme (15 ml/L) resulted in a better specific weight growth rate than the low concentration (0.6-1.1 ml/L) in this study. However, at a concentration of 5-10 ml/L in the study by **Sikku *et al.* (2023)** showed a lower value (2.33-2.98%) compared to the treatment in this study. Thus, there is no linear pattern that shows that the higher the concentration of bio-enzyme given, the higher the growth of Nile fish fry.

Bio-enzyme research has also been conducted on catfish fry cultivation. The results of **Kusumawati *et al.* (2018)** research show that the addition of bio-enzymes to catfish fry ponds maintained for 4 weeks (approximately 28 days) has a length of 5.9-8.0 cm with a weight of 2.01-4.32 g from a length of 5.1-5.8 cm and a weight of 1.12-1.22 g on day 0. Meanwhile, the results of measurements on the 26th day of this study have shown that the weight and length data were 8.76 g and 9.22 cm respectively in the control treatment; 8.14 g and 10.12 cm in the 500 ml bio-enzyme treatment (0.6 ml/L); and 7.90 g and 9.82 cm in the 1000 ml bio-enzyme treatment (1.1 ml/L) (Figure 4). This difference may be caused by the contribution of bio-enzymes in the research of **Kusumawati *et al.* (2018)** found lower pH and DO values, ranging from 5.85 to 6.89 and DO from 2.95 to 3.71 ppm in the fourth week of culture. This study, however, found pH values ranging from 8.33 to 9.08 and DO from 4.8 to 6.8 ppm. Based on SNI 6484.4:2014, water quality for catfish fry culture has a pH of 6.5 to 8 and DO > 3 ppm.

The description shows that specifically, the contribution of bio-enzyme supplementation to the growth performance of Nile fish fry was better than that of catfish fry (Table 1 and Table 2). This was in line with the statement by **Herliwati *et al.* (2024)** that the addition of bio-enzymes can improve water quality, which can impact the survival of catfish fry. However, the addition of bio-enzymes to the cultivation water medium only affected the growth of Nile fish and not catfish. Similar conditions were also found in the results of research by **Mado *et al.* (2023)** and **Natsir *et al.* (2024)** who found increased growth of Nile fish with the application of bio-enzymes to the cultivation water medium. However, according to **Kusumawati *et al.* (2018)** and **Bambang *et al.* (2024)**, catfish growth did not change with the same application. This is possible because catfish have a tolerance to less than ideal waters (**Hidayati *et al.*, 2021**), so that when water quality improves, its metabolism does not affect. Meanwhile, Nile fish experience increased metabolism when quality improves, resulting in better growth. Nile fish fry experience increased growth and survival when oxygen and ammonia levels are more appropriate (**Islami *et al.*, 2017**). When compared to the pH, DO, and TDS conditions in Fig. 3 and Fig. 4, it can be seen that supplementing bio-enzymes into the culture medium results in better water quality for Nile fish fry cultivation than for catfish fry. Furthermore, Fig. 3 and Fig. 4 also show that the total dissolved solids in catfish fry ponds were higher than in Nile fish fry ponds. High dissolved solids can hinder the growth performance of fish fry.

Rahman *et al.* (2020) explained that total dissolved solids are organic and inorganic substances dissolved in water that can be a factor causing water turbidity. According to **Damayanti (2014)**, increased TDS concentrations in water can originate from feed waste and accumulated metabolic waste such as fish feces. According to **Al Zamzami *et al.* (2019)**, the length of maintenance time can contribute to an increase in dissolved organic matter, one of which comes from feed waste and fish feces. Furthermore, various studies have revealed that the presence of nitrogen waste can produce ammonia, nitrate, and nitrite

compounds that can cause oxygen depletion, disease, and even death in fish (Munguti *et al.*, 2021; Li *et al.*, 2023; Mustafa *et al.*, 2024; Kurniawan *et al.*, 2025).

Relatively low bio-enzyme concentrations were considered ineffective in reducing waste contamination, particularly in catfish fry ponds. This is visually confirmed by the cloudier and more odorous water in catfish fry ponds compared to Nile fish fry ponds. This is thought to inhibit the growth of catfish fry in rearing ponds. Kusumawati *et al.* (2018) explained that administering bio-enzymes at low concentrations does not significantly reduce ammonia and nitrite concentrations. This contributed to the slow growth of catfish fry.

Higher bio-enzyme concentrations, duration, and frequency of bio-enzyme administration need to be tested for their effectiveness in improving the quality of aquaculture water and its effect on optimal growth for fish fry. This is in line with Putri *et al.* (2024) who stated that the longer the time and the higher the eco-enzyme dosage, the higher the effectiveness in reducing waste contamination as indicated by the biological oxygen demand (COD) and total suspended solids (TSS) in the environment. Yatma *et al.* (2024) also revealed that increasing the eco-enzyme dosage used was able to significantly reduce the dissolved solids (TDS) content and improve water quality. Furthermore, water quality and nutrient contribute to fish morphological and its growth (Serdiati *et al.*, 2024; Mustikasari *et al.*, 2020).

Discussion about Agribusiness Perspective of Bio-Enzyme Supplementation for Aquaculture

From an agribusiness perspective, bio-enzyme supplementation in aquaculture presents a high-potential growth sector that aligns with key trends: sustainable food production, environmental regulation, and consumer preference for clean-label seafood. Strategic investment in research and development, production, and farmer outreach can unlock long-term profitability and ecological benefits.

Perspective in agribusiness opportunities, including market potential, value addition, product development and diversification, challenges from an agribusiness viewpoint, and investment and agribusiness models. In Indonesia (and many other countries) Nile fish and catfish are major freshwater/pond-aquaculture species. Grower demand is strong for both local consumption and export. A value proposition; higher growth/lower feed cost/improved survival is very attractive to fish farmers, in particular Nile fish and catfish. Use the enzyme supplementation in aquaculture can improve output and profit. In other hand, enzyme supplementation can reduce feed costs (better feed conversion ratio or FCR), improves animal health and survival rates — leading to higher yields and profits, reduces environmental impact, enhancing the farm's market reputation and compliance with green regulations (Varshini & Gayathri, 2023).

Commented [H7]: Make agribusiness analysis deeper

Using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming. The benefits using bio-enzymes in aquaculture from an agribusiness perspective:

Improved Productivity and Profitability

Enhanced Feed Conversion Ratio (FCR) where bioenzymes help fish and shrimp digest feed more efficiently, reducing feed costs — a major operational expense. Faster growth rates where enzymes promote better nutrient absorption, accelerating growth and reducing time to market. Lower mortality rates where enzymes improve gut health and immunity, reducing disease outbreaks.

Reduction in Input Costs

Less dependency on antibiotics and chemicals where enzymes can reduce disease pressure naturally. Better water quality management where bioenzymes help degrade organic waste and ammonia, reducing the need for chemical water treatments.

Sustainable Branding & Market Access

Eco-friendly production where enzyme use aligns with growing consumer demand for sustainable and antibiotic-free aquaculture. Certification ready where farms using bioenzymes can more easily meet organic or eco-certification standards.

Waste Reduction and Circular Economy

Utilization of agricultural byproducts where some bioenzymes are produced using agri-waste (e.g., fruit peels, crop residues), supporting circular bioeconomy models. Efficient waste breakdown where help manage sludge and organic waste, reducing environmental impact.

New Revenue Streams

Local production of bioenzymes where farmers or agri-entrepreneurs can engage in low-cost production of bioenzymes using fermentation, creating micro-enterprises. Diversification for agro-industries where companies in biotech, agriculture, or food waste management can expand into aquaculture support services.

Regulatory and Environmental Compliance

Supports green policies where use of bioenzymes aligns with government and international sustainability goals in aquaculture. Reduces pollution where better water quality means less eutrophication and contamination of surrounding ecosystems.

CONCLUSION

The addition or supplementation of bio-enzymes to Nile fish and catfish fry ponds produced different contributions. Bio-enzyme supplementation to Nile fish fry ponds compared to the control treatment without bio-enzymes showed a higher contribution to weight growth than to length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation to catfish fry ponds compared to the control without bio-enzymes showed a higher contribution to length growth than to weight growth. The results of water quality measurements as a medium for cultivating Nile fish and catfish fry, namely pH, temperature, DO, TDS, and EC showed that all parameters were dynamic. This means that water quality fluctuates during cultivation and does not reflect the relationship between bio-enzyme supplementation to the cultivation medium and water quality. In agribusiness perspective, using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming.

Commented [H8]: Make sure the conclusion

ACKNOWLEDGMENT

The authors would like to thank Bangka Belitung University for supporting and funding this research and publication through Grant of Penelitian Unggulan 2025 with grant number 1116/UN50/M/PP/2025.

CONFLICTS OF INTEREST

All the authors declare no conflict of interest.

FUNDING

This research and publication were supported and funded through the Grant of Penelitian Unggulan 2025 grant number: 1116/UN50/M/PP/2025.

REFERENCES

- Ajayi, A. O. and Okoh, A. I. (2014). Bacteriological study of pond water for aquaculture purposes. *Journal of Food Agriculture and Environment*, 12(2): 1260-1265.
- Al Zamzami, I. M.; Sari, A. H. W. and Perwira, I. Y. (2019). Fluktuasi bahan organik dan residu terlarut dalam budidaya ikan koi di Tulungagung Jawa Timur. *Current Trends in Aquatic Science*, 2(1): 79-84.
- Ardianita, N.; Andriani, R. and Maya, N. P. (2023). Efektivitas ekoenzim terhadap kelulushidupan ikan lele (*Clarias* sp.) pada media yang tercemar limbah batik. *Biology Natural Resources Journal*, 2(1): 24-27.

- Bambang, B.; Umar, N. A. and Indrawati, E.** (2024). Pengaruh ekoenzim dosis berbeda terhadap perbaikan kualitas air budidaya dalam pembesaran ikan lele mutiara *Clarias* sp. *Journal of Aquaculture and Environment*, 6(2): 84-89.
- Damayanti, H. O.** (2014). Tinjauan kualitas dan dampak ekonomi konsentrasi total dissolved solid (TDS) air di area pertambakan Desa Bulumanis Kidul. *Jurnal Litbang: Media Informasi Penelitian, Pengembangan dan IPTEK*, 10(2): 103-113.
- Ducatelle, R.; Goossens, E.; De Meyer, F.; Eeckhaut, V.; Antonissen, G.; Haesebrouck, F. and Van Immerseel, F.** (2018). Biomarkers for monitoring intestinal health in poultry: Present status and future perspectives. *Veterinary Research*, 49: 43.
- Galintin, O.; Rasit, N. and Hamzah, S.** (2021). Production and characterization of eco enzyme produced from fruit and vegetable wastes and its influence on the aquaculture sludge. *Biointerface Research in Applied Chemistry*, 11(3): 10205-10214.
- Herliwati, H.; Rahman, M.; Ahmadi, A. and Tony, F.** (2024). Variations in the addition of eco enzyme in feed on the survival of siamese catfish (*Pangasius hyphthalmus*) seeds raised in aquariums. *Fish Scientiae*, 14(2): 145-154.
- Hidayati, S. N.; Laili, S. and Santoso, H.** (2021). Pengaruh kualitas air kolam terpal terhadap pertumbuhan ikan lele dumbo (*Clarias gariepinus*). *Jurnal Ilmiah Biosaintropis*, 6(2): 19-25.
- Islami, A. N.; Hasan, Z. and Anna, Z.** (2017). Pengaruh perbedaan siphonisasi dan aerasi terhadap kualitas air, pertumbuhan, dan kelangsungan hidup pada budidaya ikan nila (*Oreochromis niloticus*) stadia benih. *Jurnal Perikanan Kelautan*, 8(1): 73-82.
- Koniyo, Y.** (2020). Analisis kualitas air pada lokasi budidaya ikan air tawar di Kecamatan Suwawa Tengah. *Jurnal Technopreneur*, 8(1): 52-58.
- Kulla, O. L. S.; Yuliana, E. and Supriyono, E.** (2020). Analisis kualitas air dan kualitas lingkungan untuk budidaya ikan di Danau Laimadat, Nusa Tenggara Timur. *Pelagicus*, 1(3): 135-144.
- Kurniawan, A.; Mustikasari, D.; Kurniawan, A.; Muntoro, M.; Setiadi, J. and Kholishah, N. A.** (2024). The preliminary study about physico-chemical property of bio-enzyme produced from orange fruits waste treated with different concentrations of probiotic. *Jurnal Ilmu Lingkungan*, 22(4), 861-867.
- Kurniawan, A.; Mustikasari, D.; Muntoro.; Setiadi, J.; Wulandari, U. A.; Pratama, V.; Fadia, S.; Maheswari, D. A.; Fitri, C. and Iqbal, D. H.** (2025). Effectiveness

of decaying fruits as a carbon source on fish productivity and water quality improvement in aquaculture: A narrative review. *Egyptian Journal of Aquatic Biology & Fisheries*, 29(4): 2459-2481.

Kusumawati, A. A.; Suprpto, D. and Haeruddin, H. (2018). Pengaruh ekoenzim terhadap kualitas air dalam pembesaran ikan lele (*Clarias gariepinus*), 7(4): 307-314.

Li, H.; Cui, Z.; Cui, H.; Bai, Y.; Yin, Z. and Qu, K. (2023). Hazardous substances and removal in review. *Aquaculture*, 569: 739399.

Mado, A. M.; Umar, N. A. and Aqmal, A. (2023). Pengaruh eco-enzym terhadap kualitas air pada sistem budidaya akuaponik ikan nila *Oreochromis niloticus* dan tanaman sawi *Brassica juncea*. *Journal of Aquaculture and Environment*, 5(2): 66-72.

Munguti, J. M.; Sendi, J. G.; Obiero, K. O.; Ogello, E. O.; Kyule, D. N.; Liti, D. M. and Musalia, L. M. (2021). Aqua-feed wastes: Impact on natural systems and practical mitigations-A review. *Journal of Agricultural Science*, 13(1): 111-121.

Mustafa, S. A.; Al-Rudainy, A. J. and Salman, N. M. (2024). Effect of environmental pollutants on fish health: An overview. *Egyptian Journal of Aquatic Research*, 50(2): 225-233.

Mustikasari, D.; Suryaningsih, S. and Nuryanto, A. (2020). Morphological variation of blue panchax (*Aplocheilichthys panchax*) lives in different habitat assessed using truss morphometric. *Biosaintifika: Journal of Biology & Biology Education*, 12(3): 399-407.

Nalle, M. M. D.; Kamiasi, Y. and Edo, S. I. (2021). Budidaya ikan lele dan ikan nila oleh Kelompok Tani Tunfeu Desa Ponain Kecamatan Amarasi Kabupaten Kupang di masa Covid-19. *Jurnal Pengabdian Perikanan Indonesia*, 1(2): 151-156.

Natsir, N. I.; Jayadi, J. and Hadijah, S. (2024). Aplikasi eco-enzyme terhadap pertumbuhan dan tingkat kelangsungan hidup ikan nila (*Oreochromis niloticus*). *Jurnal Akuakultur Nusantara*, 1(2): 96-106.

Neupane, K. and Khadka, R. (2019). Production of garbage enzyme from different fruit and vegetable wastes and evaluation of its enzymatic and antimicrobial efficacy. *Tribhuvan University Journal of Microbiology*, 6: 113-118.

Putri, M. C.; Widiarini, R. and Ramadanintyas, K. N. (2024). Effectiveness of eco enzymes in reducing chemical parameters of tofu factory liquid waste in Klumutan Village, Saradan District. *Journal of Social Research*, 3(9): 1-10.

- Rahman, M. F. I.; Wibisana, H. and Zainab, S. (2020).** Analisa dan pemetaan total padatan terlarut di pesisir Pantai Pasuruan dengan citra satelit Terra Modis. *Journal of Civil Engineering*, 16(2): 144-156.
- Rasit, N.; Fern, L. H. and Ghani, W. A. W. A. K. (2019).** Production and characterization of eco enzyme produced from tomato and orange wastes and its influence on the aquaculture sludge. *International Journal of Civil Engineering and Technology*, 10(3): 967-980.
- Serdiati, N.; Islamy, R. A.; Mamat, N. B.; Hasan, V. and Valen, F. S. (2024).** Nutritional value of alligator weed (*Alternanthera philoxeroides*) and its application for herbivorous aquaculture feed. *International Journal of Agriculture and Biosciences*, 13(3): 318-324.
- Siegers, W. H.; Prayitno, Y. and Sari, A. (2019).** Pengaruh kualitas air terhadap pertumbuhan ikan nila nirwana (*Oreochromis* sp.) pada tambak payau. *The Journal of Fisheries Development*, 3(2): 95-104.
- Sikku, E. S.; Umar, N. A. and Indrawati, E. (2023).** Penggunaan eco-enzim dengan dosis berbeda pada teknologi akuaponik sederhana untuk optimalisasi pertumbuhan ikan nila *Oreochromis niloticus*. *Journal of Aquaculture and Environment*, 5(2): 60-65.
- Singh, H.; Prasad, A. and Nal, K. V. (2019).** Study and analysis of bio enzymes based natural products. *International Journal of Engineering Applied Sciences and Technology*, 4(8): 205-207.
- Tumwesigye, Z.; Tumwesigye, W.; Opio, F.; Kemigabo, C. and Mujuni, B. (2022).** The effect of water quality on aquaculture productivity in Ibanda District, Uganda. *Aquaculture Journal*, 2(1): 23-36.
- Varshini, B. and Gayathrif, V. (2023).** Role of eco-enzymes in sustainable development. *Nature Environment and Pollution Technology*, 22(3): 1299-1310.
- Yatma, M. A.; Hamzani, S. and Zubaidah, T. (2024).** The effect of eco enzyme on improving river water quality based on turbidity, temperature, pH, and TDS parameters. *Global Health & Environmental Perspectives*, 1(3): 164-168.



Growth Performance of Tilapia and Catfish Fry Cultivated with Bio-Enzyme Supplementation and Its Agribusiness Perspective

Muntoro¹, Andri Kurniawan^{2*}, Ardiansyah Kurniawan³, Ufi Ayu Wulandari³, Didin Erlan³, Venisa Pratama³, Syafiq Muhammad Syakir³, Danella Aisah Maheswari², Safitri Fadiah², Cahya Fitri²

¹Department of Agribusiness, Faculty of Agriculture, Fisheries and Marine, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

²Department of Biology, Faculty of Science and Engineering, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

³Department of Aquaculture, Faculty of Agriculture, Fisheries and Marine, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

*Corresponding Author: andri_pangkal@yahoo.co.id

ARTICLE INFO

Article History:

Received:

Accepted:

Online:

Keywords:

Aquaculture,
Bio-Enzyme;
Effectiveness; Growth
Rate; Fish Fry

ABSTRACT

Tilapia and catfish are freshwater fish commodities that are widely cultivated by the community because of their relatively easy maintenance, low cost, and short time. The quality of the cultivation water environment is an important factor and influences the growth performance and life of these commodities. This study aimed to analyze the growth performance of tilapia and catfish fry cultivated with the addition or supplementation of bio-enzymes with different concentrations in the maintenance water. This study used an experimental method with three treatments: control (without bio-enzymes), 500 ml/900 L or 0.6 ml/L bio-enzymes, and 1000 ml/900 L or 1.1 ml/L bio-enzymes. The results illustrate that bio-enzyme supplementation into tilapia fry ponds compared to the control treatment without bio-enzymes showed a higher contribution to weight growth than length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation into catfish fry ponds compared to the control without bio-enzymes showed a higher contribution to length growth than weight growth. Supplementation with 500 ml of bio-enzyme resulted in the highest absolute weight growth of tilapia fry, which was 16.95 g with a daily weight growth rate of 0.42 g/day and a specific weight growth rate of 3.35% and 100% survival rate. Supplementation with 500 ml of bio-enzyme resulted in the highest absolute length growth of catfish fry, which was 9.20 cm with a daily length growth rate of 0.23 cm/day and a specific length growth rate of 2.23% and 100% survival rate. In agribusiness perspective, only tilapia in consumption size (at 135 days old) treated with bio-enzymes (500 ml and 1000 ml) were recommended for sale, particularly in consumption size. Meanwhile, for catfish, only in fingerlings phase were recommended for selling with treatment bio-enzyme 500 ml.

INTRODUCTION

Fish farming, known as aquaculture, is one of the leading fish food sources with the fastest growth in productivity worldwide (Sumaila *et al.*, 2022; Serra *et al.*, 2024). Aquaculture stands as a vital provider of animal protein essential for human nutrition, with fish exhibiting rapid growth and offering optimal calorie-protein ratios for human consumption (Melenchón, *et al.*, 2022). Freshwater fish farming has been widely developed by communities, both on a household and large scale (Mzula *et al.*, 2021; Angermayr *et al.*, 2023). Among the freshwater fish commonly cultivated by the community are tilapia and catfish (Febrianty, 2021; Nalle *et al.*, 2021).

The development of fish farming productivity is greatly influenced by water quality, which significantly supports fish growth performance (Ajayi & Okoh, 2014; Tumwesigye *et al.*, 2022). Water quality, manifested in the form of physical, chemical, and biological parameters, affects the entire life cycle, including survival, growth, and reproduction (Siegers *et al.*, 2019; Koniyo, 2020; Kulla *et al.*, 2020).

Aquaculture water quality is generally contaminated by organic matter from leftover feed and metabolic waste in the form of fish feces. One effort to maintain and improve water quality in fish farming is the addition of enzymes to the culture environment.

Bio-enzymes or Bioz are fermented products made from fruits or vegetables that are widely used for various purposes due to their probiotic microorganism content and bioactive compounds (Ducatelle *et al.*, 2018). It is produced in 3 months fermentation process with ratio 1 : 3: 10 for sugar : waste : water (Rasit *et al.*, 2019; Galintin *et al.*, 2021; Kurniawan *et al.*, 2024). Bio-enzymes are also known as ecoenzymes, garbage enzymes, terrazymes, fruit enzymes, flower enzymes, and so on (Neupane & Khadka, 2019; Singh *et al.*, 2019). It is a fermentation product

One use of bio-enzymes is to accelerate the waste degradation process and improve environmental quality (Ardianita *et al.*, 2023). However, several studies have shown that the use of bio-enzymes (ecoenzymes) in fish farming environments does not show a different effect on reducing ammonia, nitrate, and nitrite and does not result in good fish growth (Kusumawati *et al.*, 2018; Bambang *et al.*, 2024). Based on the above description, this study was conducted to confirm previous studies by analyzing the growth performance of tilapia and catfish fry cultivated with the addition or supplementation of bio-enzymes with different concentrations in the maintenance water.

MATERIALS AND METHODS

This research was conducted at the Hatchery, Bangka Belitung University, from June to August 2025. The samples used in this study were 100 tilapia fry measuring 6-8 cm and

catfish fry measuring 5-7 cm in each pond. Both types of samples were cultivated in concrete ponds measuring 2 m long, 1.5 m wide, and 0.6 m high, filled with water to a depth of 30 cm.

The research treatment consisted of two, namely cultivation media without added bio-enzymes and cultivation media added bio-enzymes. Based on pre research, effectiveness of bio-enzyme supplementation into the cultivation media was carried out before fish maintenance with doses of 500 ml and 1000 ml in each pond containing 900 L of water or treatments of 0.6 ml/L and 1.1 ml/L. The test sample fish fry were maintained in the research pond for 40 days by being fed pellets *ad satiation*, namely feeding until the fish were full and stopped eating.

Fish sample measurements were conducted on 10% of the population in each treatment. The research parameters observed in fish were absolute weight growth (g), absolute length growth (cm), daily weight growth rate (g/day), daily length growth rate (cm/day), specific weight growth rate (%), specific length growth rate (%), and survival (%). The water quality parameters observed were pH, temperature (°C), dissolved oxygen (DO) (ppm), total dissolved solids (TDS) (ppm), and electrical conductivity (EC) (mS/cm). The research data were analyzed descriptively and presented in tables and graphs.

RESULTS AND DISCUSSION

The results of the study showed that bio-enzyme supplementation into the pond compared to the control treatment without bio-enzyme can have an effect or no effect on the growth performance of fish fry. Bio-enzyme supplementation into tilapia fry ponds contributed more to higher weight growth than length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation into catfish fry ponds resulted in higher length growth than weight growth. Supplementation of 500 ml of bio-enzyme resulted in the highest absolute weight growth of tilapia fry, namely 16.95 g with a daily weight growth rate of 0.42 g/day and a specific weight growth rate of 3.35% and 100% survival. Supplementation of 500 ml of bio-enzyme resulted in the highest absolute length growth of catfish fry, namely 9.20 cm with a daily length growth rate of 0.23 cm/day and a specific length growth rate of 2.23% and 100% survival.

Growth of Tilapia

Supplementation of bio-enzymes into the cultivation medium, namely the maintenance water in the tilapia fry pond resulted in higher weight growth of the fry at the end of the observation compared to the control treatment without bio-enzymes. However, the length growth of the tilapia fry reared in the pond with bio-enzyme supplementation was lower than the control treatment without bio-enzymes (Fig. 1). The absolute growth, daily growth rate, and specific growth rate of the tilapia fry were shown in Table 1. The

research data also did not show any symptoms of disease in the fry reared, but there were 3 fish that died during maintenance in the 1000 ml bio-enzyme treatment, while in the other treatments no dead fry were found during maintenance (Table 1).




Fig. 1. Growth in weight and length of tilapia fry

Table 1. Calculation of growth and survival of tilapia fry

Parameter	Tilapia Fry		
	Control	Bio-enzyme	
		500 ml	1000 ml
Absolute weight gain (g)	11.94	16.95	16.47
Absolute length growth (cm)	17.75	12.98	10.20
Daily weight gain rate (g/day)	0.30	0.42	0.41
Daily length growth rate (cm/day)	0.44	0.32	0.26
Specific weight growth rate (%)	1.88	3.35	3.13
Specific length growth rate (%)	2.98	2.74	2.23
Survival rate (%)	100	100	97

Growth of Catfish

Supplementation of bio-enzymes into the culture medium, namely the maintenance water in the catfish fry pond, resulted in lower weight growth of the fry at the end of the observation compared to the control treatment without bio-enzymes. However, the length growth of the catfish fry reared in the pond with bio-enzyme supplementation was higher than the control treatment without bio-enzymes (Fig. 2). The absolute growth, daily growth rate, and specific growth rate of the catfish fry were shown in Table 1. The research data also showed no symptoms of disease and no dead fry were found during the maintenance (Table 2).




Fig. 2. Growth in weight and length of catfish fry

Table 2. Calculation of growth and survival of catfish fry

Parameter	Catfish Fry		
	Control	Bio-enzyme	
		500 ml	1000 ml
Absolute weight gain (g)	15.48	15.40	13.28
Absolute length growth (cm)	7.55	9.20	7.85
Daily weight gain rate (g/day)	0.39	0.39	0.33
Daily length growth rate (cm/day)	0.19	0.23	0.20
Specific weight growth rate (%)	5.65	5.69	5.23
Specific length growth rate (%)	1.94	2.23	1.99
Survival rate (%)	100	100	100

Water quality as a medium for cultivating tilapia and catfish fry

The results of water quality measurements used as a culture medium for Nile fish and catfish fry—pH, temperature, DO, TDS, and EC—showed that all parameters were dynamic. This indicates that water quality fluctuates during cultivation and does not reflect a relationship between bio-enzyme supplementation in the culture medium and water quality.

Water quality in the tilapia fry rearing ponds showed a pH value that tended to be higher than that in the catfish fry ponds. The water temperature in the tilapia fry rearing ponds was relatively lower than that in the catfish fry ponds. Dissolved oxygen in the water in the tilapia fry ponds was relatively better than that in the catfish fry ponds. Total dissolved solids and electrical conductivity in the tilapia and catfish fry control ponds were relatively higher at certain times and relatively lower at other times compared to the TDS in the bio-enzyme treatment ponds.

The water quality in the tilapia fry pond had the highest pH value in the 500 ml bio-enzyme treatment, the highest temperature in the 1000 ml bio-enzyme treatment, the highest DO in the 500 ml bio-enzyme treatment, and the highest TDS and EC in the control treatment. The water quality in the catfish fry pond had the highest pH value in the 500 ml bio-enzyme treatment, the highest temperature in the control treatment, and the highest DO, TDS, and EC in the control treatment. The water quality conditions and comparison between the water quality in the tilapia and catfish fry ponds were shown in Fig. 3 and Fig. 4.




Fig. 3. Water quality as a medium for tilapia cultivation




Fig. 4. Water quality as a medium for catfish cultivation

Sikku *et al.* (2023) conducted a study on the application of 5-15 ml/L of ecoenzyme (bio-enzyme) in tilapia fry (3 cm in length) cultivation pond maintained for 45 days. The absolute weight produced ranged from 0.98-1.76 g with a specific weight growth rate ranging from 2.33-4.19%. The water pH value during maintenance ranged from 7.6-9.1. When compared with the specific weight growth rate in this study which ranged from 3.13-3.35% (Table 1), then supplementation with a high concentration of bio-enzyme (15 ml/L) resulted in a better specific weight growth rate than the low concentration (0.6-1.1 ml/L) in this study. However, at a concentration of 5-10 ml/L in the study by **Sikku *et al.* (2023)** showed a lower value (2.33-2.98%) compared to the treatment in this study. Thus, there is no linear pattern that shows that the higher the concentration of bio-enzyme given, the higher the growth of tilapia fry.

Bio-enzyme research has also been conducted on catfish fry cultivation. The results of **Kusumawati *et al.* (2018)** research show that the addition of bio-enzymes to catfish fry ponds maintained for 4 weeks (approximately 28 days) has a length of 5.9-8.0 cm with a weight of 2.01-4.32 g from a length of 5.1-5.8 cm and a weight of 1.12-1.22 g on day 0. Meanwhile, the results of measurements on the 26th day of this study have shown that the weight and length data were 8.76 g and 9.22 cm respectively in the control treatment; 8.14 g and 10.12 cm in the 500 ml bio-enzyme treatment (0.6 ml/L); and 7.90 g and 9.82 cm in the 1000 ml bio-enzyme treatment (1.1 ml/L) (Figure 4). This difference may be caused by the contribution of bio-enzymes in the research of **Kusumawati *et al.* (2018)** found lower pH and DO values, ranging from 5.85 to 6.89 and DO from 2.95 to 3.71 ppm in the fourth week of culture. This study, however, found pH values ranging from 8.33 to 9.08 and DO from 4.8 to 6.8 ppm. Based on SNI 6484.4:2014, water quality for catfish fry culture has a pH of 6.5 to 8 and DO > 3 ppm.

The description shows that specifically, the contribution of bio-enzyme supplementation to the growth performance of tilapia fry was better than that of catfish fry (Table 1 and Table 2). This was in line with the statement by **Herliwati *et al.* (2024)** that the addition of bio-enzymes can improve water quality, which can impact the survival of catfish fry. However, the addition of bio-enzymes to the cultivation water medium only affected the growth of tilapia and not catfish. Similar conditions were also found in the results of research by **Mado *et al.* (2023)** and **Natsir *et al.* (2024)** who found increased growth of tilapia with the application of bio-enzymes to the cultivation water medium. However, according to **Kusumawati *et al.* (2018)** and **Bambang *et al.* (2024)**, catfish growth did not change with the same application. This is possible because catfish have a tolerance to less than ideal waters (**Hidayati *et al.*, 2021**), so that when water quality improves, its metabolism does not affect. Meanwhile, tilapia experience increased metabolism when quality improves, resulting in better growth. Tilapia fry experience increased growth and survival when oxygen and ammonia levels are more appropriate (**Islami *et al.*, 2017**). When compared to the pH, DO, and TDS conditions in Fig. 3 and Fig. 4, it can be seen that supplementing bio-enzymes into the culture medium results in better water quality for tilapia fry cultivation than for catfish fry. Furthermore, Fig. 3 and Fig. 4 also show that the total dissolved solids in catfish fry ponds were higher than in tilapia fry ponds. High dissolved solids can hinder the growth performance of fish fry.

Rahman *et al.* (2020) explained that total dissolved solids are organic and inorganic substances dissolved in water that can be a factor causing water turbidity. According to **Damayanti (2014)**, increased TDS concentrations in water can originate from feed waste and accumulated metabolic waste such as fish feces. According to **Al Zamzami *et al.* (2019)**, the length of maintenance time can contribute to an increase in dissolved organic matter, one of which comes from feed waste and fish feces. Furthermore, various studies have revealed that the presence of nitrogen waste can produce ammonia, nitrate, and nitrite

compounds that can cause oxygen depletion, disease, and even death in fish (**Munguti *et al.*, 2021; Li *et al.*, 2023; Mustafa *et al.*, 2024; Kurniawan *et al.*, 2025**).

Relatively low bio-enzyme concentrations were considered ineffective in reducing waste contamination, particularly in catfish fry ponds. This is visually confirmed by the cloudier and more odorous water in catfish fry ponds compared to tilapia fry ponds. This is thought to inhibit the growth of catfish fry in rearing ponds. **Kusumawati *et al.* (2018)** explained that administering bio-enzymes at low concentrations does not significantly reduce ammonia and nitrite concentrations. This contributed to the slow growth of catfish fry.

Higher bio-enzyme concentrations, duration, and frequency of bio-enzyme administration need to be tested for their effectiveness in improving the quality of aquaculture water and its effect on optimal growth for fish fry. This is in line with **Putri *et al.* (2024)** who stated that the longer the time and the higher the eco-enzyme dosage, the higher the effectiveness in reducing waste contamination as indicated by the biological oxygen demand (COD) and total suspended solids (TSS) in the environment. **Yatma *et al.* (2024)** also revealed that increasing the eco-enzyme dosage used was able to significantly reduce the dissolved solids (TDS) content and improve water quality. Furthermore, water quality and nutrient contribute to fish morphological and its growth (**Serdiati *et al.*, 2024; Mustikasari *et al.*, 2020**).

Discussion about Agribusiness Perspective of Bio-Enzyme Supplementation for Aquaculture

From an agribusiness perspective, bio-enzyme supplementation in aquaculture presents a high-potential growth sector that aligns with key trends: sustainable food production, environmental regulation, and consumer preference for clean-label seafood. Strategic investment in research and development, production, and farmer outreach can unlock long-term profitability and ecological benefits.

Perspective in agribusiness opportunities, including market potential, value addition, product development and diversification, challenges from an agribusiness viewpoint, and investment and agribusiness models. In Indonesia (and many other countries) tilapia and catfish are major freshwater/pond-aquaculture species. Grower demand is strong for both local consumption and export. A value proposition; higher growth/lower feed cost/improved survival is very attractive to fish farmers, in particular tilapia and catfish. Use the enzyme supplementation in aquaculture can improve output and profit. In other hand, enzyme supplementation can reduce feed costs (better feed conversion ratio or FCR), improves animal health and survival rates — leading to higher yields and profits, reduces environmental impact, enhancing the farm's market reputation and compliance with green regulations (**Varshini & Gayathri, 2023**).

Using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming. The benefits using bio-enzymes in aquaculture from an agribusiness perspective:

Improved Productivity and Profitability

Enhanced Feed Conversion Ratio (FCR) where bioenzymes help fish and shrimp digest feed more efficiently, reducing feed costs — a major operational expense. Faster growth rates where enzymes promote better nutrient absorption, accelerating growth and reducing time to market. Lower mortality rates where enzymes improve gut health and immunity, reducing disease outbreaks.

Profitability of this research result can be calculated with assumption referring to Table 1 and Table 2. Tilapia and catfish fish fry were cultivated 45 days for fingerlings size and 135 days for consumption size with survival rate 100%. The potential economic value of selling of fish fry and consumption sizes was shown in Table 3.

The economic value of selling tilapia at the fingerling phase or consumption size, as presented in Table 3, indicated that only tilapia treated with bio-enzymes were recommended for sale. The potential of selling price from bio-enzyme-supplemented fish was higher than normal price at local market. Consumption size with treatment bio-enzyme 500 ml had potential of selling price 52,000 IDR and bio-enzyme 1000 ml had potential of selling price 47,565 IDR compared price at local market 35,000 IDR. Meanwhile, selling of fingerlings phase of bio-enzyme-supplemented tilapia was not recommended due to the size of fish fry treated with bio-enzymes was not better than the control.

For catfish, the control and bio-enzyme treatment of 500 ml and 1000 ml of enzyme did not show significant differences. Even though, fingerlings phase (at 45 days old) treated with bio-enzyme 500 ml had potential of selling price 495 IDR, higher than price at local market (300 IDR). Thus, selling of fingerlings phase of bio-enzyme-supplemented catfish (500 ml bio-enzyme) was recommended.

Weight of catfish fry in the control treatment was bigger than the bio-enzyme treatment so that the addition of bio-enzymes had no effect on fry growth. As well as the weight of the fish, the addition of bioenzymes did not result in significant weight gain compared to the control treatment. Therefore, selling of fingerlings phase and also consumption size of bio-enzyme-treated fish were not recommended.

Table 3. Economic value of tilapia and catfish in fingerlings and consumption size

Fish (Phase or Size)	Comparing Between Control and Bio-enzym Supplementation (500 ml and 1000 ml)		Price (IDR) in local market in Bangka, Indonesia	Potential of Selling Price (IDR)	Selling Recommendation of Bio-enzyme- supplemented Fish
	Weight (g)	Length (cm)			
Tilapia					
Fingerlings phase (at 45 days old)	5.01 x 100 fishes	Control > Treatment	200/fish	No	Not recommended for selling in fingerlings phase
	4.53 x 100 fishes				
Consumption size (at 135 days old)	15.03 x 100 fishes	-	35,000/kg (35/g)	52,000	Recommended for selling in consumption size (treatment bio-enzyme 500 ml)
	13.59 x 100 fishes			47,565	Recommended for selling in consumption size (treatment bio-enzyme 1000 ml)
Catfish					
Fingerlings phase (at 45 days old)	Control > Treatment	1.65	300/fish	495	Recommended for selling in consumption size (treatment bio-enzyme 500 ml)
		0.3		90	Not recommended for selling in fingerlings phase
Consumption size (at 135 days old)	-	4.95 x 100 fishes	23,000/kg (23/g)	11,385	Not recommended for selling in consumption size
		0.9 x 100 fishes		2,070	

Reduction in Input Costs

Less dependency on antibiotics and chemicals where enzymes can reduce disease pressure naturally. Better water quality management where bioenzymes help degrade organic waste and ammonia, reducing the need for chemical water treatments.

Sustainable Branding & Market Access

Eco-friendly production where enzyme use aligns with growing consumer demand for sustainable and antibiotic-free aquaculture. Certification ready where farms using bioenzymes can more easily meet organic or eco-certification standards.

Waste Reduction and Circular Economy

Utilization of agricultural byproducts where some bioenzymes are produced using agri-waste (e.g., fruit peels, crop residues), supporting circular bioeconomy models. Efficient waste breakdown where help manage sludge and organic waste, reducing environmental impact.

New Revenue Streams

Local production of bioenzymes where farmers or agri-entrepreneurs can engage in low-cost production of bioenzymes using fermentation, creating micro-enterprises. Diversification for agro-industries where companies in biotech, agriculture, or food waste management can expand into aquaculture support services.

Regulatory and Environmental Compliance

Supports green policies where use of bioenzymes aligns with government and international sustainability goals in aquaculture. Reduces pollution where better water quality means less eutrophication and contamination of surrounding ecosystems.

CONCLUSION

The addition or supplementation of bio-enzymes to tilapia and catfish fry ponds produced different contributions. Bio-enzyme supplementation to tilapia fry ponds compared to the control treatment without bio-enzymes showed a higher contribution to weight growth than to length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation to catfish fry ponds compared to the control without bio-enzymes showed a higher contribution to length growth than to weight growth. The results of water quality measurements as a medium for cultivating tilapia and catfish fry, namely pH, temperature, DO, TDS, and EC showed that all parameters were dynamic. This means that water quality fluctuates during cultivation and does not reflect the relationship between bio-enzyme supplementation to the cultivation medium and water quality. In agribusiness

perspective, using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming. However, based on the economic value, only tilapia in consumption size (at 135 days old) treated with bio-enzymes (500 ml and 1000 ml) were recommended for sale, particularly in consumption size. Meanwhile, for catfish, only in fingerlings phase were recommended for selling with treatment bio-enzyme 500 ml.

ACKNOWLEDGMENT

The authors would like to thank Bangka Belitung University for supporting and funding this research and publication through Grant of Penelitian Unggulan 2025 with grant number 1116/UN50/M/PP/2025.

CONFLICTS OF INTEREST

All the authors declare no conflict of interest.

FUNDING

This research and publication were supported and funded through the Grant of Penelitian Unggulan 2025 grant number 1116/UN50/M/PP/2025.

REFERENCES

- Ajayi, A. O. and Okoh, A. I. (2014). Bacteriological study of pond water for aquaculture purposes. *Journal of Food Agriculture and Environment*, 12(2): 1260-1265.
- Angermayr, G.; Palacio, A. and Chaminade, C. (2023). Small-scale freshwater aquaculture, income generation and food security in rural Madagascar. *Sustainability*, 15(21): 15439.
- Al Zamzami, I. M.; Sari, A. H. W. and Perwira, I. Y. (2019). Fluktuasi bahan organik dan residu terlarut dalam budidaya ikan koi di Tulungagung Jawa Timur. *Current Trends in Aquatic Science*, 2(1): 79-84.
- Ardianita, N.; Andriani, R. and Maya, N. P. (2023). Efektivitas ekoenzim terhadap kelulushidupan ikan lele (*Clarias* sp.) pada media yang tercemar limbah batik. *Biology Natural Resources Journal*, 2(1): 24-27.
- Bambang, B.; Umar, N. A. and Indrawati, E. (2024). Pengaruh ekoenzim dosis berbeda terhadap perbaikan kualitas air budidaya dalam pembesaran ikan lele mutiara *Clarias* sp. *Journal of Aquaculture and Environment*, 6(2): 84-89.

- Damayanti, H. O.** (2014). Tinjauan kualitas dan dampak ekonomi konsentrasi total dissolved solid (TDS) air di area pertambakan Desa Bulumanis Kidul. Jurnal Litbang: Media Informasi Penelitian, Pengembangan dan IPTEK, 10(2): 103-113.
- Ducatelle, R.; Goossens, E.; De Meyer, F.; Eeckhaut, V.; Antonissen, G.; Haesebrouck, F. and Van Immerseel, F.** (2018). Biomarkers for monitoring intestinal health in poultry: Present status and future perspectives. Veterinary Research, 49: 43.
- Febrianty, I.** (2021). Public perception of catfish aquaculture ponds in Banjar District, South Kalimantan, Indonesia. Aquaculture, Aquarium, Conservation & Legislation, 14(1): 227-232.
- Galintin, O.; Rasit, N. and Hamzah, S.** (2021). Production and characterization of eco enzyme produced from fruit and vegetable wastes and its influence on the aquaculture sludge. Biointerface Research in Applied Chemistry, 11(3): 10205-10214.
- Herliwati, H.; Rahman, M.; Ahmadi, A. and Tony, F.** (2024). Variations in the addition of eco enzyme in feed on the survival of siamese catfish (*Pangasius hypophthalmus*) seeds raised in aquariums. Fish Scientiae, 14(2): 145-154.
- Hidayati, S. N.; Laili, S. and Santoso, H.** (2021). Pengaruh kualitas air kolam terpal terhadap pertumbuhan ikan lele dumbo (*Clarias gariepinus*). Jurnal Ilmiah Biosaintropis, 6(2): 19-25.
- Islami, A. N.; Hasan, Z. and Anna, Z.** (2017). Pengaruh perbedaan siphonisasi dan aerasi terhadap kualitas air, pertumbuhan, dan kelangsungan hidup pada budidaya ikan nila (*Oreochromis niloticus*) stadia benih. Jurnal Perikanan Kelautan, 8(1): 73-82.
- Koniyo, Y.** (2020). Analisis kualitas air pada lokasi budidaya ikan air tawar di Kecamatan Suwawa Tengah. Jurnal Technopreneur, 8(1): 52-58.
- Kulla, O. L. S.; Yuliana, E. and Supriyono, E.** (2020). Analisis kualitas air dan kualitas lingkungan untuk budidaya ikan di Danau Laimadat, Nusa Tenggara Timur. Pelagicus, 1(3): 135-144.
- Kurniawan, A.; Mustikasari, D.; Kurniawan, A.; Muntoro, M.; Setiadi, J. and Kholishah, N. A.** (2024). The preliminary study about physico-chemical property of bio-enzyme produced from orange fruits waste treated with different concentrations of probiotic. Jurnal Ilmu Lingkungan, 22(4), 861-867.

- Kurniawan, A.; Mustikasari, D.; Muntoro.; Setiadi, J.; Wulandari, U. A.; Pratama, V.; Fadia, S.; Maheswari, D. A.; Fitri, C. and Iqbal, D. H. (2025). Effectiveness of decaying fruits as a carbon source on fish productivity and water quality improvement in aquaculture: A narrative review. *Egyptian Journal of Aquatic Biology & Fisheries*, 29(4): 2459-2481.
- Kusumawati, A. A.; Suprpto, D. and Haeruddin, H. (2018). Pengaruh ekoenzim terhadap kualitas air dalam pembesaran ikan lele (*Clarias gariepinus*), 7(4): 307-314.
- Li, H.; Cui, Z.; Cui, H.; Bai, Y.; Yin, Z. and Qu, K. (2023). Hazardous substances and removal in review. *Aquaculture*, 569: 739399.
- Mado, A. M.; Umar, N. A. and Aqmal, A. (2023). Pengaruh eco-enzym terhadap kualitas air pada sistem budidaya akuaponik ikan nila *Oreochromis niloticus* dan tanaman sawi *Brassica juncea*. *Journal of Aquaculture and Environment*, 5(2): 66-72.
- Melenchón, F.; de Mercado, E.; Pula, H. J.; Cardenete, G.; Barroso, F. G.; Fabrikov, D.; Lourenço, H. M.; Pessoa, M. F.; Lagos, L.; Weththasinghe, P.; Cortés, M. and Tomás-Almenar, C. (2022). Fishmeal dietary replacement up to 50%: A comparative study of two insect meals for rainbow trout (*Oncorhynchus mykiss*). *Animals*, 12(2): 179.
- Munguti, J. M.; Sendi, J. G.; Obiero, K. O.; Ogello, E. O.; Kyule, D. N.; Liti, D. M. and Musalia, L. M. (2021). Aqua-feed wastes: Impact on natural systems and practical mitigations-A review. *Journal of Agricultural Science*, 13(1): 111-121.
- Mustafa, S. A.; Al-Rudainy, A. J. and Salman, N. M. (2024). Effect of environmental pollutants on fish health: An overview. *Egyptian Journal of Aquatic Research*, 50(2): 225-233.
- Mustikasari, D.; Suryaningsih, S. and Nuryanto, A. (2020). Morphological variation of blue panchax (*Aplocheilichthys panchax*) lives in different habitat assessed using truss morphometric. *Biosaintifika: Journal of Biology & Biology Education*, 12(3): 399-407.
- Mzula, A.; Wambura, P. N.; Mdegela, R. H. and Shirima, G. M. (2021). Present status of aquaculture and the challenge of bacterial diseases in freshwater farmed fish in Tanzania; A call for sustainable strategies. *Aquaculture and Fisheries*, 6(3): 247-253.
- Nalle, M. M. D.; Kamlasi, Y. and Edo, S. I. (2021). Budidaya ikan lele dan ikan nila oleh Kelompok Tani Tunfeu Desa Ponain Kecamatan Amarasi Kabupaten Kupang di masa Covid-19. *Jurnal Pengabdian Perikanan Indonesia*, 1(2): 151-156.

- Natsir, N. I.; Jayadi, J. and Hadijah, S. (2024).** Aplikasi eco-enzyme terhadap pertumbuhan dan tingkat kelangsungan hidup ikan nila (*Oreochromis niloticus*). Jurnal Akuakultur Nusantara, 1(2): 96-106.
- Neupane, K. and Khadka, R. (2019).** Production of garbage enzyme from different fruit and vegetable wastes and evaluation of its enzymatic and antimicrobial efficacy. Tribhuvan University Journal of Microbiology, 6: 113-118.
- Putri, M. C.; Widiarini, R. and Ramadanintyas, K. N. (2024).** Effectiveness of eco enzymes in reducing chemical parameters of tofu factory liquid waste in Klumutan Village, Saradan District. Journal of Social Research, 3(9): 1-10.
- Rahman, M. F. I.; Wibisana, H. and Zainab, S. (2020).** Analisa dan pemetaan total padatan terlarut di pesisir Pantai Pasuruan dengan citra satelit Terra Modis. Journal of Civil Engineering, 16(2): 144-156.
- Rasit, N.; Fern, L. H. and Ghani, W. A. W. A. K. (2019).** Production and characterization of eco enzyme produced from tomato and orange wastes and its influence on the aquaculture sludge. International Journal of Civil Engineering and Technology, 10(3): 967-980.
- Serdiati, N.; Islamy, R. A.; Mamat, N. B.; Hasan, V. and Valen, F. S. (2024).** Nutritional value of alligator weed (*Alternanthera philoxeroides*) and its application for herbivorous aquaculture feed. International Journal of Agriculture and Biosciences, 13(3): 318-324.
- Serra, V.; Pastorelli, G.; Tedesco, D. E. A.; Turin, L. and Guerrini, A. (2024).** Alternative protein sources in aquafeed: Current scenario and future perspectives. Veterinary and Animal Science, 25: 100381.
- Siegers, W. H.; Prayitno, Y. and Sari, A. (2019).** Pengaruh kualitas air terhadap pertumbuhan ikan nila nirwana (*Oreochromis sp.*) pada tambak payau. The Journal of Fisheries Development, 3(2): 95-104.
- Sikku, E. S.; Umar, N. A. and Indrawati, E. (2023).** Penggunaan eco-enzim dengan dosis berbeda pada teknologi akuaponik sederhana untuk optimalisasi pertumbuhan ikan nila *Oreochromis niloticus*. Journal of Aquaculture and Environment, 5(2): 60-65.
- Singh, H.; Prasad, A. and Nal, K. V. (2019).** Study and analysis of bio enzymes based natural products. International Journal of Engineering Applied Sciences and Technology, 4(8): 205-207.

- Tumwesigye, Z.; Tumwesigye, W.; Opiyo, F.; Kemigabo, C. and Mujuni, B.** (2022). The effect of water quality on aquaculture productivity in Ibanda District, Uganda. *Aquaculture Journal*, 2(1): 23-36.
- Varshini, B. and Gayathri, V.** (2023). Role of eco-enzymes in sustainable development. *Nature Environment and Pollution Technology*, 22(3): 1299-1310.
- Yatma, M. A.; Hamzani, S. and Zubaidah, T.** (2024). The effect of eco enzyme on improving river water quality based on turbidity, temperature, pH, and TDS parameters. *Global Health & Environmental Perspectives*, 1(3): 164-168.

Egyptian Journal of Aquatic Biology & Fisheries

Zoology Department, Faculty of Science,

Ain Shams University,

Cairo, Egypt.

ISSN 1110 - 6131

<http://ejabf.journals.ekb.eg/>



Oct. 28, 2025

Acceptance

Dear **Muntoro, Andri Kurniawan, Ardiansyah Kurniawan, Ufi Ayu Wulandari, Didin Erlan, Venisa Pratama³, Syafiq Muhammad Syakir, Danella Aisah Maheswari, Safitri Fadia, Cahya Fitri**

I am pleased to inform you that your paper entitled “**Growth Performance of Tilapia and Catfish Fry Cultivated with Bio-Enzyme Supplementation and Its Agribusiness Perspective**” (Manuscript ID: **EJABF- 2510-6839**) has been accepted for publication in the **Egyptian Journal of Aquatic Biology and Fisheries (EJABF)**.

Congratulations on your successful submission, and we look forward to featuring your valuable research in our journal.

Regards,

Prof. Magdy T. Khalil

Co-Editor-In Chief of EJABF Journal,

Faculty of Science,

Ain Shams University,

Cairo, Egypt.





Growth Performance of Tilapia and Catfish Fry Cultivated with Bio-Enzyme Supplementation and Its Agribusiness Perspective

Muntoro¹, Andri Kurniawan^{2*}, Ardiansyah Kurniawan³, Ufi Ayu Wulandari³, Didin Erlan³, Venisa Pratama³, Syafiq Muhammad Syakir³, Danella Aisah Maheswari², Safitri Fadia², Cahya Fitri²

¹Department of Agribusiness, Faculty of Agriculture, Fisheries and Marine, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

²Department of Biology, Faculty of Science and Engineering, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

³Department of Aquaculture, Faculty of Agriculture, Fisheries and Marine, Bangka Belitung University, Jl Kampus Terpadu UBB, Balunijuk 33127, Bangka Belitung, Indonesia

*Corresponding Author: andri_pangkal@yahoo.co.id

ARTICLE INFO

Article History:

Received: July 20, 2025

Accepted: Sep. 20, 2025

Online: Oct. 29, 2025

Keywords:

Aquaculture,
Bio-Enzyme,
Effectiveness,
Growth rate,
Fish fry

ABSTRACT

Tilapia and catfish are freshwater fish commodities that are widely cultivated by the community because of their relatively easy maintenance, low cost, and short time. The quality of the cultivation water environment is an important factor and influences the growth performance and life of these commodities. This study aimed to analyze the growth performance of tilapia and catfish fry cultivated with the addition or supplementation of bio-enzymes with different concentrations in the maintenance water. This study used an experimental method with three treatments: control (without bio-enzymes), 500 ml/900 L or 0.6 ml/L bio-enzymes, and 1000 ml/900 L or 1.1 ml/L bio-enzymes. The results illustrate that bio-enzyme supplementation into tilapia fry ponds compared to the control treatment without bio-enzymes showed a higher contribution to weight growth than length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation into catfish fry ponds compared to the control without bio-enzymes showed a higher contribution to length growth than weight growth. Supplementation with 500 ml of bio-enzyme resulted in the highest absolute weight growth of tilapia fry, which was 16.95 g with a daily weight growth rate of 0.42 g/day and a specific weight growth rate of 3.35% and 100% survival rate. Supplementation with 500 ml of bio-enzyme resulted in the highest absolute length growth of catfish fry, which was 9.20 cm with a daily length growth rate of 0.23 cm/day and a specific length growth rate of 2.23% and 100% survival rate. In agribusiness perspective, only tilapia in consumption size (at 135 days old) treated with bio-enzymes (500 ml and 1000 ml) were recommended for sale, particularly in consumption size. Meanwhile, for catfish, only in fingerlings phase were recommended for selling with treatment bio-enzyme 500 ml.

INTRODUCTION

Fish farming, known as aquaculture, is one of the leading fish food sources with the fastest growth in productivity worldwide (**Sumaila *et al.*, 2022; Serra *et al.*, 2024**). Aquaculture stands as a vital provider of animal protein essential for human nutrition, with fish exhibiting rapid growth and offering optimal calorie-protein ratios for human consumption (**Melenchón, *et al.*, 2022**). Freshwater fish farming has been widely developed by communities, both on a household and large scale (**Mzula *et al.*, 2021; Angermayr *et al.*, 2023**). Among the freshwater fish commonly cultivated by the community are tilapia and catfish (**Febrianty, 2021; Nalle *et al.*, 2021**).

The development of fish farming productivity is greatly influenced by water quality, which significantly supports fish growth performance (**Ajayi & Okoh, 2014; Tumwesigye *et al.*, 2022**). Water quality, manifested in the form of physical, chemical, and biological parameters, affects the entire life cycle, including survival, growth, and reproduction (**Siegers *et al.*, 2019; Koniyo, 2020; Kulla *et al.*, 2020**).

Aquaculture water quality is generally contaminated by organic matter from leftover feed and metabolic waste in the form of fish feces. One effort to maintain and improve water quality in fish farming is the addition of enzymes to the culture environment.

Bio-enzymes or Bioz are fermented products made from fruits or vegetables that are widely used for various purposes due to their probiotic microorganism content and bioactive compounds (**Ducatelle *et al.*, 2018**). It is produced in 3 months fermentation process with ratio 1 : 3: 10 for sugar : waste : water (**Rasit *et al.*, 2019; Galintin *et al.*, 2021; Kurniawan *et al.*, 2024**). Bio-enzymes are also known as ecoenzymes, garbage enzymes, terrazymes, fruit enzymes, flower enzymes, and so on (**Neupane & Khadka, 2019; Singh *et al.*, 2019**). It is a fermentation product

One use of bio-enzymes is to accelerate the waste degradation process and improve environmental quality (**Ardianita *et al.*, 2023**). However, several studies have shown that the use of bio-enzymes (ecoenzymes) in fish farming environments does not show a different effect on reducing ammonia, nitrate, and nitrite and does not result in good fish growth (**Kusumawati *et al.*, 2018; Bambang *et al.*, 2024**). Based on the above description, this study was conducted to confirm previous studies by analyzing the growth performance of tilapia and catfish fry cultivated with the addition or supplementation of bio-enzymes with different concentrations in the maintenance water.

MATERIALS AND METHODS

This research was conducted at the Hatchery, Bangka Belitung University, from June to August 2025. The samples used in this study were 100 tilapia fry measuring 6-8 cm and catfish fry measuring 5-7 cm in each pond. Both types of samples were cultivated in concrete ponds measuring 2 m long, 1.5 m wide, and 0.6 m high, filled with water to a depth of 30 cm.

The research treatment consisted of two, namely cultivation media without added bio-enzymes and cultivation media added bio-enzymes. Based on pre research, effectiveness of bio-enzyme supplementation into the cultivation media was carried out before fish maintenance with doses of 500 ml and 1000 ml in each pond containing 900 L of water or treatments of 0.6 ml/L and 1.1 ml/L. The test sample fish fry were maintained in the research pond for 40 days by being fed pellets *ad satiation*, namely feeding until the fish were full and stopped eating.

Fish sample measurements were conducted on 10% of the population in each treatment. The research parameters observed in fish were absolute weight growth (g), absolute length growth (cm), daily weight growth rate (g/day), daily length growth rate (cm/day), specific weight growth rate (%), specific length growth rate (%), and survival (%). The water quality parameters observed were pH, temperature (°C), dissolved oxygen (DO) (ppm), total dissolved solids (TDS) (ppm), and electrical conductivity (EC) (mS/cm). The research data were analyzed descriptively and presented in tables and graphs.

RESULTS AND DISCUSSION

The results of the study showed that bio-enzyme supplementation into the pond compared to the control treatment without bio-enzyme can have an effect or no effect on the growth performance of fish fry. Bio-enzyme supplementation into tilapia fry ponds contributed more to higher weight growth than length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation into catfish fry ponds resulted in higher length growth than weight growth. Supplementation of 500 ml of bio-enzyme resulted in the highest absolute weight growth of tilapia fry, namely 16.95 g with a daily weight growth rate of 0.42 g/day and a specific weight growth rate of 3.35% and 100% survival. Supplementation of 500 ml of bio-enzyme resulted in the highest absolute length growth of catfish fry, namely 9.20 cm with a daily length growth rate of 0.23 cm/day and a specific length growth rate of 2.23% and 100% survival.

Growth of tilapia

Supplementation of bio-enzymes into the cultivation medium, namely the maintenance water in the tilapia fry pond resulted in higher weight growth of the fry at the end of the observation compared to the control treatment without bio-enzymes. However,

the length growth of the tilapia fry reared in the pond with bio-enzyme supplementation was lower than the control treatment without bio-enzymes (Fig. 1). The absolute growth, daily growth rate, and specific growth rate of the tilapia fry are shown in Table (1). The research data also did not show any symptoms of disease in the fry reared, but there were 3 fish that died during maintenance in the 1000 ml bio-enzyme treatment, while in the other treatments no dead fry were found during maintenance (Table 1).




Fig. 1. Growth in weight and length of tilapia fry

Table 1. Calculation of growth and survival of tilapia fry

Parameter	Tilapia Fry		
	Control	Bio-enzyme	
		500 ml	1000 ml
Absolute weight gain (g)	11.94	16.95	16.47
Absolute length growth (cm)	17.75	12.98	10.20
Daily weight gain rate (g/day)	0.30	0.42	0.41
Daily length growth rate (cm/day)	0.44	0.32	0.26
Specific weight growth rate (%)	1.88	3.35	3.13
Specific length growth rate (%)	2.98	2.74	2.23
Survival rate (%)	100	100	97

Growth of catfish

Supplementation of bio-enzymes into the culture medium, namely the maintenance water in the catfish fry pond, resulted in lower weight growth of the fry at the end of the observation compared to the control treatment without bio-enzymes. However, the length growth of the catfish fry reared in the pond with bio-enzyme supplementation was higher than the control treatment without bio-enzymes (Fig. 2). The absolute growth, daily growth rate, and specific growth rate of the catfish fry were shown in Table (1). The research data also showed no symptoms of disease and no dead fry were found during the maintenance (Table 2).




Fig. 2. Growth in weight and length of catfish fry

Table 2. Calculation of growth and survival of catfish fry

Parameter	Catfish Fry		
	Control	Bio-enzyme	
		500 ml	1000 ml
Absolute weight gain (g)	15.48	15.40	13.28
Absolute length growth (cm)	7.55	9.20	7.85
Daily weight gain rate (g/day)	0.39	0.39	0.33
Daily length growth rate (cm/day)	0.19	0.23	0.20
Specific weight growth rate (%)	5.65	5.69	5.23
Specific length growth rate (%)	1.94	2.23	1.99
Survival rate (%)	100	100	100

Water quality as a medium for cultivating tilapia and catfish fry

The results of water quality measurements used as a culture medium for the Nile fish and catfish fry—pH, temperature, DO, TDS, and EC—showed that all parameters were dynamic. This indicates that water quality fluctuates during cultivation and does not reflect a relationship between bio-enzyme supplementation in the culture medium and water quality.

Water quality in the tilapia fry rearing ponds showed a pH value that tended to be higher than that in the catfish fry ponds. The water temperature in the tilapia fry rearing ponds was relatively lower than that in the catfish fry ponds. Dissolved oxygen in the water in the tilapia fry ponds was relatively better than that in the catfish fry ponds. Total dissolved solids and electrical conductivity in the tilapia and catfish fry control ponds were relatively higher at certain times and relatively lower at other times compared to the TDS in the bio-enzyme treatment ponds.

The water quality in the tilapia fry pond had the highest pH value in the 500 ml bio-enzyme treatment, the highest temperature in the 1000 ml bio-enzyme treatment, the highest DO in the 500 ml bio-enzyme treatment, and the highest TDS and EC in the control treatment. The water quality in the catfish fry pond had the highest pH value in the 500 ml bio-enzyme treatment, the highest temperature in the control treatment, and the highest DO, TDS, and EC in the control treatment. The water quality conditions and comparison between the water quality in the tilapia and catfish fry ponds are shown in Figs. (3, 4).




Fig. 3. Water quality as a medium for tilapia cultivation




Fig. 4. Water quality as a medium for catfish cultivation

Sikku *et al.* (2023) conducted a study on the application of 5-15 ml/L of ecoenzyme (bio-enzyme) in tilapia fry (3 cm in length) cultivation pond maintained for 45 days. The absolute weight produced ranged from 0.98-1.76 g with a specific weight growth rate ranging from 2.33-4.19%. The water pH value during maintenance ranged from 7.6-9.1. When compared with the specific weight growth rate in this study which ranged from 3.13-3.35% (Table 1), then supplementation with a high concentration of bio-enzyme (15 ml/L) resulted in a better specific weight growth rate than the low concentration (0.6-1.1 ml/L) in this study. However, at a concentration of 5-10 ml/L in the study by **Sikku *et al.* (2023)** showed a lower value (2.33-2.98%) compared to the treatment in this study. Thus, there is no linear pattern that shows that the higher the concentration of bio-enzyme given, the higher the growth of tilapia fry.

Bio-enzyme research has also been conducted on catfish fry cultivation. The results of **Kusumawati *et al.* (2018)** research show that the addition of bio-enzymes to catfish fry ponds maintained for 4 weeks (approximately 28 days) has a length of 5.9-8.0 cm with a weight of 2.01-4.32 g from a length of 5.1-5.8 cm and a weight of 1.12-1.22 g on day 0. Meanwhile, the results of measurements on the 26th day of this study have shown that the weight and length data were 8.76 g and 9.22 cm, respectively, in the control treatment; 8.14 g and 10.12 cm in the 500 ml bio-enzyme treatment (0.6 ml/L); and 7.90 g and 9.82 cm in the 1000 ml bio-enzyme treatment (1.1 ml/L) (Fig. 4). This difference may be caused by the contribution of bio-enzymes in the research of **Kusumawati *et al.* (2018)** found lower pH and DO values, ranging from 5.85 to 6.89 and DO from 2.95 to 3.71 ppm in the fourth week of culture. This study, however, found pH values ranging from 8.33 to 9.08 and DO from 4.8 to 6.8 ppm. Based on SNI 6484.4:2014, water quality for catfish fry culture has a pH of 6.5 to 8 and DO > 3 ppm.

The description shows that specifically, the contribution of bio-enzyme supplementation to the growth performance of tilapia fry was better than that of catfish fry (Tables 1, 2). This was in line with the statement by **Herliwati *et al.* (2024)** that the addition of bio-enzymes can improve water quality, which can impact the survival of catfish fry. However, the addition of bio-enzymes to the cultivation water medium only affected the growth of tilapia and not catfish. Similar conditions were also found in the results of research by **Mado *et al.* (2023)** and **Natsir *et al.* (2024)**, who found increased growth of tilapia with the application of bio-enzymes to the cultivation water medium. However, according to **Kusumawati *et al.* (2018)** and **Bambang *et al.* (2024)**, catfish growth did not change with the same application. This is possible because catfish have a tolerance to less than ideal waters (**Hidayati *et al.*, 2021**), so that when water quality improves, its metabolism does not affect. Meanwhile, tilapia experience increased metabolism when quality improves, resulting in better growth. Tilapia fry experience increased growth and survival when oxygen and ammonia levels are more appropriate (**Islami *et al.*, 2017**). When compared to the pH, DO, and TDS conditions (Figs. 3 and 4), it can be seen that supplementing bio-enzymes into the culture medium results in better water quality for tilapia fry cultivation than for catfish fry. Furthermore, Figs. (3 and 4) also show that the total dissolved solids in catfish fry ponds were higher than in tilapia fry ponds. High dissolved solids can hinder the growth performance of fish fry.

Rahman *et al.* (2020) explained that total dissolved solids are organic and inorganic substances dissolved in water that can be a factor causing water turbidity. According to **Damayanti (2014)**, increased TDS concentrations in water can originate from feed waste and accumulated metabolic waste such as fish feces. According to **Al Zamzami *et al.* (2019)**, the length of maintenance time can contribute to an increase in dissolved organic matter, one of which comes from feed waste and fish feces. Furthermore, various studies have revealed that the presence of nitrogen waste can produce ammonia, nitrate, and nitrite

compounds that can cause oxygen depletion, disease, and even death in fish (**Munguti et al., 2021; Li et al., 2023; Mustafa et al., 2024; Kurniawan et al., 2025**).

Relatively low bio-enzyme concentrations were considered ineffective in reducing waste contamination, particularly in catfish fry ponds. This is visually confirmed by the cloudier and more odorous water in catfish fry ponds compared to tilapia fry ponds. This is thought to inhibit the growth of catfish fry in rearing ponds. **Kusumawati et al. (2018)** explained that administering bio-enzymes at low concentrations does not significantly reduce ammonia and nitrite concentrations. This contributed to the slow growth of catfish fry.

Higher bio-enzyme concentrations, duration, and frequency of bio-enzyme administration need to be tested for their effectiveness in improving the quality of aquaculture water and its effect on optimal growth for fish fry. This is in line with **Putri et al. (2024)**, who stated that the longer the time and the higher the eco-enzyme dosage, the higher the effectiveness in reducing waste contamination as indicated by the biological oxygen demand (COD) and total suspended solids (TSS) in the environment. **Yatma et al. (2024)** also revealed that increasing the eco-enzyme dosage used was able to significantly reduce the dissolved solids (TDS) content and improve water quality. Furthermore, water quality and nutrient contribute to fish morphological and its growth (**Mustikasari et al., 2020; Serdiati et al., 2024**).

Discussion about Agribusiness Perspective of Bio-Enzyme Supplementation for Aquaculture

From an agribusiness perspective, bio-enzyme supplementation in aquaculture presents a high-potential growth sector that aligns with key trends: sustainable food production, environmental regulation, and consumer preference for clean-label seafood. Strategic investment in research and development, production, and farmer outreach can unlock long-term profitability and ecological benefits.

Perspective in agribusiness opportunities, including market potential, value addition, product development and diversification, challenges from an agribusiness viewpoint, and investment and agribusiness models. In Indonesia (and many other countries) tilapia and catfish are major freshwater/pond-aquaculture species. Grower demand is strong for both local consumption and export. A value proposition; higher growth/lower feed cost/improved survival is very attractive to fish farmers, in particular tilapia and catfish. Use the enzyme supplementation in aquaculture can improve output and profit. In other hand, enzyme supplementation can reduce feed costs (better feed conversion ratio or FCR), improves animal health and survival rates — leading to higher yields and profits, reduces environmental impact, enhancing the farm's market reputation and compliance with green regulations (**Varshini & Gayathri, 2023**).

Using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming. The benefits using bio-enzymes in aquaculture from an agribusiness perspective:

Improved Productivity and Profitability

Enhanced Feed Conversion Ratio (FCR) where bioenzymes help fish and shrimp digest feed more efficiently, reducing feed costs — a major operational expense. Faster growth rates where enzymes promote better nutrient absorption, accelerating growth and reducing time to market. Lower mortality rates where enzymes improve gut health and immunity, reducing disease outbreaks.

Profitability of this research result can be calculated with assumption referring to Tables (1 and 2). Tilapia and catfish fish fry were cultivated 45 days for fingerlings size and 135 days for consumption size with survival rate 100%. The potential economic value of selling of fish fry and consumption sizes is shown in Table (3).

The economic value of selling tilapia at the fingerling phase or consumption size, as presented in Table (3), indicates that only tilapia treated with bio-enzymes were recommended for sale. The potential of selling price from bio-enzyme-supplemented fish was higher than normal price at local market. Consumption size with treatment bio-enzyme 500 ml had potential of selling price 52,000 IDR and bio-enzyme 1000 ml had potential of selling price 47,565 IDR compared price at local market 35,000 IDR. Meanwhile, selling of fingerlings phase of bio-enzyme-supplemented tilapia was not recommended due to the size of fish fry treated with bio-enzymes was not better than the control.

For catfish, the control and bio-enzyme treatment of 500 ml and 1000 ml of enzyme did not show significant differences. Even though, fingerlings phase (at 45 days old) treated with bio-enzyme 500 ml had potential of selling price 495 IDR, higher than price at local market (300 IDR). Thus, selling of fingerlings phase of bio-enzyme-supplemented catfish (500 ml bio-enzyme) was recommended.

Weight of catfish fry in the control treatment was bigger than the bio-enzyme treatment so that the addition of bio-enzymes had no effect on fry growth. As well as the weight of the fish, the addition of bioenzymes did not result in significant weight gain compared to the control treatment. Therefore, selling of fingerlings phase and also consumption size of bio-enzyme-treated fish were not recommended

Table 3. Economic value of tilapia and catfish in fingerlings and consumption size

Fish (Phase or Size)	Comparing Between Control and Bio-enzym Supplementation (500 ml and 1000 ml)		Price (IDR) in local market in Bangka, Indonesia	Potential of Selling Price (IDR)	Selling Recommendation of Bio-enzyme- supplemented Fish
	Weight (g)	Length (cm)			
Tilapia					
Fingerlings phase (at 45 days old)	5.01 x 100 fishes	Control > Treatment	200/fish	No	Not recommended for selling in fingerlings phase
	4.53 x 100 fishes				
Consumption size (at 135 days old)	15.03 x 100 fishes	-	35,000/kg (35/g)	52,000	Recommended for selling in consumption size (treatment bio-enzyme 500 ml)
	13.59 x 100 fishes			47,565	Recommended for selling in consumption size (treatment bio-enzyme 1000 ml)
Catfish					
Fingerlings phase (at 45 days old)	Control > Treatment	1.65	300/fish	495	Recommended for selling in consumption size (treatment bio-enzyme 500 ml)
		0.3		90	Not recommended for selling in fingerlings phase
Consumption size (at 135 days old)	-	4.95 x 100 fishes	23,000/kg (23/g)	11,385	Not recommended for selling in consumption size
		0.9 x 100 fishes		2,070	

Reduction in Input Costs

Less dependency on antibiotics and chemicals where enzymes can reduce disease pressure naturally. Better water quality management where bioenzymes help degrade organic waste and ammonia, reducing the need for chemical water treatments.

Sustainable Branding & Market Access

Eco-friendly production where enzyme use aligns with growing consumer demand for sustainable and antibiotic-free aquaculture. Certification ready where farms using bioenzymes can more easily meet organic or eco-certification standards.

Waste Reduction and Circular Economy

Utilization of agricultural byproducts where some bioenzymes are produced using agri-waste (e.g., fruit peels, crop residues), supporting circular bioeconomy models. Efficient waste breakdown where help manage sludge and organic waste, reducing environmental impact.

New Revenue Streams

Local production of bioenzymes where farmers or agri-entrepreneurs can engage in low-cost production of bioenzymes using fermentation, creating micro-enterprises. Diversification for agro-industries where companies in biotech, agriculture, or food waste management can expand into aquaculture support services.

Regulatory and Environmental Compliance

Supports green policies where use of bioenzymes aligns with government and international sustainability goals in aquaculture. Reduces pollution where better water quality means less eutrophication and contamination of surrounding ecosystems.

CONCLUSION

The addition or supplementation of bio-enzymes to tilapia and catfish fry ponds produced different contributions. Bio-enzyme supplementation to tilapia fry ponds compared to the control treatment without bio-enzymes showed a higher contribution to weight growth than to length growth. The opposite occurred for catfish fry, namely bio-enzyme supplementation to catfish fry ponds compared to the control without bio-enzymes showed a higher contribution to length growth than to weight growth. The results of water quality measurements as a medium for cultivating tilapia and catfish fry, namely pH, temperature, DO, TDS, and EC showed that all parameters were dynamic. This means that water quality fluctuates during cultivation and does not reflect the relationship between bio-enzyme supplementation to the cultivation medium and water quality. In agribusiness

perspective, using bio-enzymes in aquaculture offers several agribusiness advantages, making them an increasingly attractive input in sustainable fish farming. However, based on the economic value, only tilapia in consumption size (at 135 days old) treated with bio-enzymes (500 ml and 1000 ml) were recommended for sale, particularly in consumption size. Meanwhile, for catfish, only in fingerlings phase were recommended for selling with treatment bio-enzyme 500 ml.

ACKNOWLEDGMENT

The authors would like to thank Bangka Belitung University for supporting and funding this research and publication through Grant of Penelitian Unggulan 2025 with grant number 1116/UN50/M/PP/2025.

CONFLICTS OF INTEREST

All the authors declare no conflict of interest.

FUNDING

This research and publication were supported and funded through the Grant of Penelitian Unggulan 2025 grant number 1116/UN50/M/PP/2025.

REFERENCES

- Ajayi, A. O. and Okoh, A. I. (2014).** Bacteriological study of pond water for aquaculture purposes. *Journal of Food Agriculture and Environment*, 12(2): 1260-1265.
- Angermayr, G.; Palacio, A. and Chaminade, C. (2023).** Small-scale freshwater aquaculture, income generation and food security in rural Madagascar. *Sustainability*, 15(21): 15439.
- Al Zamzami, I. M.; Sari, A. H. W. and Perwira, I. Y. (2019).** Fluktuasi bahan organik dan residu terlarut dalam budidaya ikan koi di Tulungagung Jawa Timur. *Current Trends in Aquatic Science*, 2(1): 79-84.
- Ardianita, N.; Andriani, R. and Maya, N. P. (2023).** Efektivitas ekoenzim terhadap kelulushidupan ikan lele (*Clarias* sp.) pada media yang tercemar limbah batik. *Biology Natural Resources Journal*, 2(1): 24-27.
- Bambang, B.; Umar, N. A. and Indrawati, E. (2024).** Pengaruh ekoenzim dosis berbeda terhadap perbaikan kualitas air budidaya dalam pembesaran ikan lele mutiara *Clarias* sp. *Journal of Aquaculture and Environment*, 6(2): 84-89.

- Damayanti, H. O.** (2014). Tinjauan kualitas dan dampak ekonomi konsentrasi total dissolved solid (TDS) air di area pertambakan Desa Bulumanis Kidul. *Jurnal Litbang: Media Informasi Penelitian, Pengembangan dan IPTEK*, 10(2): 103-113.
- Ducatelle, R.; Goossens, E.; De Meyer, F.; Eeckhaut, V.; Antonissen, G.; Haesebrouck, F. and Van Immerseel, F.** (2018). Biomarkers for monitoring intestinal health in poultry: Present status and future perspectives. *Veterinary Research*, 49: 43.
- Febrianty, I.** (2021). Public perception of catfish aquaculture ponds in Banjar District, South Kalimantan, Indonesia. *Aquaculture, Aquarium, Conservation & Legislation*, 14(1): 227-232.
- Galintin, O.; Rasit, N. and Hamzah, S.** (2021). Production and characterization of eco enzyme produced from fruit and vegetable wastes and its influence on the aquaculture sludge. *Biointerface Research in Applied Chemistry*, 11(3): 10205-10214.
- Herliwati, H.; Rahman, M.; Ahmadi, A. and Tony, F.** (2024). Variations in the addition of eco enzyme in feed on the survival of siamese catfish (*Pangasius hypophthalmus*) seeds raised in aquariums. *Fish Scientiae*, 14(2): 145-154.
- Hidayati, S. N.; Laili, S. and Santoso, H.** (2021). Pengaruh kualitas air kolam terpal terhadap pertumbuhan ikan lele dumbo (*Clarias gariepinus*). *Jurnal Ilmiah Biosaintropis*, 6(2): 19-25.
- Islami, A. N.; Hasan, Z. and Anna, Z.** (2017). Pengaruh perbedaan siphonisasi dan aerasi terhadap kualitas air, pertumbuhan, dan kelangsungan hidup pada budidaya ikan nila (*Oreochromis niloticus*) stadia benih. *Jurnal Perikanan Kelautan*, 8(1): 73-82.
- Koniyo, Y.** (2020). Analisis kualitas air pada lokasi budidaya ikan air tawar di Kecamatan Suwawa Tengah. *Jurnal Technopreneur*, 8(1): 52-58.
- Kulla, O. L. S.; Yuliana, E. and Supriyono, E.** (2020). Analisis kualitas air dan kualitas lingkungan untuk budidaya ikan di Danau Laimadat, Nusa Tenggara Timur. *Pelagicus*, 1(3): 135-144.
- Kurniawan, A.; Mustikasari, D.; Kurniawan, A.; Muntoro, M.; Setiadi, J. and Kholishah, N. A.** (2024). The preliminary study about physico-chemical property of bio-enzyme produced from orange fruits waste treated with different concentrations of probiotic. *Jurnal Ilmu Lingkungan*, 22(4), 861-867.

- Kurniawan, A.; Mustikasari, D.; Muntoro.; Setiadi, J.; Wulandari, U. A.; Pratama, V.; Fadia, S.; Maheswari, D. A.; Fitri, C. and Iqbal, D. H. (2025).** Effectiveness of decaying fruits as a carbon source on fish productivity and water quality improvement in aquaculture: A narrative review. *Egyptian Journal of Aquatic Biology & Fisheries*, 29(4): 2459-2481.
- Kusumawati, A. A.; Suprpto, D. and Haeruddin, H. (2018).** Pengaruh ekoenzim terhadap kualitas air dalam pembesaran ikan lele (*Clarias gariepinus*), 7(4): 307-314.
- Li, H.; Cui, Z.; Cui, H.; Bai, Y.; Yin, Z. and Qu, K. (2023).** Hazardous substances and removal in review. *Aquaculture*, 569: 739399.
- Mado, A. M.; Umar, N. A. and Aqmal, A. (2023).** Pengaruh eco-enzym terhadap kualitas air pada sistem budidaya akuaponik ikan nila *Oreochromis niloticus* dan tanaman sawi *Brassica juncea*. *Journal of Aquaculture and Environment*, 5(2): 66-72.
- Melenchón, F.; de Mercado, E.; Pula, H. J.; Cardenete, G.; Barroso, F. G.; Fabrikov, D.; Lourenço, H. M.; Pessoa, M. F.; Lagos, L.; Weththasinghe, P.; Cortés, M. and Tomás-Almenar, C. (2022).** Fishmeal dietary replacement up to 50%: A comparative study of two insect meals for rainbow trout (*Oncorhynchus mykiss*). *Animals*, 12(2): 179.
- Munguti, J. M.; Sendi, J. G.; Obiero, K. O.; Ogello, E. O.; Kyule, D. N.; Liti, D. M. and Musalia, L. M. (2021).** Aqua-feed wastes: Impact on natural systems and practical mitigations-A review. *Journal of Agricultural Science*, 13(1): 111-121.
- Mustafa, S. A.; Al-Rudainy, A. J. and Salman, N. M. (2024).** Effect of environmental pollutants on fish health: An overview. *Egyptian Journal of Aquatic Research*, 50(2): 225-233.
- Mustikasari, D.; Suryaningsih, S. and Nuryanto, A. (2020).** Morphological variation of blue panchax (*Aplocheilichthys panchax*) lives in different habitat assessed using truss morphometric. *Biosaintifika: Journal of Biology & Biology Education*, 12(3): 399-407.
- Mzula, A.; Wambura, P. N.; Mdegela, R. H. and Shirima, G. M. (2021).** Present status of aquaculture and the challenge of bacterial diseases in freshwater farmed fish in Tanzania; A call for sustainable strategies. *Aquaculture and Fisheries*, 6(3): 247-253.
- Nalle, M. M. D.; Kamlasi, Y. and Edo, S. I. (2021).** Budidaya ikan lele dan ikan nila oleh Kelompok Tani Tunfeu Desa Ponain Kecamatan Amarasi Kabupaten Kupang di masa Covid-19. *Jurnal Pengabdian Perikanan Indonesia*, 1(2): 151-156.

- Natsir, N. I.; Jayadi, J. and Hadijah, S. (2024).** Aplikasi eco-enzyme terhadap pertumbuhan dan tingkat kelangsungan hidup ikan nila (*Oreochromis niloticus*). Jurnal Akuakultur Nusantara, 1(2): 96-106.
- Neupane, K. and Khadka, R. (2019).** Production of garbage enzyme from different fruit and vegetable wastes and evaluation of its enzymatic and antimicrobial efficacy. Tribhuvan University Journal of Microbiology, 6: 113-118.
- Putri, M. C.; Widiarini, R. and Ramadanintyas, K. N. (2024).** Effectiveness of eco enzymes in reducing chemical parameters of tofu factory liquid waste in Klumutan Village, Saradan District. Journal of Social Research, 3(9): 1-10.
- Rahman, M. F. I.; Wibisana, H. and Zainab, S. (2020).** Analisa dan pemetaan total padatan terlarut di pesisir Pantai Pasuruan dengan citra satelit Terra Modis. Journal of Civil Engineering, 16(2): 144-156.
- Rasit, N.; Fern, L. H. and Ghani, W. A. W. A. K. (2019).** Production and characterization of eco enzyme produced from tomato and orange wastes and its influence on the aquaculture sludge. International Journal of Civil Engineering and Technology, 10(3): 967-980.
- Serdiati, N.; Islamy, R. A.; Mamat, N. B.; Hasan, V. and Valen, F. S. (2024).** Nutritional value of alligator weed (*Alternanthera philoxeroides*) and its application for herbivorous aquaculture feed. International Journal of Agriculture and Biosciences, 13(3): 318-324.
- Serra, V.; Pastorelli, G.; Tedesco, D. E. A.; Turin, L. and Guerrini, A. (2024).** Alternative protein sources in aquafeed: Current scenario and future perspectives. Veterinary and Animal Science, 25: 100381.
- Siegers, W. H.; Prayitno, Y. and Sari, A. (2019).** Pengaruh kualitas air terhadap pertumbuhan ikan nila nirwana (*Oreochromis sp.*) pada tambak payau. The Journal of Fisheries Development, 3(2): 95-104.
- Sikku, E. S.; Umar, N. A. and Indrawati, E. (2023).** Penggunaan eco-enzim dengan dosis berbeda pada teknologi akuaponik sederhana untuk optimalisasi pertumbuhan ikan nila *Oreochromis niloticus*. Journal of Aquaculture and Environment, 5(2): 60-65.
- Singh, H.; Prasad, A. and Nal, K. V. (2019).** Study and analysis of bio enzymes based natural products. International Journal of Engineering Applied Sciences and Technology, 4(8): 205-207.

- Tumwesigye, Z.; Tumwesigye, W.; Opiyo, F.; Kemigabo, C. and Mujuni, B. (2022).** The effect of water quality on aquaculture productivity in Ibanda District, Uganda. *Aquaculture Journal*, 2(1): 23-36.
- Varshini, B. and Gayathri, V. (2023).** Role of eco-enzymes in sustainable development. *Nature Environment and Pollution Technology*, 22(3): 1299-1310.
- Yatma, M. A.; Hamzani, S. and Zubaidah, T. (2024).** The effect of eco enzyme on improving river water quality based on turbidity, temperature, pH, and TDS parameters. *Global Health & Environmental Perspectives*, 1(3): 164-168.