

▶ Conference Proceedings Citation Index (Web of Science) <a>C</a>

▶ Materials Science & Engineering Database (ProQuest) ☑

▶ SciTech Premium Collection (ProQuest) ☑

▶ Technology Collection (ProQuest) ☑

▶ EBSCO (EBSCO Discovery Service) <a>™</a>

▶ DOAJ []

▶ Inspec ☑

▶ Google Scholar ☑

▶ Polymer Library ☑

▶ Wanfang Data ☑

### Crossref indexation

This journal is indexed in Crossref.

The Crossref system is used for linking citations across publishers. To ensure direct linking to and from its contents, EDP Sciences registers its scientific research journals to Crossref. Online archives have also been indexed.

Any Crossref members can create direct links from its online references to the EDP Sciences contents, using Crossref database. Reciprocally, the online references of EDP Sciences articles include direct links to the cited articles. Such links can be created on condition that the reference of the cited article matches a Crossref data. In particular these direct links only concern references whose publishers have previously indexed their own data to Crossref.

Read more about Crossref .



### MATEC Web of Conferences

ef55N: 2261-236X


Copyright / Published by: EDP Sciences





MATEC Web of Conferences

All issues Series Forthcoming About



All issues ▶ Volume 101 (2017) ▶ MATEC Web Conf., 101 (2017) 04010 ▶ Abstract

### Open Access

Issue MATEC Web Conf.

Volume 101, 2017

Sriwijaya International Conference on Engineering, Science and Technology (SICEST 2016)

Article Number 04010 Number of page(s) 6

Section Applied Technology for Sustainable Environment
DOI https://doi.org/10.1051/matecconf/201710104010

Published online 09 March 2017

MATEC Web of Conferences 101, 04010 (2017)

## The Impact of tin mining in Bangka Belitung and its reclamation studies

Eddy Nurtjahya<sup>1\*</sup>, Jennifer Franklin<sup>2</sup>, Umroh<sup>3</sup> and Fournita Agustina<sup>4</sup>

- <sup>1</sup> University of Bangka Belitung, Biology Department, 33172 Merawang, Indonesia
- <sup>2</sup> University of Tennessee, Forestry, Wildlife and Fisheries Department, 274 Ellington Plant Sciences Building, TN 37996, USA
- <sup>3</sup> University of Bangka Belitung, Fisheries Department, 33172 Merawang, Indonesia
- <sup>4</sup> University of Bangka Belitung, Agribusiness Department, 33172 Merawang, Indonesia

© The Authors, published by EDP Sciences, 2017



This is an Open Access article distributed under the terms of the Creative Commons Attribution License 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### **MATEC Web of Conferences**

elSSN: 2261-236X

Copyright / Published by: EDP Sciences

<sup>\*</sup> Corresponding author: eddy\_nurtjahya@yahoo.com

### **Editorial board**

### Rachid Bennacer

École Normale Supérieure, Cachan, France

website

### Chérifa Boukacem-Zeghmouri

Université Claude Bernard Lyon 1, Villeurbanne, France

website

### Vladimir Buzek

Slovak Academy of Sciences, Bratislava, Slovakia

website

### Heidi Gautschi

Haute Ecole Pédagogique de Lausanne, Switzerland

### Éric Lichtfouse

French National Institute for Agricultural Research (INRA), Aix-en-Provence, France website

### Ming-Jun Zhang

DGUT-CNAM Institute, Dongguan University of Technology, Guangdong Province, P.R. China website

### **Zhien Zhang**

The Ohio State University, Columbus, USA

### **MATEC Web of Conferences**

elSSN: 2261-236X

Copyright / Published by: EDP Sciences



## The Impact of tin mining in Bangka Belitung and its reclamation studies

Eddy Nurtjahya<sup>1,\*</sup>, Jennifer Franklin<sup>2</sup>, Umroh<sup>3</sup> and Fournita Agustina<sup>4</sup>

Abstract. Tin mining in Bangka Belitung has been exploited for hundred years. The province is the second largest tin producer in the world. Secondary data from studies which took place in all four regencies in Bangka Island were discussed to show the impact of mining activities and its reclamation studies. In order to add plant selection criteria for regevetate mined soil, the greenhouse and laboratory experiment was carried out with fourteen herbs and grass species in Tennessee. The mining activities increase the wealth of the community, but the other hand they change and decrease the environmental stability, and cause horizontal conflicts. Offshore mining reduced water quality, change sea bed caused the change of biodiversity. Onshore mining activity reduces biodiversity and causes floods and damages infrastructure. While the more economic species are demanded, planting local tree species is challenging. An evaluation with local tree species concluded that best adapted species based on anatomical and physiological measurements was not those that showed the best performance in the field. The greenhouse and laboratory findings indicate that some physiological characteristics i.e. plant height and cover, transpiration rate, and foliar pigments may be used to select plant adaptability to mined soil.

### 1 Introduction

Bangka Belitung Islands produced approximately 106,000 t of tin in 2013 or more than one third of global tin supply, and the majority is exported [1]. Tin mining is the most significant economic-driver in the province, taking place inland and offshore, including in protected forests and marine ecosystems [1].

Following the issuance of a 1999 Ministry of Trade and Industry decree that tin is not an export item to be monitored and regulated, the Bangka regent issued a decree in 2001 giving permission for the people to mine tin

There were 80 dredges and nearly 4000 floating tin mines off the shore of Bangka Island in 2013 [2], and are up to 50,000 artisanal small scale mines (ASM) and approximately 30 independent smelters [1]. Tin production from ASM contributes up to 80% of Indonesian tin exports [3].

Tin mining activities increase the wealth of the local people, but most of the activities neglect good mining practices, safety and land reclamation [3]. The most accidents at inland mine sites are due to landslides, and non-standard diving devices [4]. The mining activity decrease the environmental stability, causes pollution, and cause horizontal conflicts. Offshore mining reduced water quality, change sea bed caused the change of biodiversity.

An evaluation of ten native tree species concluded that four of ten species showed highest survival rates and cover development [7] but measurement of anatomical and physiological measurements of five year saplings of those species, best adapted species were not those that showed the best performance in the field (Table 1) [8].

The identification of traits that can be used to identify species for potential use in mine reclamation would greatly speed the search for appropriate native species. The early growth, pigment content, and transpiration rates of 14 herbaceous species are studied to determine whether these traits can be used to predict ground cover success on mine reclamation sites.

Anatomy and physiology parameters: stomatal density, leaf thickness, palisade thickness, sponge thickness, upper epidermal thickness, lower epidermal thickness, upper cuticle thickness, lower cuticle thickness, xylem diameter, xylem bundles, root diameter, root conductity ratio. Morphology parameters: survival rate and cover area (1 year old), and height, stem diameter, and cover area (5 year old)

<sup>&</sup>lt;sup>1</sup>University of Bangka Belitung, Biology Department, 33172 Merawang, Indonesia

<sup>&</sup>lt;sup>2</sup>University of Tennessee, Forestry, Wildlife and Fisheries Department, 274 Ellington Plant Sciences Building, TN 37996, USA

<sup>&</sup>lt;sup>3</sup>University of Bangka Belitung, Fisheries Department, 33172 Merawang, Indonesia

<sup>&</sup>lt;sup>4</sup>University of Bangka Belitung, Agribusiness Department, 33172 Merawang, Indonesia

While the more plant species planted in mined soils in the last decade, economic species such as rubber, oil palm, and some fruit species are demanded. Planting local tree species is challenging. There is an additional challenge in cases where reclaimed soils are contaminated with metals, are highly acidic, or coarsely textured. Natural succession takes a long time [5]. Soil amendments and land preparation are the major costs

<sup>\*</sup> Corresponding author: eddy nurtjahya@yahoo.com

<sup>©</sup> The Authors, published by EDP Sciences. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (http://creativecommons.org/licenses/by/4.0/).

**Table 1.** The most adaptive local tree species on sandy tin tailing at one and five year old based on anatomy and physiology, and morphology measurements.

	1 year old [7]	5 year old [8]
Anatomy and Physiology	not measured	Vitex pinnata Calophyllum inophyllum Syzygium grande
Morphology	Hibiscus tiliaceus Ficus superba Calophyllum inophyllum Syzygium grande Vitex pinnata	Syzygium grande Calophyllum inophyllum Vitex pinnata

### 2 Methods

Bangka Island, with a population of nearly one million, is located off the eastern coast of South Sumatra Island [9]. The island has a surface area of 11,900 km² and is mainly lowland below 50 m; its climatic differences within the island are small. Its climate belonging to the Af-type Köppen-Geiger climate classification [10], with an average temperature of 26.3°C, average humidity of 61.7% and average annual rainfall of approximately 2,400 mm [7].

The authors gathered secondary data from studies which took place in all four regencies in Bangka Island (Figure 1), and in the green house and laboratory in the University of Tennessee, Knoxville, USA to indicate that some physiological characteristics may be used to select plant adaptability to mined soil.

The experiment was carried out with fourteen herbs and grass species whose seeds are widely available commercially within the southeastern United States. Germination test was conducted in the laboratory, and the plants were grown in the green house. Plant height and cover were measured at weeks 2, 4, 6, and 8 after seeding. Transpiration rate was measured beginning eleven weeks after planting. All analysis was completed using SPSS © Statistical Software (version 18.1, SPSS Inc., Chicago, USA) [28]. Leaf tissue was extracted in 80% (v÷v) acetone for chlorophylls a and b [11], and equation of [12] for carotenoids.

### 3 Socio-economic impacts

### 3.1. Positive impacts

The positive impact of tin mining is economic [13-15]. The increase of income is shown by the number of motorcycles and cars, and from 1999 to 2011, motorcycle and car taxes increased 15-fold. The number

of people who perform the hajj pilgrimage increased almost 10% from 2001 to 2012 [16].

The income percentage for tin miners compared to overall income per month of people in Lubuk Kelik, Bangka; for ex-pepper farmers in Silip, Bangka; and for ex-rubber farmers in Bencah, Central Bangka are 90% and above. Pepper and rubber plantations contribute less than 3 % each of overall monthly income [17]. The net monthly income of fishermen in Rebo and Bubus beaches, Bangka, is just about one-third of the income of their colleagues working in tin mining [18].



**Fig. 1.** Inland and offshore study sites in Bangka Island from secondary data [19]

### 3.2 Negative impacts

Tin mining also causes societal conflicts at mining sites. Most of the conflict in both inland and offshore mining is between locals and immigrants [16; 4]. Attitude changes and conflicts are reported in the hamlets and villages of the studied area [19].

The drop-out rate from elementary to senior high school has increased. In 2011, the province of Bangka Belitung had the second-largest student drop-out rate in the country because of children's involvement in mining or following their parents when they move to new mining sites [16].


In some areas, fishermen and farmers have changed their professions to become miners. Fishing boats are modified to become mobile floating dredges in Bangka [18]. Rubber plantations and pepper plantations have been mined in some areas in Central Bangka and South Bangka [17].

Flooding in many areas of the province is believed to be caused by tin mining. The original small stream channels have been changed by the mining activity.

### 4 Water qualities and offshore biotas

Offshore tin mining has reduced water quality. This is shown by a 40% total soluble solid (TSS) increase, a 75% sedimentation rate increase, a 25% water pH decrease and a 50% dissolved oxygen (DO) increase [20] (Figure 2). It is reported that, owing to tin mining, the concentrations of lead (Pb) (0.223 ppm) and TSS in solution offshore at Batu Belubang (705 ppm) were

above the ministerial regulations of (Kepmen LH No. 51 tahun 2004) 0.008 ppm and 400 ppm respectively [21].



**Fig. 2.** Increase and decrease percentage of water quality and offshore biotas [20].

In another study, offshore mining was found to cause a 40% reduction of the number plankton species [18]. The number of species of seagrass in mined water was about 70% of the number in less mined water [18]. The dominant substrate in mined water was sand and rubble, in contrast to macroalga *Halimeda* sp. in less mined water [18].

The number of coral reef-associated fish in mined water was 30% of that in less mined water [18]. Coral reef life coverage was less than 25% in mined water compared to more than 90% in less mined water [18]. However, the growth rate of the coral reef species *Acropora digitata* transplanted to Teluk Limau Beach, Bangka, was 2.2–2.4 mm/month [22].

Because of floating small-scale artisanal tin mining units (*TI apung*), the number of fish caught has decreased, causing some fishermen not to go fishing, and no need to go further with no guarantee of a good catch [23]. The habitat changes have caused the economic benthic mollusc species *Laevistrombus canarium* L. (*siput gonggong*) of the family Strombidae to be replaced by the bivalve species *Anadara granosa* [24]. Small pelagic and demersal fish production decreased at three offshore mined sites in three regencies over the period 2009–2010 across the island, from 10 – 70% [19].

### 5 Soil degradation and inland biotas

Alluvial tin deposits - cassiterite (SnO<sub>2</sub>) - were revealed after stripping the vegetation above the upper soil and removing the non-tin deposit overburden. The extraction is done by pouring a large volume of highly pressured water over the sediment. Heavy tin ore separated from

light material by gravity. Non-tin sediment settles in a lower area with acidic pH.

Dredger exploits tin deposits located offshore up to 70 m depth with bucket wheel dredging. ASM use small traditional gravel pumps to pump tin-ore deposits to floating dredge units (*TI apung*) or modified small fishing boats.

Inland mining decreases soil properties, changing sand faction up to 97% (Table 2). The concentrations of phosphate, potassium and sodium in undisturbed land are higher than in mined soil, and are gradually decreasing as the site is abandoned [5]. C-organics are less than 2%, and the cation-exchange capacity (CEC) of tin-mined lands is very low (0.4–3.9 units) [5]. The soil temperature may reach 45°C during the day [25], and evaporation on sandy tailings may reach 4 L/m²/day or double than of undisturbed soil [26].

**Table 2.** Soil properties of 0-, 7-, 11-, 38- year old tin-mined land, and riparian forest [7].

Sites	Sand	Silt	Clay	pН	С	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	CEC
Sites		(%)		H <sub>2</sub> O	(0)	%)	(1	ng/100	g)
0	94	2	4	4.8	0.2	0	2	3	0.4
7	94	4	3	4.8	1	0.1	49	3	3.3
11	83	5	13	4.9	0.2	0	11	4	2
38	96	2	2	5.1	0.3	0	5	2	1
forest	78	13	10	4.7	1.6	0.2	22	5	5.8

C (Walk & Black); N (Kjeldahl); Cation-exchange (NH4-Acetate 1 N, pH 7); CEC (Ca+Mg+K+Na)

The water and the sediment from the washing process bring acidic material, which may reach the pH below 3. The acidity negatively affects soil flora and fauna [5]. A river that receives tin sedimentation has nearly 30% less fish species compared to a river free from tin mining [27].

Mining activity changes the vegetation structure and composition. The vegetation structure after 38 years of natural succession on old tin-mined land was less than 2%, similar to that of a riparian forest on Bangka Island [5] (Table 3). The number of arbuscular mycorrhizal fungi (AMF) spores increases with the abandonment of tin-mined land, and the number of phosphate solubilising bacteria (PSB) shows different readings with the period of abandonment [5] (Table 4).

Land recovery and coral reef transplantation are costly. The revenue from tin through land function change is lower than for non-mining land uses: protected forest, rubber plantation and beach [20]. The expenditure to convert one hectare of previously tin-mined site into rice field is estimated at Rp. 31 million, with the land preparation component representing the major portion, and almost half of the costs are for soil amendment [6].

**Table 3.** Number of individuals (A), species (B), and families (C) of 0-, 7-, 11-, 38-year old tin-mined land and riparian forest [7].

A. Number of individual per hectare

C:4	Number of individuals / ha								
Sites	seedlings	saplings	poles	trees	total				
0	0	0	0	0	0				
7	890	0	0	0	890				
11	1675	45	0	0	1720				
38	2125	55	0	0	2180				
forest	2665	4155	305	170	7295				

### B. Number of species

C!4	Number of species							
Sites	seedlings	saplings	poles	trees	total			
0	0	0	0	0	0			
7	6	0	0	0	6			
11	7	2	0	0	8			
38	15	1	0	0	16			
forest	42	66	24	11	85			

### C. Number of families

S:4	Number of families							
Sites	seedlings	saplings	poles	trees	total			
0	0	0	0	0	0			
7	4	0	0	0	4			
11	4	2	0	0	5			
38	12	1	0	0	13			
forest	24	30	14	8	44			

**Table 4.** Number of phosphate solubilizing bacteria (PSB) and arbuscular mycorrhizal fungus (AMF) of 0-, 7-, 11-, 38-year old tin-mined land and riparian forest [7].

Sites	PSB (10 <sup>5</sup> c/ g soil)	AMF spores/ 50 g soil	AMF genera	Glomus spp. (%)
0	5	1	1	100
7	10.3	69	4	67
11	6	87	4	59
38	3.2	372	3	95
forest	7	30	4	57

### 6 Physiological characters

Reclaimed mined land is a highly variable and often challenging environment for the establishment of plants. Although sites differ greatly depending on climate, local geology and reclamation methods there are several characteristics that are commonly encountered on reclaimed mines: relatively low water holding capacity due to coarsely textured soils with little or no organic content, and low nutrient availability.

Species that have been widely adopted for mine reclamation are likely to be ones that are tolerant of a wide range of environmental conditions, and in particular, those that are tolerant of low water and nutrient availability [28].

For water stressed environments, the most drought-adapted species at the individual plant scale has the lowest daily transpirational water consumption [29]. Transpiration reduction also means increasing water use efficiency [30] which may show better adaptability in unfavorable soil conditions. Other species may adapt to drought through morphological changes such as leaf size [31] or possibly anatomical characters [8] rather than through physiological mechanisms.

Although rapid growth of vegetation is often cited as a desirable characteristic to control erosion on reclaimed sites, we found that the most frequently used species tended to have a moderate growth rate. Only *Trifolium pratense* L. showed higher values that are within the typical reported range for that species [32] (Table 5).

Better able to endure the stressful, low nutrient environments of mine sites, species of moderate growth rate may have been favored for reclamation due to their ability to persist and spread. This finding provides a starting point for further testing.

**Table 5.** Summary of traits potentially desirable in reclamation ground covers in Eastern United States [28].

	A	В	C	D	E	F	G	Н
T. pratense	₽	₽	₽	₽	₽	₽	₽	₽
H. esculentus	√□	√□	<b>√</b> □			√□	√□	√□
T. repens	₽	₽	₽			₽	₽	₽
L. multiflorum		√□	<b>√</b> □		√□	√□	√□	
L. corniculatus			₽	₽		₽	₽	₽
B. napus	√□	√□				√□	√□	
B. perviridis	₽	₽	₽				₽	
L. perenne	₽		₽	₽				₽
P. virgatum			<b>√</b> □	√□		√□	√□	
V. unguiculata		₽		₽		₽	₽	
D. glomerata			₽	₽		₽		
P. fagopyrum		√□	√□		_	√□	_	
S. scoparium			<b>√</b> □			√□		<b>√</b> □
S. nutans	√□		√□		_	√□	_	

A= germination >30%, B= cover > 50% within 2 weeks, C= number of germination x pH, D= number of cover x soil type, E= number of height x soil type, F= number of transpiration x soil type, G= low transpiration rate, H= foliar pigments > 0.4 mg/g

### 7 Conclusions

The mining activities increase the wealth of the community, but the other hand they change and decrease the environmental stability, and socio economic impact stimulate horizontal conflicts. Offshore mining reduced water quality, change sea bed caused the change of biodiversity, and increase the mortality of coral reefs and their associated fishes. Inland mining activity reduces biodiversity and causes floods and damages infrastructure.

While the more economic species such as rubber, oil palm, and some fruit species are demanded, planting local tree species is challenging. An evaluation with local tree species concluded that best adapted species based on anatomical and physiological measurements was not those that showed the best performance in the field.

The work of finding physiological characteristics to predict ground cover success on mine reclamation sites has being conducted. Plant height and cover, transpiration rate, and foliar pigments may be used to select plant adaptability to mined soil. Species most widely used in reclamation tended to be perennials of moderate rate.

The first author gratefully acknowledges the funding of this research by Directorate General of Higher Education, Republic of Indonesia (Hibah Bersaing 092/SP2H/PP/DP2M/III/2007;086/SP2H/PP/DP2M/III/2008), and to thank Fulbright for funding the research (68150141). The first author would like to thank Department of Forestry, Wildlife and Fisheries, University of Tennessee kindly provided facilities to undertake the research. University of Bangka Belitung is thanked for its permission allowing the first author to conduct research. The authors acknowledge Mr. A. Akbar and Ms. W.E.A. Putri, Dr. Z.Q. Liu, and Mr. J. Seaton for the valuable data and help.

### References

- 1. IDH Tin working group communiqué To share recent study findings and consult Indonesian stakeholders about their sustainability views and priorities for action (2013).
- Bangka Pos, 2013, Idil: arogansi bupati akan dibayar mahal. Bangka Pos 20 Maret 2013, viewed 19 April 2013.
- 3. International Tin Research Institute (ITRI) Indonesia's position in the global tin industry. *Proceedings ITRI Indonesia Tin Forum*, 11 December 2013, Pangkalpinang (2013).
- Walhi Friends of the Earth Indonesia, Mari buat Bangka Belitung lebih baik. *Proceedings ITRI Indonesia Tin Forum*, 11 December 2013, Pangkalpinang, Indonesia.
- 5. E. Nurtjahya, D. Setiadi, E. Guhardja, M. Muhadiono, Y. Setiadi, Succession on tin-mined land in Bangka Island, Blumea, **54** (2009), no. 1–3, pp. 131–138.
- E. Nurtjahya, M.M. Nur, E. Mulyono, Rice field cultivation on tin-mined land in Bangka Island, Indonesia. *Proceedings of Mine Closure 2009*, A.B. Fourie, and M. Tibbett (eds), September 2009, Perth,2009 Australian Centre for Geomechanics, Perth, pp. 549-560.
- 7. E. Nurtjahya, Revegetasi lahan pasca tambang timah dengan beragam jenis pohon lokal di Pulau Bangka, Dissertation, Bogor, Institut Pertanian Bogor, 163 pp, (2008).
- 8. E. Nurtjahya, R. Robika, D. Dorly, Can anatomical and physiological characters predict plant adaptation on tin-mined land in Bangka Island? *Proceedings of the 6<sup>th</sup> International Conference on Mine Closure 2011*, A.B. Fourie, and M. Tibbett and A. Beersing (eds), September 2011, Alberta, Canada,2011 Australian Centre for Geomechanics, Perth, pp. 75-83.
- 9. Badan Pusat Statistik (BPS), *Bangka Belitung dalam Angka*, Pangkalpinang (2012), Indonesia.
- PT Timah Tbk, ANDAL, RKL, dan RPL. Kegiatan penambangan timah dan pasir laut di perairan Pulau Bangka Kabupaten Bangka, Propinsi Sumatera Selatan (1997).
- Z.D. Sesták, J. Čatskŷ, P.G. Jarvis, Determination of chlorophylls a and b, in *Plant photosynthetic production: Manual of Methods*, Z. Sesták, J. Čatskŷ, and P.G. Jarvis, eds., Junk NV, The Hague (1971), pp. 672–701. B.H.

- 12. Davies, Analysis of carotenoid pigments. In: T.W. Goodwin (ed.) *Chemistry and Biochemistry of Plant Pigments*, pp. 489-532. Academic Press, New York, (1965).
- 13. L. Juniarti, *Disharmonisasi keluarga pada pekerja tambang inkonvensional (TI) di Dusun Parit 19 Kecamatan*, Dissertation, Jurusan Sosiologi, Universitas Bangka Belitung, Pangkalpinang, Indonesia, (2014).
- Indra, Tambang inkonvensional dan implikasinya terhadap perilaku sosial masyarakat di Desa Tempilang Kabupaten Bangka Barat, Dissertation, Jurusan Sosiologi, Universitas Bangka Belitung, Pangkalpinang, Indonesia, (2013).
- B. Romeo, Tambang inkonvensional dan perubahan perilaku ekonomi masyarakat Desa Semulut Kecamatan Jebus, Dissertation, Jurusan Sosiologi, Universitas Bangka Belitung, Pangkalpinang, Indonesia, (2011).
- E. Erman, Dampak penambangan timah dan respon masyarakat local. *Proceeding ITRI Indonesia Tin Forum*, 11 December 2013, Pangkalpinang, Indonesia, (2013).
- 17. E. Nurtjahya, F. Agustina, W.A.E. Putri, Neraca ekologi penambangan timah di Pulau Bangka Studi kasus pengalihan lahan di ekosistem darat, Berkala Penelitian Hayati, **14**(1) (2008), pp. 29–38.
- 18. E. Nurtjahya, F. Agustina, A. Akbar, Kajian manfaat sosial ekonomi penambangan timah inkonvensional dan kerusakan lingkungan dan keanekaragaman hayati yang ditimbulkannya di Pulau Bangka, Laporan Penelitian Hibah Bersaing, Universitas Bangka Belitung (2008), 70 p.
- E. Nurtjahya, F. Agustina, Managing the socioeconomic impact of tin mining on Bangka Island, Indonesia — preparation for closure. *Proceedings Mine Closure 2015*, A.B. Fourie and M. Tibbett, L. Sawatsky and D. Van Zyl (eds), Juni 2015, Vancouver, Canada, Australian Centre for Geomech anics, Perth, pp. 817–826.
- E. Nurtjahya, U. Umroh, F. Agustina, Impact of tin mining on the biota of Bangka Island, Indonesia a proof to convince the tin supply chain of smartphone companies. *Proceedings Mine Closure 2014*, A.B. Fourie, M. Tibbett, and I. Weiersby (eds), October 2014, Johannesburg, SouthAfrica, Australian Centre for Geomechanics, P erth (to be published).
- 21. H. Wahyuni, S.B. Sasongko, D.P. Sasongko, Kandungan logam berat pada air, sedimen dan plankton di daerah penambangan masyarakat Desa Batu Belubang Kabupaten Bangka Tengah, in Proceedings Seminar Nasional Pengelolaan Sumberdaya Alam dan Lingkungan (2013), 489– 494.

- 22. H. Sodikin, *Pertumbuhan Acropora digitata pada transplantasi karang di pantai Teluk Limau Sungailiat Provinsi Kepulauan Bangka Belitung*, Dissertation, Pangkalpinang: Jurusan Perikanan Universitas Bangka Belitung (2011), 36 p.
- 23. P. Pratama, *Alih profesi nelayan menjadi penambang timah di Lingkungan Nelayan 2 Sungailiat*, Dissertation, Jurusan Sosiologi, Pangkalpinang: Universitas Bangka Belitung (2014).
- 24. F. Yulianda, F. Achmad, A.H. Armin, H. Sri, K. Kusharjani, *Ekologi ekosistem perairan laut tropis*, PUSDIKLAT Kehutanan Departemen Kehutanan RI, Bogor: SECEM Korea International Cooperation Agency (2009).
- 25. E. Nurtjahya, D. Setiadi, E. Guhardja, M. Muhadiono, Y. Setiadi, Revegetation of tin-mined land using various local tree species in Bangka Island, Indonesia. Proceedings of the 2008 National Meeting of the American Society of Mining and Reclamation New Opportunities to Apply Our Science, R.I. Barnhisel (ed), June 2008, Lexington, USA, ASMR, pp. 739-755.
- 26. E. Nurtjahya, Ex tin-mined research a local tree species search (2010) (unpublished).
- K. Muslih, E.M. Adiwilaga, S. Adiwibowo, Pengaruh penambangan timah terhadap keanekaragaman ikan sungai dan kearifan local masyarakat di Kabupaten Bangka. *Prosiding* Pertemuan Ilmiah Tahunan MLI I (2013).
- 28. E. Nurtjahya, J. Franklin, Some physiological characteristics to estimate species potential as a mine reclamation ground cover. Int. J. Min. Reclam. Environ (2016) (to be published).
- 29. C. Tong, J.-Z. Gong, R. Marrs, I. Zhang, W.-Q Wang, Pattern of transpiration of four shrub species and water consumption from shrub stands in an eco–reclamation catchment in northwest China, Arid Land Res. Manag. 22 (2008), pp. 242–254.
- 30. A.M. Gonzalez-Rodriguez, Z. Baruch, D. Palomo, G. Cruz-Trujillo, M.S. Jimenez, D. Morales, Ecophysiology of the invader Pennisetum setaceum and three native grasses in the Canary Islands, Acta Oecol. **36** (2010), pp. 248–254.
- 31. F.M. DaMatta, Exploring drought tolerance in coffee: a physiological approach with some insights for plant breeding, Braz. J. Plant Physi. **16** (2004), pp. 1–6.
- 32. L. Simova-Stoilova, K. Demirevska, A. Kingston-Smith, U. Feller, Involvement of the leaf antioxidant system in the response to soil flooding in two Trifolium genotypes differing in their tolerance to water logging, Plant Sci. **183** (2012), pp. 43–49.

6

Corresponding author: eddy\_nurtjahya@yahoo.com

### Preface

Sriwijaya International Conference on Engineering, Science and Technology 2016 (SICEST 2016), is the first international conference organized by the Faculty of Engineering Universitas Sriwijaya. Theme of the event was "Building a better future through innovation in engineering, science and technology".

Through SICEST conference, the organizing committee has successfully recognized audience to the 300 presented articles from 650 submissions. The forum was arranged in 7 symposia and attended by more than 450 participants. They who contributed were scientists, researchers, engineers, students, professional stakeholders, consultants, government officials, marketers and professional users. All the symposia attended by various participants from 20 countries and from 137 reputable affiliations.

In order to make SICEST conference proceedings widely usable by the scientific community, we are proud having a partnership with MATEC as one of reputable partner with highly flexible publishing solutions. *MATEC Web of Conferences* are believed could increase the visibility and enhance the profile of SICEST conference, as well as to highlight the work of the paper contributors. It also provides a long-term open-access record of the conferences.

After a thorough consideration based on the quality of paper submitted, SICEST Scientific Committee has selected 115 articles for publication via Matec Web of Conferences. We have conducted peer-review in accordance with the policy on publishing integrity of the journal, in order to safeguard good scientific practice in publishing.

All submitted articles to SICEST 2016 were gone through a double blind peer review by competent reviewers. After that stage, many papers were invited to revise and resubmit in maximum of 3 weeks. The final decision was made on the basis of the revised paper only. Reviews were made explicit how the reviewers think the paper could be improved.

We express our deepest gratitude to all members of the international scientific committee, keynote and invited speakers, reviewers, sponsors, all paper contributors, participants in the conference and members of the SICEST organizing committee for their valuable contribution to the successful organization of SICEST2016.

### Dr. Muhammad Faizal

Executive Chairman of SICEST 2016 Sriwijaya International Conference on Engineering, Science and Technology

Palembang-Indonesia, February 2017





## Statement of Peer review

In submitting conference proceedings to MATEC Web of Conferences, I certify to the Publisher that I adhere to the **Policy on Publishing Integrity** of the journal in order to safeguard good scientific practice in publishing.

- 1. All articles have been subjected to peer review administered by the proceedings editors.
- 2. Reviews have been conducted by expert referees, who have been requested to provide unbiased and constructive comments aimed, whenever possible, at improving the work.
- 3. Proceedings editors have taken all reasonable steps to ensure the quality of the materials they publish and their decision to accept or reject a paper for publication has been based only on the merits of the work and the relevance to the journal.


Title, date and place of the conference

Sriwijaya International Conference on Engineering, Science and Technology (SICEST 2016)

November 9-10, 2016 Santika Hotel Bangka Island, Indonesia


Proceedings editor(s):

Prof. Iskhaq Iskandar Suryadi Ismadji, PhD Tuty Emilia Agustina, PhD Irsyadi Yani, PhD Dr. Leily Nurul Komariah Dr. Saloma Hasyim



### https://www.matec-

 $\frac{conferences.org/articles/matecconf/abs/2017/15/contents/contents.html \# section\_10.1051/matecconf/201710101001$ 



### https://www.matec-

 $\frac{conferences.org/articles/matecconf/abs/2017/15/contents/contents.html \# section\_10.1051/matecconf/201710101001$ 



# Authors Affiliations

versity of Science & Technology - Taiwan

sity of Agriculture and Forestry - Vietnam ersity Technology Thonburi - Thailand

University of Tennessee – USA	Yokohama National University – Japan	Tokyo University of Agriculture and Technology - Japan
Ecole Centrale de Nantes – France	De La Salle University – Philippines	National Taiwan University of Science & Technology - Tai
University of Twente – Netherlands	Can Tho University – Vietnam	King Mongkut's University Technology Thonburi - Thailan
Firat University – Turkey	Academia Sinica – Taiwan	Thai Nguyen University of Agriculture and Forestry – Vietn
Obafemi Awolowo University - Nigeria	Chonbuk National University - Korea	Nanyang Technological University - Singapore
Federal University of Technology - Nigeria	The University of Queensland - Australia	Korea Institute of Industrial Technology - Korea
King Saud University – Saudi Arabi	Sejong University – Korea	University of MARA Technology Malaysia
Sultan Qaboos University - Oman	The University of Hong Kong - Hong Kong	Ethiopian Health and Nutrition Research Institute - Ethiopi
KE College – India	Universiti Tun Hussein Onn – Malaysia	OTHERS
Bharata Martha College – India	Universiti Kebangsaan Malaysia – Malaysia	Ecophile – Korea
Curtin University - Australia	Universiti Malaysia Pahang – Malaysia	German – Malaysian Institute - Malaysia
Universiti Teknologi Malaysia – Malaysia	Universiti Sultan Zainal Abidin – Malaysia	PUB (Singapore National Water Agency) - Singapore
Tokyo University of Science – Japan	National University of Singapore - Singapore	Surbana Jurong Consultants Pte. Ltd - Singapore

Surbana Jurong C	Surbana Jurong Consultants Pte. Ltd – Singapore	
	re – Singapore	



COMPANY	COMPANY AND OTHERS
PT. Krakatau Steel	PT. Pamapersada Nusantara
PT Wijaya Karya Beton, Tbk	Pertamina Upstream Technology Center
Nanocenter Indonesia	PT. Batamindo Investment Cakrawala
PT Adaro Indonesia	PT. Newmont Nusa Tenggara
GeoPangea Research Group	Sekampung River Organization (Mesuji)
PT Teno Indonesia	Watershed Management Technology Centre
PT Ecophile Green Indonesia	PT. Sumiasih Oleochemicals



## 

Prof. Ahmad Fauzi Ismail	Prof. Osman Adiguzel	Dr. Suryadi Ismadji	Prof. Iskhaq Iskandar, Ph.D	Prof. Safian Sharif	Tuty Emilia Agustina, PhD	Nurly Gofar, PhD

Vanyang Technological University-Singapore Universiti Teknologi Malaysia-Malaysia Universiti Teknologi Malaysia-Malaysia Widya Mandala University-Indonesia Sriwijaya University-Indonesia Sriwijaya University-Indonesia Firat University-Turkey

# 

ryanti F Nata, Ph.D (Lambung Mangkurat Univ) Makbul Anwari, PhD (King Abdul Aziz University, Or. Budhi Kuswan Susilo, (Sriwijaya University) Dr. Leily Nurul Komariah (Sriwijaya University) Dr. Wiryanto Dewobroto (Pelita Harapan Univ) Abu Bakar Siddik, PhD (Sriwijaya University) Dr. Poedji Loekitowati (Sriwijaya University) Dr. Saloma (Sriwijaya University-Indonesia) Dr. Widya Iskandar (Sriwijaya University) Dr. Joni Arliansyah (Sriwijaya University) Dr. Tri Kurnia Dewi (Sriwijaya University rsyadi Yani, Ph.D (Sriwijaya University) Or. Elda Melwita (Sriwijaya University) Dr. Erna Yuliwati (Sriwijaya University) Dr. Mohamed Iqbal (IUUM Malaysia) Dr. Ivan Affandi (SPS Technology) Prof. Solehudin Husein (UTM) Prof. Andreas Wibowo (PU) (ingdom of Saudi Arabiia) Dr. Agung Murti Nugroho ndonesia) Amrifan Saladin Mohruni, PhD (Sriwijaya University) Assoc. Prof. Nik Rumzi bin Nik Idris (UTM Malaysia) Dr. Cynthia Fabian Madrazo (De La Salle University, Salahuddin Husein, PhD (Gadjah Mada University) Prof. Siti Nurmaini (Sriwijaya University-Indonesia) Dr. Dina Muthmainah (Ministry of Marine Affairs & Setyawati Yani, Ph.D (UMI Makasar-Indonesia) Dr. Nouruddeen Bashir Umar (UTM-Malaysia) Dr. Noor Azlinda binti Ahmad(UTM Malaysia) Prof.Ahmed Fazary (King Khalid Univ- Saudi Dr. Ildrem (Padjajaran University-Indonesia) Prof. Muhammad Said (Sriwijaya University) Dr. Ardiyanshah Syahrom (UTM – Malaysia) Dr. Dasapta Erwin Irawan (ITB - Indonesia) Prof. Husni Husin (Syiah Kuala University) Joko Nugroho, Ph.D. (ITB-Indonesia) Dr. Agus Subekti (LIPI -Indonesia) Prof. Arief Junaidi (ITS-Indonesia) Dr. Esa Prakasa (LIPI-Indonesia) Fisheries Republic of Indonesia) Phillipine) Arabia) 00000 000000000000 Or. Ruddy Kurniawan (Andalas University-Indonesia) Dr. Risfidian Mohadi (Sriwijaya University-Indonesia) Prof. Siti Nurmaini (Sriwijaya University-Indonesia) Moh Seddik Meddah, PhD (Sultan Qaboos Univ-Dr. Phil Lewis (Oklahoma State University-USA) Rudi Heriansyah, PhD (Umm Al-Quro University, Prof. Wen Chien Lee (National Chung Cheng Prof. Subriyer Nasir (Sriwijaya University) Assoc. Prof. Jafri bin Din (UTM Malaysia) Dr. SD. Sumbogo Murti (BPPT-Indonesia) Prof. Anis Saggaf (Sriwijaya University) Prof. Praveen Linga (NUS - Singapore) Prof. Shafian Sharif (UTM-Malaysia) Dhemi Harlan, Ph.D (ITB-Indonesia) Prof. Zainal Salam (UTM Malaysia) Dr. Rizgon Fajar (BPPT-Indonesia) Prof. Yazid Bindar (ITB-Indonesia) Dr. Dedi Apriadi (ITB-Indonesia)) Kingdom of Saudi Arabia) University-Taiwan) 00000000000 

Dr. Heni Fitriani (Sriwijaya University)

# ORGANIZING COMMITTEI

**Executive Chairman** 

Dr. Muhammad Faizal

Technical Committee Chairman

Amrifan S Mohruni, PHD

Program & Conference Manager

Dr. Leily Nurul Komariah

Publication & Paper Management Tuty Emilia Agustina, Ph.D

Elda Melwita, Ph.D

Ika Yuliantina, Marwani Zachrie, Lia Cundari, Heriyanto, Hermawan Yuliansyah, Deni Chaerudin

Symposium Chair Dr. Saloma Hasyim

Dr. Diah Kusuma Pratiwi Heni Fitriani, Ph.D Irsyadi Yani, Ph.D

Moh. Abu Bakar Siddik. Ph.D Dr. Budhi Kuswan Soesilo Widya Fransiska F, Ph.D

Secretariate

Bazlina Dawami, Cindi Ramayanti, Winny Andalia, Alexander Ivan, Sucia Aprisah, Yangia Septa, Mia Odina,

Program Staff:

Irwin Bizzy, Dr. Susila Arita, Baharuddin, Yoga Permana, Amir Arifin, PhD

Cooperation & Sponsorship: Farida Ali, Dr. Restu Juniah, Baharuddin, Dr. Hanafiah,

Production Team: Amrillah Nugrahsyah, Ihsan Riady, Bella Febrianti, Edriansyah, Chandra Kartawijaya, Agusyanda, Maidawati

Secretariate Office: Graha Bukit Asam Sriwijaya University 1st Floor JI. Srijaya Negara Bukit Besar Palembang