



IJC 2019

by Andri Kurniawan

Submission date: 09-Sep-2021 01:24PM (UTC+0700)

Submission ID: 1644321051

File name: Andri_IJC 2019.pdf (300.34K)

Word count: 4554

Character count: 22830

Short Communication:

Andri Kurniawan^{1,2,*}, Oedjijono², Tamad³, and Uyi Sulaeman⁴

¹Department of Aquaculture, University of Bangka Belitung,
Jl. Balunijuk, Merawang, Bangka, Bangka Belitung 33172, Indonesia

²Faculty of Biology, Jenderal Soedirman University,
Jl. Dr. Suparno no. 63, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia

³Department of Agriculture, Jenderal Soedirman University,
Jl. Dr. Suparno no. 61, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia

⁴Department of Chemistry, Faculty of Science and Technology, Jenderal Soedirman University,
Jl. Dr. Suparno no. 61, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia

* Corresponding author:

tel: +62-81351714747

email: andri-kurniawan@ubb.ac.id

Received: March 1, 2018

Accepted: July 17, 2018

DOI: 10.22146/ijc.33613

Abstract: The heavy metals distribution of ex-tin mining ponds were investigated. The time chronosequence was determined at the pond of age < 1 year (Station A), the pond of age 5–10 years (Station B), and the pond of age > 15 years (Station C). The results showed sixteen heavy metals of As, Co, Cu, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V could be detected in the ponds. The As, and in Station C showed higher concentration compared to the metals such as Cr, Fe, Hf, Sn, Ta, Te, and V in Station A and Station B showed higher concentration compared to the Station C. The positive, negative, and dynamic correlation pattern could be found in distribution of heavy metal to time chronosequence. The concentration of Ta and V showed a positive correlation because their concentration decrease, whereas concentration of As, Cu, Ga, Mn, and Zn showed a negative correlation because their concentration increase along in time chronosequence. The dynamic correlation could be found that concentration of Co, Ni, Pb, Sn, and Th decrease from Station A to Station B and then increase in Station C, whereas concentration of Cr, Fe, Hf, and Te increase from Station A to Station B and then decrease in Station C.

Keywords: distribution pattern; heavy metals; chronosequence; ex-tin mining ponds

INTRODUCTION

Tin mining activity is an anthropogenic activity that has contributed to the ecological problems. It has been known as the major sources of heavy metal contamination such as heavy metals in as one of water resource were contaminated by heavy metals that can be accumulated in food chains and secondary activities of the water [1-3]. Some of the Mn, Fe, Ni, Sn, As, Cd were detected in site of ex-tin mining [4-5].

The each of waters had their that [6]. These conditions were influenced by ecological succession in time chronosequence [7] and the ecological changes of ex-tin mining ponds have occurred for a long time [8]. The implication of ecological changes as long as time chronosequence was interactions and reactions of chemical complexation between organic compounds, environmental factors, and inorganic compounds, include presence of heavy metals [9].

The researches of heavy metals in ex-tin mining ecosystems have been done, however, not for correspondence of heavy metals distributions and ex-tin mining ponds in time chronosequence. While an understanding this correspondence was crucial to predict ecosystems condition and to detect environment change. The understanding becomes an important part of aquatic ecosystem management as water sources for secondary activities and habitable for organisms.

■ **EXPERIMENTAL SECTION**

Materials

The study areas were located in Bangka Regency, Bangka Belitung Archipelago Province of Indonesia. The considered study sites cover three ponds of ex-tin mining with chronosequence ranging in time. The study areas were encoded as Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years).

The coordinate of Station A were 01°59' S in points 36.0"; 36.2"; 36.4"; 36.5"; 36.6" and 106°06' E in points 36.5"; 36.9"; 37.3"; 37.4"; 37.5". The coordinate of Station B were 01°59' S in points 41.3"; 41.4"; 41.5"; 42.4"; 42.5" and 106°06' E in points 39.2"; 39.5"; 41.4"; 42.7"; 43.1". The coordinate of Station C were 01°55' S in points 40.9"; 58.9"; 59.1"; 59.2"; 59.5" and 106°06' E in points 19.5"; 19.7"; 19.9"; 22.4"; 29.2".

In the each of research stations points, water sampling was done < 4 m in depth (station code A.1; B.1; and C.1) and composite sampling were done to water and sediment > 4 m in depth (station code A.2; B.2; and C.2).

Instrumentation

The water samples and composite samples were analyzed pH value by pH meter (PH-009(I)-A) Sun Care. The speciation and concentration of heavy metals analyzed by X-Ray Fluorescence (Rigaku) with three light spreader metals of copper (Cu), molybdenum (Mo), and aluminum (Al). Also, the pH and moisture of soil around the ponds were measured by pH-moisture meter. The correspondence between and

the was analyzed by PAST (*Palaentological Statistics*) 3.

■ **RESULTS AND DISCUSSION**

pH Value of Water and Soil

Tin mining activity had generated the ex-tin mining ponds as one of the water resources. However, water quality in the ponds had serious ecological problem. The ex-mining activity produced acid mine drainage (AMD) that can caused the long-term impairment of water quality and biodiversity. The pH value of ex-tin mining water and soil pH value indicated that the acid could be found in Station A (pond in age < 1 year) and Station B (pond in age 5–10 years) and the neutral value could be found in Station C (pond in age > 15 years). In the each of research stations points, water samples < 4m in depth indicated more acid than the composite samples (water and sediment > 4 m in depth) (Fig. 1).

The () was [10]. In acid drainage, presence and mobility of elements such , As, , Zn, also ionic activity and sulphide




Fig 1. Soil and water pH value of . The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth)

minerals oxidation caused pH value decreasing [11-12]. Furthermore, the low pH condition implicated in oxidation state, toxicity, and mobility of elements [9].

In long time chronosequence more than 15 years in the Station C caused changes water quality of ex-tin mining ponds were more stable than Station A and Station B. The indicator for it was pH value increased to neutral conditions (7.0-7.4) and other parameters that indicated water quality was more habitable for organisms' life such as fishes. The pH value and other parameters changes, indirectly, caused ecological productivity increasing. The chronosequence in long time caused changes in nutrient availability and resource supply for ecosystem productivity [13].

The pH value the determination of . Furthermore, acid mine drainage had a major contribution in heavy metal contamination [14]. There was

distribution where pH value decreasing caused increasing in heavy metal desorption [15].

The Pattern of Heavy Metals Distribution

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C) (Table 1).

There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that . The average of As concentration were 1.7475 ppm (Station C) > 1.6075 ppm (Station B) > 1.0775 ppm (Station A). The concentration of Co were 3.32 ppm (Station C) > 2.525 ppm (Station A) > not detected (Station B). The concentration of Cu were 6.375 ppm (Station C) > 5.4825 ppm (Station B) > 4.9775 ppm (Station A).

Table 1. Distribution in water

No	Heavy Metals	Heavy Metals Concentration (ppm)					
		In Station A		In		In	
		.1	A.1	B.1	.2	C.1	C.2
1	As	nd	2.155	nd	3.215	2.4	1.095
2	Ga	15	4.275	nd	4.47	9.76	nd
3	V						.67
4	Te			102	nd	nd	5.8
5	Cr	nd	4.735	nd	4.945	nd	0.67
6	Zn	nd	6.05	nd	8.1	8.17	3.905
7	Sn	47	82.8	50.3	51.4	61.8	57.15
8	Mn	nd	13.05	nd	13.6	30.8	14.85
9	Fe	21.4	1481.3	148	2344	1590	849.05
10	Co	nd	5.05	nd	nd	6.64	nd
11	Ni	nd	4.09	nd	2.875	3.48	1.905
12	Cu	3.08	6.875	3.19	7.775	8.94	3.81
13	Pb	nd	6.7	nd	6	8.21	4.62
14	Th	nd	4.605	nd	4.38	13.5	7.05
15	Hf	6.71	3.21	8.21	3.32	nd	9.38
16	Ta	4.94	2.615	4.16	2.995	nd	2.805

*) nd (not detected). The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth)

The concentration of Ga were 4.88 ppm (Station C) > 2.235 ppm (Station B) > 2.1375 ppm (Station A). The concentration of Mn were 22.825 ppm (Station C) > 6.8 ppm (Station B) > 6.525 ppm (Station A). The concentration of Ni were 2.6925 ppm (Station C) > 2.045 ppm (Station A) > 1.4375 ppm (Station B). The concentration of Pb were 6.415 ppm (Station C) > 3.35 ppm (Station A) > 3.0 ppm (Station B). The concentration of Th were 10.275 ppm (Station C) > 2.3025 ppm (Station A) > 2.19 ppm (Station B). The concentration of Zn were 6.0375 ppm (Station C) > 4.05 ppm (Station B) > 3.025 ppm (Station A).

In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V. The average of Cr concentration were 2.4725 ppm (Station B) > 2.3675 ppm (Station A) > 0.335 ppm (Station C). The concentration of Fe were 1246 ppm (Station B) > 1219.53 ppm (Station C) > 751.35 ppm (Station A). The concentration of Hf were 5.765 ppm (Station B) > 4.96 ppm

(Station A) > 4.69 ppm (Station C). The concentration of Sn were 64.9 ppm (Station A) > 59.475 ppm (Station C) > 50.85 ppm (Station B). The concentration of Ta were 3.7775 ppm (Station A) > 3.5775 ppm (Station B) > 1.4025 ppm (Station C). The concentration of Te were 51 ppm (Station B) > 2.9 ppm (Station C) > not detected (Station A). The concentration of V were 0.42 ppm (Station A) > 0.335 ppm (Station C) > not detected (Station B).

Their concentration in [redacted] showed three patterns distribution that a positive correlation, a negative correlation, and a dynamic correlation. The positive correlation indicated that concentration of heavy metals decreased along chronosequence in time in ex-tin mining ponds. The negative correlation indicated that concentration of heavy metals increased along chronosequence in time in ex-tin mining ponds. The dynamic correlation indicated that concentration of




Fig 2. Distribution of [redacted] of [redacted]. The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth)

heavy metals on decreased or increased along chronosequence in time in ex-tin mining ponds.

The elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. The elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increasing along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then increased concentration in ponds with age > 15 years. The elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then decreased concentration in ponds with age > 15 years (Fig. 2).

The chronosequence in long time caused changes in heavy metals concentration were decreased as the primary succession progression [16]. The low pH value implicated to dissolved oxygen (DO) [17] and low DO in acid mine drainage implicated to biological oxygen demand (BOD) by microorganisms for their growth and other ecological factor changes such as nitrogen and phosphate [7,18-19]. Nitrogen (ammonia, nitrite, and nitrate) and phosphate were

primary succession, led to changes in the biogeochemical cycles of nutrients in ponds [20-22], indirectly implicated to dissolved and suspended solid [23-24] and ecosystem eutrophic levels [25-31]. Therefore, pH value changes, availability of elements included heavy metals, environmental characteristics, and eutrophication had a positive inter-relationships in their correspondence.

These conditions, however, had a negative correlation to heavy metals residue decreasing in Station C. The heavy metals in Station C were dominated As, Zn, Mn, Co, Ni, Cu, Pb, and Th were higher than Station A and Station B, although these metals and include Ga, Cr, Sn, Fe, Hf, and Ta were detected in all of the research station. The results indicated that chronosequence in time of ponds caused ecological changes and had positive correlations, but not for

Correspondence Analysis Distribution

The correspondence analysis also showed that there was not a positive correlation between in water and




Fig 3. Correspondence analysis between distribution of of . The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth)

heavy metals distribution (Fig. 3). The ecological changes in chronosequence succession of ex-tin mining ponds had not significantly affect on speciation and concentration of heavy metals. Thus, there was the other factor that had affected

soil water contributed distribution heavy [32]. were , include and the [15]. The clay mud were found to be dominant in the sediment of Station A and Station B, however, sand was found to be dominant in Station C.

The clay and mud had a positive correlation and sand had a negative correlation to a presence, affinity, and minerals structure [33-34]. The clay and mud had a (CEC), and of were also adsorbed the , accumulated, carried diagenetically [35-36]. Therefore, Station A Station B were lower than Station C.

CONCLUSION

1 affected ecological dynamic such as pH value change and heavy metals distribution. The pH value of ex-tin mining ponds moved from acid to neutral condition, however, it had been taken for a long time of the chronosequence. The consequence of low pH value can implicated to microorganisms' growth, nutrients, and environment factors in primary succession. Further, it implicated to heavy metals interaction, whereas the heavy metals residue should be in low concentration in long time chronosequence of ex-tin mining ponds.

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and

neutral condition (in Station C). There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that Ga, Th, addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that

There were three patterns in that a positive correlation, a negative correlation, and a dynamic correlation. The chronosequence in time had affects on heavy metals distribution. Elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. Elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increased along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5-10 years and then increased concentration in ponds with age > 15 years. Elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5-10 years and then decreased concentration in ponds with age > 15 years.

The chronosequence in time can be a factor that contributed to heavy metals distribution. However, the sediment factor had a significant effect to distribution of heavy metals in the ex-tin mining ponds, where clay and mud were more effective than sand in accumulation, adsorption, and interaction with heavy metals and carried them diagenetically to bottom sediments.

ACKNOWLEDGMENTS

7 scholarship 13 of) for the funding with Project No: 120.G/UN50.3.1/PP/2018.

REFERENCES

[1] de Silva, A.A.L., de Carvalho, M.A.R., de Souza, S.A.L., Dias, P.M.T., Filho, R.G.S., Saramago, C.S.M., Bento, C.A.M., and Hofer, E., 2012, Heavy

- metal tolerance (Cr, Ag and Hg) in bacteria isolated from sewage, *Braz. J. Microbiol.*, 43 (4), 1620–1631.
- [2] Guan, Y., Shao, C., and Ju, M., 2014, Heavy metal contamination assessment and partition for industrial and mining gathering areas, *Int. J. Environ. Res. Public Health*, 11 (7), 7286–7303.
- [3] Kurniawan, A., 2017, Chronosequence effect of post tin mining ponds to metals residu and micro ecosystem change, *Omni-Akuatika*, 13 (1), 60–65.
- [4] Ashraf, M.A., Maah, M.J., and Yusoff, I., 2012, Speciation of heavy metals in the sediments of former tin mining catchment, *Iran. J. Sci. Technol. Trans. A Sci.*, 36 (2), 163–180.
- [5] Daniel, V.N., Chudusu, E.S., Chup, J.A., and Pius, N.D., 2014, Variations of heavy metals in agricultural soils irrigated with tin water in Heipang District of Barkin Ladi, Plateau State, Nigeria, *Int. J. Sci. Technol.*, 3 (5), 255–263.
- [6] Ashraf, M.A., Maah, M.J., and Yusoff, I., 2012, Morphology, geology and water quality assessment of former tin mining catchment, *Sci. World J.*, 2012, 369206.
- [7] Urbanová, M., Kopecký, J., Valášková, V., Ságová-Marečková, M., Elhottová, D., Kyselková, M., Moënné-Loccoz, Y., and Baldrian, P., 2011, Development of bacterial community during spontaneous succession on spoil heaps after brown coal mining, *FEMS Microbiol. Ecol.*, 78 (1), 59–69.
- [8] Kurniawan, A., 2016, Microorganism communities response of ecological changes in post tin mining ponds, *RRJoMV*, 6 (1), 17–26.
- [9] Violante, A., Cozzolino, V., Perelomov, L., Caporale, A.G., and Pigna, M., 2010, Mobility and bioavailability of heavy metals and metalloids in soil environments, *J. Soil Sci. Plant Nutr.*, 10 (3), 268–292.
- [10] Akcil, A., and Koldas, S., 2006, Acid mine drainage (AMD): causes, treatment and case studies, *J. Cleaner Prod.*, 14 (12-13), 1139–1145.
- [11] Tan, G.L., Shu, W.S., Hallberg, K.B., Li, F., Lan, C.Y., and Huang, L.N., 2007, Cultivation-dependent and cultivation-independent characterization of the microbial community in acid mine drainage associated with acidic Pb/Zn mine tailings at Lechang, Guangdong, China, *FEMS Microbiol. Ecol.*, 59 (1), 118–126.
- [12] Campaner, V.P., Luiz-Silva, W., and Machado, W., 2014, Geochemistry of acid mine drainage from a coal mining area and processes controlling metal attenuation in stream waters, Southern Brazil, *An. Acad. Bras. Ciênc.*, 86 (2), 539–554.
- [13] Bardgett, R.D., Bowman, W.D., Kaufmann, R., and Schmidt, S.K., 2005, A temporal approach to linking aboveground and belowground ecology, *Trends Ecol. Evol.*, 20 (11), 634–641.
- [14] Ahmad, A.K., and Al-Mahaqeri, S.A., 2015, Human health risk assessment of heavy metals in fish species collected from catchments of former tin mining, *Int. J. Res. Stud. Sci. Eng. Technol.*, 2 (4), 9–21.
- [15] Zeng, F., Ali, S., Zhang, H., Ouyang, Y., Qiu, B., Wu, F., and Zhang, G., 2011, The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants, *Environ. Pollut.*, 159 (1), 84–91.
- [16] Huang, L.N., Tang, F.Z., Song, Y.S., Wan, C.Y., Wang, S.L., Liu, W.Q., and Shu, W.S., 2010, Biodiversity, abundance, and activity of nitrogen-fixing bacteria during primary succession on a copper mine tailings, *FEMS Microbiol. Ecol.*, 78 (3), 439–450.
- [17] Abdel-Raouf, N., Al-Homaidan, A.A., and Ibraheem, I.B.M., 2012, Microalgae and wastewater treatment, *Saudi J. Biol. Sci.*, 19 (3), 257–275.
- [18] Tschërko, D., Hammesfahr, U., Zeltner, G., Kandeler, E., and Böcker, R., 2005, Plant succession and rhizosphere microbial communities in a recently deglaciated alpine terrain, *Basic Appl. Ecol.*, 6 (4), 367–383.
- [19] Fierer, N., Nemergut, D., Knight, R., and Craine, J.M., 2010, Changes through time: Integrating microorganisms into the study of succession, *Res. Microbiol.*, 161 (8), 635–642.
- [20] Quirós, R., 2003, The relationship between nitrate and ammonia concentrations in the pelagic zone of lakes, *Limnetica*, 22 (1-2), 37–50.
- [21] Michalski, R., and Kurzyca, I., 2006, Determination of nitrogen species (nitrate, nitrite and ammonia

- ions) in environmental samples by ion chromatography, *Pol. J. Environ. Stud.*, 15 (1), 5–18.
- [22] Mihale, M.J., 2015, Nitrogen and phosphorus dynamics in the waters of the Great Ruaha River, Tanzania, *J. Water Resour. Ocean Sci.*, 4 (5), 59–71.
- [23] Akan, J.C., Audu, S.I., Mohammed, Z., and Ogugbuaja, V.O., 2013, Assessment of heavy metals, pH, organic matter and organic carbon in roadside soils in Makurdi Metropolis, Benue State, Nigeria, *J. Environ. Prot.*, 4 (6), 618–628.
- [24] Kuriata-Potasznik, A., Szymczyk, S., Skwierawski, A., Glińska-Lewczuk, K., and Cymes, I., 2016, Heavy metal contamination in the surface layer of bottom sediments in a flow-through lake: A case study of Lake Symsar in Northern Poland, *Water*, 8 (8), 358.
- [25] Maher, W., Krikowa, F., Wruck, D., Louie, H., Nguyen, T., and Huang, W.Y., 2002, Determination of total phosphorus and nitrogen in turbid waters by oxidation with alkaline potassium peroxodisulfate and low pressure microwave digestion, autoclave heating or the use of closed vessels in a hot water bath: Comparison with Kjeldahl digestion, *Anal. Chim. Acta*, 463 (2), 283–293.
- [26] Song, Y., Hahn, H.H., and Hoffmann, E., 2002, Effects of solution conditions on the precipitation of phosphate for recovery a thermodynamic evaluation, *Chemosphere*, 48 (10), 1029–1034.
- [27] Kumar, M.M.S.Y., Galil, M.S.A., Suresha, M.S., Sathish, M.A., and Nagendrappa, G., 2007, A simple spectrophotometric determination of phosphate in sugarcane juices, water, and detergent samples, *E-J. Chem.*, 4 (4), 467–473.
- [28] Topçu, A., and Pulatsü, S., 2014, Phosphorus fractions and cycling in the sediment of a shallow eutrophic pond, *J. Agric. Sci.*, 20 (1), 63–70.
- [29] Sengupta, S., Nawaz, T., and Beaudry, J., 2015, Nitrogen and phosphorus recovery from wastewater, *Curr. Pollut. Rep.*, 1 (3), 155–166.
- [30] Cerozi, B.S., and Fitzsimmons, K., 2016, The effect of pH on phosphorus availability and speciation in an aquaponics nutrient solution, *Bioresour. Technol.*, 219, 778–781.
- [31] Lei, Y., Song, B., van der Weijden, R.D., Saakes, M., and Buisman, C.J.N., 2017, Electrochemical induced calcium phosphate precipitation: importance of local pH, *Environ. Sci. Technol.*, 51 (19), 11156–11164.
- [32] De Saedeleer, V., Cappuyns, V., De Cooman, W., and Swennen, R., 2010, Influence of major elements on heavy metal composition of river sediments, *Geol. Belg.*, 13 (3), 257–268.
- [33] Jiang, M., Zeng, G., Zhang, C., Ma, X., Chen, M., Zhang, J., Lu, L., Yu, Q., Hu, L., and Liu, L., 2012, Assessment of heavy metal contamination in the surrounding soils and surface sediments in Xiawangang River, Qingshuitang District, *PLoS One*, 8 (8), e71176.
- [34] Alamgir, M., Islam, M., Hossain, N., Kibria, M.G., and Rahman, M.M., 2015, Assessment of heavy metal contamination in urban soils of Chittagong City, Bangladesh, *Int. J. Plant Soil Sci.*, 7 (6), 362–372.
- [35] Zhang, C., Yu, Z., Zeng, G., Jiang, M., Yang, Z., Cui, F., Zhu, M., Shen, L., and Hu, L., 2014, Effects of sediment geochemical properties on heavy metal bioavailability, *Environ. Int.*, 73, 270–281.
- [36] Yao, Q., Wang, X., Jian, H., Chen, H., and Yu, Z., 2015, Characterization of the particles size fraction associated with heavy metals in suspended sediments of the Yellow River, *Int. J. Environ. Res. Public Health*, 12 (6), 6725–6744.

ORIGINALITY REPORT

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

9%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	sinta3.ristekdikti.go.id Internet Source	2%
2	sucra.repo.nii.ac.jp Internet Source	1%
3	journal.unnes.ac.id Internet Source	1%
4	Submitted to Griffith University Student Paper	1%
5	Submitted to Universitas Brawijaya Student Paper	1%
6	Submitted to Manchester Metropolitan University Student Paper	1%
7	gtg.webhost.uoradea.ro Internet Source	1%
8	Nigar Alkan, Ali Alkan, Ahmet Demirak, Moez Bahloul. "Metals/metalloid in Marine Sediments, Bioaccumulating in Macroalgae	<1%

and a Mussel", Soil and Sediment
Contamination: An International Journal, 2020

Publication

9

www.e3s-conferences.org

Internet Source

<1%

10

www.bio.uni-frankfurt.de

Internet Source

<1%

11

Muhammad Aqeel Ashraf, Mohd. Jamil Maah, Ismail Yusoff. "Morphology, Geology and Water Quality Assessment of Former Tin Mining Catchment", The Scientific World Journal, 2012

Publication

<1%

12

Submitted to Sunway Education Group

Student Paper

<1%

13

Upik Nurbaiti, Darminto, Triwikantoro, Mochamad Zainuri, Suminar Pratapa. "Synthesis and characterization of silica sand-derived nano-forsterite ceramics", Ceramics International, 2018

Publication

<1%

14

ee.hnu.cn

Internet Source

<1%

15

DAVID JOHNSON. "Use of estimated glomerular filtration rate to assess level of kidney function", Nephrology, 10/2005

Publication

<1%

16

Helena Doležalová Weissmannová, Jiří Pavlovský. "Indices of soil contamination by heavy metals – methodology of calculation for pollution assessment (minireview)", *Environmental Monitoring and Assessment*, 2017

Publication

<1%

17

Submitted to University of KwaZulu-Natal

Student Paper

<1%

18

R. Narendran, K. Kathiresan, R.S. Sathishkumar, K. Kayalvizhi, A. Sundaramanickam. "Bioremoval of toxic substances in synthetic wastewater using *Trichoderma pubescens* (NPK2), isolated from mangrove soil", *Biocatalysis and Agricultural Biotechnology*, 2019

Publication

<1%

19

Zhining Huang, Yankui Tang, Kaixuan Zhang, Yanhui Chen, Ying Wang, Lingmin Kong, Tianyu You, Zhanqi Gu. "Environmental risk assessment of manganese and its associated heavy metals in a stream impacted by manganese mining in South China", *Human and Ecological Risk Assessment: An International Journal*, 2016

Publication

<1%

20

eprints.utar.edu.my

Internet Source

<1%

21 isdesr.org Internet Source <1%

22 link.springer.com Internet Source <1%

23 Zhang, Chang, Zhi-gang Yu, Guang-ming Zeng, Min Jiang, Zhong-zhu Yang, Fang Cui, Meng-ying Zhu, Liu-qing Shen, and Liang Hu. "Effects of sediment geochemical properties on heavy metal bioavailability", Environment International, 2014. Publication <1%

24 de Wit, R.. "ROBUST: The ROle of BUffering capacities in STabilising coastal lagoon ecosystems", Continental Shelf Research, 200112 Publication <1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On