

Korespondensi Artikel dengan Judul
**Short Communication: The Pattern of Heavy Metals Distribution in Time
Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia**

Dari: "Prof. Dr.rer.nat. Nuryono, MS" <nuryono_mipa@ugm.ac.id>
Tanggal: 1 Mar 2018 6.57 PM
Subjek: [IJC] Submission Acknowledgement
Kepada: andri kurniawan <andri_pangkal@yahoo.co.id>

andri kurniawan:

Thank you for submitting the manuscript, "Short Communication: The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia" to Indonesian Journal of Chemistry. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL: <https://jurnal.ugm.ac.id/ijc/author/submission/33613>

Username: xxxxxxxxxxxxxxxxxxxx

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Prof. Dr.rer.nat. Nuryono, MS
Indonesian Journal of Chemistry

Nuryono Nuryono <nuryono_mipa@ugm.ac.id>

Kepada: Andri Kurniawan

Sel, 17 Jul 2018 jam 14.07

Dear Andri Kurniawan:

We have reached a decision regarding your submission to Indonesian Journal of Chemistry, "The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia".

Our decision is to: Accept Submission

In accordance to the Journal policy, you are required to immediately pay the publication fee of Rp 2,500,000 (USD 200) by transfer to the following bank account:

Bank name : PT. BANK NEGARA INDONESIA (PERSERO) TBK

SWIFT Code : BNINIDJAXXX

Account number : 0495984067

Account holder : Nurzanah Hidayanti

Bank Address : BNI UGM Branch, Sekip Utara Yogyakarta Indonesia 55281

Please send the evidence of remittance by fax or email to the editorial

office of Indonesian Journal of Chemistry (email: ijc@ugm.ac.id).

In few days after payment, you will receipt email for the further process, i.e. copy-editing, layout and proofreading.

Thank you for your valuable contribution to the journal.

Best regards,

Nuryono Nuryono
Laboratory of Inorganic Chemistry,
Department of Chemistry,
Universitas Gadjah Mada
Phone +628156800908
Fax +62274545188
nuryono_mipa@ugm.ac.id
Editor in Chief
Indonesian Journal of Chemistry
Indexed in SCOPUS since 2012

Nuryono Nuryono <nuryono_mipa@ugm.ac.id>

Kepada:Andri Kurniawan

Cc:Nuryono Nuryono

Sen, 6 Agu 2018 jam 14.09

Andri Kurniawan:

We have now copyedited your submission "The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia" for Indonesian Journal of Chemistry. To review the proposed changes and respond to Author Queries, please follow these steps:

1. Log into the journal using URL below with your username and password (use Forgot link if needed).
2. Click on the file at 1. Initial Copyedit File to download and open copyedited version.
3. Review the copyediting, making changes using Track Changes in Word, and answer queries.
4. Save file to desktop and upload it in 2. Author Copyedit.
5. Click the email icon under COMPLETE and send email to the editor.

Please also check the comments in the enclosed manuscript.

This is the last opportunity that you have to make substantial changes. You will be asked at a later stage to proofread the galleys, but at that point only minor typographical and layout errors can be corrected.

Manuscript URL: <https://jurnal.ugm.ac.id/ijc/author/submissionEditing/33613>

Username: andri_pangkal

If you are unable to undertake this work at this time or have any questions, please contact me. Thank you for your contribution to this journal.

Nuryono Nuryono
Editor in Chief
Indonesian Journal of Chemistry
Indexed in SCOPUS since 2012

adec kurnia <andri_pangkal@yahoo.co.id>

Kepada:Nuryono Nuryono

Rab, 3 Okt 2018 jam 00.35

Dear Proofreader

I have revised fig 3 of the article and i have attached it in this email and in the article file.
Thank you for your attentions.

Regard,

andri Kurniawan

Nuryono Nuryono <nuryono_mipa@ugm.ac.id>

Kepada:Andri Kurniawan

Kam, 4 Okt 2018 jam 09.41

Andri Kurniawan:

Nuryono Nuryono has added a comment to the submission, "The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia" in Indonesian Journal of Chemistry:

Those have been added in the paper

Editor in Chief

Indonesian Journal of Chemistry

Indexed in SCOPUS since 2012

Nuryono Nuryono <nuryono_mipa@ugm.ac.id>

Kepada:Andri Kurniawan

Kam, 4 Okt 2018 jam 09.44

Andri Kurniawan:

Thank you for proofreading the galleys for your manuscript, "The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia," in Indonesian Journal of Chemistry. We are looking forward to publishing your work shortly.

If you subscribe to our notification service, you will receive an email of the Table of Contents as soon as it is published. If you have any questions,

please contact me.

Nuryono Nuryono
Laboratory of Inorganic Chemistry,
Department of Chemistry,
Universitas Gadjah Mada

andri_pangkal <andri_pangkal@yahoo.co.id>

Kepada:Nuryono Nuryono

Kam, 4 Okt 2018 jam 20.11

Dear Editor in Chief

I would like to say thank you for your attentions.

I want to subscribe to journal' notification service for receiving an email of the Table of Contents.

Thank you.

Andri

Nuryono Nuryono <nuryono_mipa@ugm.ac.id>

Kepada:Andri Kurniawan

Rab, 21 Nov 2018 jam 11.30

Readers:

Indonesian Journal of Chemistry has just published its latest issue at
<https://jurnal.ugm.ac.id/ijc>.

Thanks for the continuing interest in our work and we invite you to submit your research work.

Best regards,

Nuryono Nuryono
Laboratory of Inorganic Chemistry,
Department of Chemistry,
Universitas Gadjah Mada

Short Communication:
The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia

Andri Kurniawan^{a,b,*}, Oedjijono^b, Tamad^c, and Uyi Sulaeman^d

^a *Department of Aquaculture, University of Bangka Belitung
Jl. Balunijuk, Merawang, Bangka Belitung Archipelago Province*

^b *Faculty of Biology, University of Jenderal Soedirman
Jl. dr. Suparno, Grendeng, Purwokerto, Central Java*

^c *Departement of Agriculture, University of Jenderal Soedirman
Jl. dr. Suparno, Grendeng, Purwokerto, Central Java*

^d *Department of Chemistry, University of Jenderal Soedirman
Jl. dr. Suparno, Grendeng, Purwokerto, Central Java*

* Corresponding author, tel/fax : +6281351714747, email: andri-kurniawan@ubb.ac.id

ABSTRACT

The heavy metals distribution of ex-tin mining ponds were investigated. The time chronosequence was determined at the pond of age < 1 year (Station A), the pond of age 5-10 years (Station B), and the pond of age > 15 years (Station C). The results showed sixteen heavy metals of As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V could be detected in the ponds. The As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn in Station C showed higher concentration compared to the Station A and Station B. The Cr, Fe, Hf, Sn, Ta, Te, and V in Station A and Station B showed higher concentration compared to the Station C. The positive, negative, and dynamic correlation pattern could be found in distribution of heavy metal to time chronosequence. The Ta and V showed a positive correlation because their concentrations decrease, whereas the As, Cu, Ga, Mn, and Zn showed a negative correlation because their concentrations increase along in time chronosequence. The dynamic correlation could be found that the concentrations of Co, Ni, Pb, Sn, and Th decrease from Stasion A to Stasion B and then increase in Stasion C, whereas the concentrations of Cr, Fe, Hf, and Te increase from Station A to Stasion B and then decrease in Stasion C.

Keywords: *distribution pattern, heavy metals, chronosequence, ex-tin mining ponds*

ABSTRAK

Distribusi logam berat dari danau pascatambang timah telah dikaji. Kronosekuens waktu danau ditetapkan pada umur < 1 tahun (Stasiun A), umur 5-10 tahun (Stasiun B), dan umur > 15 tahun (Station C). Hasil menunjukkan 16 logam berat, yaitu As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, dan V terdeteksi di danau tersebut. Logam As, Co, Cu, Ga, Mn, Ni, Pb, Th, dan Zn di Stasiun C memiliki konsentrasi lebih tinggi dibandingkan Stasiun A dan Stasiun B. Logam Cr, Fe, Hf, Sn, Ta, Te, dan V di Stasiun A dan Stasiun B memiliki konsentrasi lebih tinggi dibandingkan Stasiun C. Pola korelasi positif, negatif, dan dinamis diperoleh dari distribusi logam berat terhadap kronosekuens waktu. Logam Ta dan V menunjukkan korelasi positif karena konsentrasinya meningkat, sedangkan As, Cu, Ga, Mn, dan Zn menunjukkan korelasi negatif karena konsentrasinya menurun seiring kronosekuens waktu. Korelasi dinamis terlihat pada

konsentrasi Co, Ni, Pb, Sn, dan Th yang menurun dari Stasiun A ke Stasiun B dan kemudian meningkat di Stasiun C, serta konsentrasi Cr, Fe, Hf, and Te meningkat dari Stasiun A ke Stasiun B dan kemudian meningkat di Stasiun C.

Kata kunci: pola distribusi, logam berat, kronosekuens, danau pascatambang timah

INTRODUCTION

Tin mining activity is an anthropogenic activity that has contributed to the ecological problems. It has been known as the major sources of heavy metal contamination, such as heavy metals in ex-tin mining ponds. Ex-tin mining ponds as one of water resource were contaminated by heavy metals that can be accumulated in food chains and secondary activities of the water [1, 2-3]. Some of the heavy metals such as Pb, Zn, Mn, Fe, Cr, Cu, Ni, Sn, As, and Cd were detected in site of ex-tin mining [4-5].

The each of waters had their pattern of physical and chemical characteristics that determined by climatic, geomorphological, and geochemical conditions [6]. These conditions were influenced by ecological succession in time chronosequence [7] and the ecological changes of ex-tin mining ponds have occurred for a long time [8]. The implication of ecological changes as long as time chronosequence was interactions and reactions of chemical complexation between organic compounds, environmental factors, and inorganic compounds, include a presence of heavy metals [9].

The researches of heavy metals in ex-tin mining ecosystems have been done, however, not for correspondence of heavy metals distributions and ex-tin mining ponds in time chronosequence. While an understanding this correspondence was crucial to predict ecosystems conditions and to detect environment change. The understanding of heavy metals distribution in time chronosequence of ex-tin mining ponds becomes an important part of aquatic ecosystem management as water sources for secondary activities and habitable for organisms..

EXPERIMENTAL SECTION

Materials

The study areas were located in Bangka Regency, Bangka Belitung Archipelago Province of Indonesia. The considered study sites cover three ponds of ex-tin mining with chronosequence ranging in time. The study areas were encoded as Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years).

The coordinate of Station A were 01°59' S in points 36,0"; 36,2"; 36,4"; 36,5"; 36,6" and 106°06' E in points 36,5"; 36,9"; 37,3"; 37,4"; 37,5". The coordinate of Station B were 01°59' S in points 41,3"; 41,4"; 41,5"; 42,4"; 42,5" and 106°06' E in points 39,2"; 39,5"; 41,4"; 42,7"; 43,1". The coordinate of Station C were 01°55' S in points 40,9"; 58,9"; 59,1"; 59,2"; 59,5" and 106°06' E in points 19,5"; 19,7"; 19,9"; 22,4"; 29,2".

In the each of research stations points, water sampling was done < 4m in depth (station code A.1; B.1; and C.1) and composite sampling were done to water and sediment > 4m in depth (station code A.2; B.2; and C.2).

Instrumentation

The water samples and composite samples were analyzed pH value by pH meter (PH-009(I)-A) Sun Care. The speciation and concentration of heavy metals analyzed by X-Ray Fluorescence (Rigaku) with three light spreader metals of copper (Cu), molybdenum (Mo), and aluminum (Al). Also, the pH and moisture of soil around the ponds were measured by pH-moisture meter. The correspondence heavy metals distribution in time chronosequence of ex-tin mining water was analyzed by PAST (*Palaentological Statistics*) 3.

RESULTS AND DISCUSSION

pH Value of Water and Soil

Tin mining activity had generated the ex-tin mining ponds as one of the water resources. However, water quality in the ponds had serious ecological problem. The ex-mining activity produced acid mine drainage (AMD) that can cause the long-term impairment of water quality and biodiversity. The pH value of ex-tin mining water and soil pH value indicated that the acid could be found in Station A (pond in age < 1 year) and Station B (pond in age 5-10 years) and the neutral value could be found in Station C (pond in age > 15 years). In the each of research stations points, water samples < 4m in depth indicated more acid than the composite samples (water and sediment > 4m in depth) (Figure 1).




Figure 1. Soil and water pH value in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The acid mine drainage (AMD) was produced when sulfide-bearing material is exposed to oxygen and water [10]. In acid mine drainage, presence and mobility of elements such as Al, As,

Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Zn, and also ionic activity and sulphide minerals oxidation caused pH value decreasing [11-12]. Furthermore, low pH condition implicated in oxidation state, toxicity, and mobility of elements [9].

In long time chronosequence more than 15 years in the Station C caused changes water quality of ex-tin mining ponds were more stable than Station A and Station B. The indicator for it was pH value increased to neutral conditions (7,0-7,4) and other parameters that indicated water quality was more habitable for organisms' life such as fishes. The pH value and other parameters changes, indirectly, caused ecological productivity increasing. The chronosequence in long time caused changes in nutrient availability and resource supply for ecosystem productivity [13].

The pH value was found to play the most important role in the determination of metal speciation, solubility from mineral surfaces, movement, and bioavailability of metals. Furthermore, acid mine drainage had a major contribution in heavy metal contamination [14]. There was a negative correlation between soil pH and distribution heavy metal where pH value decreasing caused increasing in heavy metal desorption [15].

Pattern of Heavy Metals Distribution

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C) (Table 2).

Table 2. Distribution of heavy metals in time chronosequence in water of ex-tin mining ponds

No	Heavy Metals	Heavy Metals Concentration (ppm)					
		In Station A		In Station B		In Station C	
		Station A.1	Station A.2	Station B.1	Station B.2	Station C.1	Station C.2
1	As	Nd	2,155	nd	3,215	2,4	1,095
2	Ga	Nd	4,275	nd	4,47	9,76	nd
3	V	Nd	0,84	nd	nd	nd	0,67
4	Te	Nd	nd	102	nd	nd	5,8
5	Cr	Nd	4,735	nd	4,945	Nd	0,67
6	Zn	Nd	6,05	nd	8,1	8,17	3,905
7	Sn	47	82,8	50,3	51,4	61,8	57,15
8	Mn	Nd	13,05	nd	13,6	30,8	14,85
9	Fe	21,4	1481,3	148	2344	1590	849,05
10	Co	Nd	5,05	nd	nd	6,64	nd
11	Ni	Nd	4,09	nd	2,875	3,48	1,905
12	Cu	3,08	6,875	3,19	7,775	8,94	3,81
13	Pb	Nd	6,7	nd	6	8,21	4,62
14	Th	Nd	4,605	nd	4,38	13,5	7,05
15	Hf	6,71	3,21	8,21	3,32	nd	9,38
16	Ta	4,94	2,615	4,16	2,995	nd	2,805

*) *nd (not detected). Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).*

There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. The average of As concentration were 1,7475 ppm (Station C) > 1,6075 ppm (Station B) > 1,0775 ppm (Station A). The concentration of Co were 3,32 ppm (Station C) > 2,525 ppm (Station A) > not detected (Station B). The concentration of Cu were 6,375 ppm (Station C) > 5,4825 ppm (Station B) > 4,9775 ppm (Station A). The concentration of Ga were 4,88 ppm (Station C) > 2,235 ppm (Station B) > 2,1375 ppm (Station A). The concentration of Mn were 22,825 ppm (Station C) > 6,8 ppm (Station B) > 6,525 ppm (Station A). The concentration of Ni were 2,6925 ppm (Station C) > 2,045 ppm (Station A) > 1,4375 ppm (Station B). The concentration of Pb were 6,415 ppm (Station C) > 3,35 ppm (Station A) > 3,0 ppm (Station B). The concentration of Th were 10,275 ppm (Station C) > 2,3025 ppm (Station A) > 2,19 ppm (Station B). The concentration of Zn were 6,0375 ppm (Station C) > 4,05 ppm (Station B) > 3,025 ppm (Station A).

In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V. The average of Cr concentration were 2,4725 ppm (Station B) > 2,3675 ppm (Station A) > 0,335 ppm (Station C). The concentration of Fe were 1246 ppm (Station B) > 1219,53 ppm (Station C) > 751,35 ppm (Station A). The concentration of Hf were 5,765 ppm (Station B) > 4,96 ppm (Station A) > 4,69 ppm (Station C). The concentration of Sn were 64,9 ppm (Station A) > 59,475 ppm (Station C) > 50,85 ppm (Station B). The concentration of Ta were 3,7775 ppm (Station A) > 3,5775 ppm (Station B) > 1,4025 ppm (Station C). The concentration of Te were 51 ppm (Station B) > 2,9 ppm (Station C) > not detected (Station A). The concentration of V were 0,42 ppm (Station A) > 0,335 ppm (Station C) > not detected (Station B).

Their concentration of heavy metals in time chronosequence in ex-tin mining ponds showed three pattern distribution that a positive correlation, a negative correlation, and a dynamic correlation. The positive correlation indicated that concentration of heavy metals decreased along chronosequence in time in ex-tin mining ponds. The negative correlation indicated that concentration of heavy metals increased along chronosequence in time in ex-tin mining ponds. The dynamic correlation indicated that concentration of heavy metals on decreased or increased along chronosequence in time in ex-tin mining ponds.

Elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. Elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increasing along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5-10 years and than increased concentration in ponds with age > 15 years. Elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5-10 years and than decreased concentration in ponds with age > 15 years (Figure 2).




Figure 2. Distribution of heavy metals in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. Station A(pond in age < 1 year), Station B(pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The chronosequence in long time caused changes in heavy metals concentrations were decreased as the primary succession progression [16]. The low pH value implicated to dissolved oxygen (DO) [17] and low DO in acid mine drainage implicated to biological oxygen demand (BOD) by microorganisms for their growth and other ecological factor changes, such as nitrogen and phosphate [7, 18-19]. Nitrogen (ammonia, nitrit, and nitrate) and phosphate were important for sustenance of various life forms, primary succession, and led to changes in the biogeochemical cycles of nutrients in ponds [20, 21-22], indirectly implicated to dissolved and suspended solid [23-24] and ecosystem eutrophic levels [25, 26-31]. Therefore, pH value changes, availability of elements included heavy metals, environmental characteristics, and eutrophication had a positive inter-relationships in their correspondence.

These conditions, however, had a negative correlation to heavy metals residue decreasing in Station C. The heavy metals in Station C were dominated As, Zn, Mn, Co, Ni, Cu, Pb, and Th were higher than Station A and Station B, although these metals and include Ga, Cr, Sn, Fe, Hf, and Ta were detected in all of the research station. The results indicated that chronosequence in time of ponds caused ecological changes and had positive correlations, but not for distribution of heavy metals.

Correspondence Analysis of Heavy Metals Distribution

The correspondence analysis also showed that there was not a positive correlation between in time chronosequence in water of ex-tin mining ponds and heavy metals distribution (Figure 3). The ecological changes in chronosequence succession of ex-tin mining ponds had not significantly affect on speciations and concentrations of heavy metals. Thus, there was the other factor that had affected the distribution of heavy metals.




Figure 3. Correspondence analysis between distribution of heavy metals and in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The soil and sediment of water contributed to the distribution of heavy metals [32]. In general, the mobility and availability of heavy metals were controlled by adsorption and desorption characteristics of soils, including cation exchange capacity and the contents of clay minerals [15]. The clay and mud were found to be dominant in the sediment of Station A and Station B, however, sand was found to be dominant in Station C.

The clay and mud had a positive correlation and sand had a negative correlation to a presence, affinity, and minerals structure [33-34]. The clay and mud had a larger surface area, higher cation exchange capability (CEC), and particles of clay in sediments were often closely associated with organic matter and also exist in the form of organic-mineral complexes. These particles adsorbed the active metals from the water phases, accumulated, and carried them to the bottom sediments diagenetically [35-36]. Therefore, distribution of heavy metals in Station A and Station B were lower than Station C.

CONCLUSION

In time chronosequence of ex-tin mining ponds had affected ecological dynamic, such as pH value change and heavy metals distribution. The pH value of ex-tin mining ponds moved from acid to neutral condition. However it had been taken for a long time of the chronosequence. The consequence of low pH value can implicate to microorganisms' growth, nutrients, and environment factors in primary succession. Further, it implied interaction to heavy metals. In long time chronosequence should decrease heavy metals residue in the ex-tin mining ponds.

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C). There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V.

There were three pattern of heavy metals distribution in time chronosequence in ex-tin mining ponds that a positive correlation, a negative correlation, and a dynamic correlation. The chronosequence in time had affects on heavy metals distribution. Elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. Elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increasing along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5-10 years and than increased concentration in ponds with age > 15 years. Elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5-10 years and than decreased concentration in ponds with age > 15 years.

The chronosequence in time can be an factor that contributed to heavy metals distribution. However, the sediment factor had a significant effect to distribution of heavy metals in the ex-tin mining ponds, where clay and mud were more effective than sand in accumulation, adsorption, and interaction with heavy metals and carried them diagenetically to bottom sediments.

ACKNOWLEDGEMENT

I would like to thank Ministry of Research, Technology and Higher Education of the Republic of Indonesia for scholarship of Pendidikan Pascasarjana Dalam Negeri (BPPDN) and research grand of doctoral disertation (PDD).

REFERENCES

1. De Lima e Silva, AA., Ribeiro de Carvalho, MA., L de Souza, SA., Dias, PMT., Da Silva Filho, RG., De Meirelles Saramago, CS., De Melo Bento, CA., Hofer, E., 2012, Heavy metal

- tolerance (Cr, Ag and Hg) in bacteria isolated from sewage, *Brazilian Journal of Microbiology*, 43(4), 1620-1631.
2. Guan, Y., Shao, C., Ju, M., 2014, Heavy metal contamination assessment and partition for industrial and mining gathering areas, *International Journal of Environmental Research and Public Health*, 11(7), 7286-7303.
 3. Kurniawan, A., 2017, Chronosequence effect of post tin mining ponds to metals residu and microecosystem change, *Omni-Akuatika*, 13(1), 60-65.
 4. Ashraf, MA., Maah, MJ., Yusoff, I., 2012, Speciation of heavy metals in the sediments of former tin mining catchment, *Iranian Journal of Science and Technology*, 36(A2), 163-180.
 5. Daniel, VN., Chudusu, ES., Chup, JA., Pius, ND., 2014, Variations of heavy metals in agricultural soils irrigated with tin water in Heipang District of Barkin Ladi, Plateau State, Nigeria, *International Journal of Science and Technology*, 3(5), 255-263.
 6. Ashraf, MA., Maah, MJ., Yusoff, I., 2012. Morphology, geology and water quality assessment of former tin mining catchment, *The Scientific World Journal*, 2012, 1-15.
 7. Urbanov´a, M., Kopecký, J., Val´as´kov´a, V., S´agov´a-Marec´kov´a, M., Elhottov´a, D., Kyselkov´a, M., Moe¨nne-Loccoz, Y., Baldrian, P., 2011, Development of bacterial community during spontaneous succession on spoil heaps after brown coal mining, *FEMS Microbiology Ecology*, 78(1), 59-69.
 8. Kurniawan, A., 2016, Microorganism communities response of ecological changes in post tin mining ponds, *Journal of Microbiology and Virology*, 6(1), 17-26.
 9. Violante, A., Cozzolino, V., Perelomov, L., Caporale, AG., Pigna, M., 2010, Mobility and bioavailability of heavy metals and metalloids in soil environments, *Journal of Soil Science and Plant Nutrition*, 10(3), 268-292.
 10. Akcil, A., and Koldas, S., 2006, Acid mine drainage (AMD): causes, treatment and case studies, *Journal of Cleaner Production*, 14(12-13), 1139-1145.
 11. Tan, G., Shu, W., Hallberg, KB., Li, F., Lan, C., Huang, L., 2007, Cultivation-dependent and cultivation-independent characterization of the microbial community inacid mine drainage associated with acidic Pb/Zn mine tailings at Lechang, Guangdong, China, *FEMS Microbiology Ecology*, 59(1), 118-126.
 12. Campaner, VP., Luiz-Silva, W., Machado, W., 2014, Geochemistry of acid mine drainage from a coal mining area and processes controlling metal attenuation in stream waters, Southern Brazil, *Anais da Academia Brasileira de Cincias*, 86(2), 539-554.
 13. Bardgett, RD., Bowman, WD., Kaufmann, R., Schmidt, SK., 2005, A temporal approach to linking aboveground and belowground ecology, *TRENDS in Ecology and Evolution*, 20(11), 634-641.
 14. Ahmad, AK., and Al-Mahaqeri, SA., 2015, Human health risk assessment of heavy metals in fish species collected from catchments of former tin mining, *International Journal of Research Studies in Science, Engineering and Technology*, 2(4), 9-21.
 15. Zeng, F., Ali, S., Zhang, H., Ouyang, Y., Qiu, B., Wu, F., Zhang, G., 2011, The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants, *Environmental Pollution*, 159(1), 84-91.
 16. Huang, L., Tang, F., Song, Y., Wan, C., Wang, S., Liu, W., Shu, W., 2010. Biodiversity, abundance, and activity of nitrogen-fixing bacteria during primary succession on a copper mine tailings, *FEMS Microbiology Ecology*, 78(3), 439-450.
 17. Abdel-Raouf, N., Al-Homaidan, AA., Ibraheem, IBM., 2012, Microalgae and wastewater treatment, *Saudi Journal of Biological Sciences* 19(3), 257-275.
 18. Tscherko, D., Hammesfahr, U., Zeltner, G., Kandeler, E., Bocker, R.. 2005, Plant succession and rhizosphere microbial communities in a recently deglaciated alpine terrain. *Basic and Applied Ecology*, 6(4), 367-383.
 19. Fierer, N., Nemergut, D., Knight, R., Craine, JM., 2010, Changes through time: integrating microorganisms into the study of succession, *Research in Microbiology*, 161(8), 635-642.
 20. Quirós, R., 2003, The relationship between nitrate and ammonia concentrations in the pelagic zone of lakes. *Limnetica*, 22(1-2), 37-50.

21. Michalski, R., and Kurzyca, I. 2006, Determination of nitrogen species (nitrate, nitrite and ammonia ions) in environmental samples by ion chromatography. *Polish Journal of Environmental Studies*, 15(1), 5-18.
22. Mihale, MJ., 2015, Nitrogen and phosphorus dynamics in the waters of the Great Ruaha River, Tanzania, *Journal of Water Resources and Ocean Science*, 4(5), 59-71.
23. Akan, JC., Audu, SI., Mohammed, Z., Ogugbuaja, VO., 2013, Assessment of heavy metals, pH, organic matter and organic carbon in roadside soils in Makurdi Metropolis, Benue State, Nigeria, *Journal of Environmental Protection*, 4(6), 618-628.
24. Kuriata-Potasznik, A., Szymczyk, S., Skwierawski, A., Glińska-Lewczuk, K., Cymes, I., 2016, Heavy metal contamination in the surface layer of bottom sediments in a flow-through lake: a case study of Lake Symsar in Northern Poland, *Water*, 8(8), 1-15.
25. Maher, W., Krikowa, F., Wruck, D., Louie, H., Nguyen, T., Huang, WY., 2002, Determination of total phosphorus and nitrogen in turbid waters by oxidation with alkaline potassium peroxodisulfate and low pressure microwave digestion, autoclave heating or the use of closed vessels in a hot water bath: comparison with kjeldahl digestion, *Analytica Chimica Acta*, 463(2), 283-293.
26. Song, Y., Hahn, HH., Hoffmann, E., 2002, Effects of solution conditions on the precipitation of phosphate for recovery a thermodynamic evaluation, *Chemosphere*, 48(10), 1029-1034.
27. Mahadevaiah., Kumar, MSY., Galil, MSA., Suresha, MS., Sathish, MA., Nagendrappa, G., 2007, A simple spectrophotometric determination of phosphate in sugarcane juices, water, and detergent samples, *E-Journal of Chemistry*, 4(4): 467-473.
28. Topcu, A., and Pulatsu, S., 2014, Phosphorus fractions and cycling in the sediment of a shallow eutrophic pond. *Journal of Agricultural Sciences*, 2014(20), 63-70.
29. Sengupta, S., Nawaz, T., Beaudry, J., 2015, Nitrogen and phosphorus recovery from wastewater. *Current Pollution Reports*, 1(3), 155-166.
30. Da Silva Cerozi, B., and Fitzsimmons, K., 2016, The effect of pH on phosphorus availability and speciation in an aquaponics nutrient solution. *Bioresource Technology*, 2016(219), 778-781.
31. Lei, Y., Song, B., Van Der Weijden, RD., Saakes, M., Buisman, CJN., 2017, Electrochemical induced calcium phosphate precipitation: importance of local pH. *Environmental Science & Technology*, 51(19): 11156-11164.
32. De Saedeleer, V., Cappuyns, V., De Cooman, W., Swennen, R., 2010, Influence of major elements on heavy metal composition of river sediments. *Geologica Belgica*, 13(3), 257-268.
33. Jiang, M., Zeng, G., Zhang, C., Ma, X., Chen, M., Zhang, J., Lu, L., Yu, Q., Hu, L., Liu, L., 2012, Assessment of heavy metal contamination in the surrounding soils and surface sediments in Xiawangang River, Qingshuitang District. *PLoS ONE*, 8(8), e71176.
34. Alamgir, M., Islam, M., Hossain, N., Kibria, MG., Rahman, MM., 2015, Assessment of heavy metal contamination in urban soils of Chittagong City, Bangladesh, *International Journal of Plant & Soil Science*, 7(6), 362-372.
35. Zhang, C., Yu, Z., Zeng, G., Jiang, M., Yang, Z., Cui, F., Zhu, M., Shen, L., Hu, L., 2014, Effects of sediment geochemical properties on heavy metal bioavailability. *Environment International*, 2014(73), 270-281.
36. Yao, Q., Wang, X., Jian, H., Chen, H., Yu, Z., 2015, Characterization of the particles size fraction associated with heavy metals in suspended sediments of the Yello River. *International Journal of Environment Research and Public Health*, 12(6), 6725-6744.

The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia

Complete with name of all authors with institution address and corresponding authors

ABSTRACT

The heavy metals distribution of ex-tin mining ponds were investigated. The time chronosequence was determined at the pond of age < 1 year (Station A), the pond of age 5-10 years (Station B), and the pond of age > 15 years (Station C). The results showed sixteen heavy metals of As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V could be detected in the ponds. The metals such as As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn in Station C showed higher concentration compared to the Station A and Station B. The metals such as Cr, Fe, Hf, Sn, Ta, Te, and V in Station A and Station B showed higher concentration compared to the Station C. The positive, negative, and dynamic correlation pattern could be found in distribution of heavy metal to time chronosequence. The concentration of Ta and V showed a positive correlation because their concentration decrease, whereas concentration of As, Cu, Ga, Mn, and Zn showed a negative correlation because their concentration increase along in time chronosequence. The dynamic correlation could be found that concentration of Co, Ni, Pb, Sn, and Th decrease from Station A to Station B and then increase in Station C, whereas concentration of Cr, Fe, Hf, and Te increase from Station A to Station B and then decrease in Station C.

Keywords: distribution pattern, heavy metals, chronosequence, ex-tin mining ponds

ABSTRAK

Distribusi logam berat dari danau pascatambang timah telah dikaji. Kronosekuensi waktu danau ditetapkan pada umur < 1 tahun (Stasiun A), umur 5-10 tahun (Stasiun B), dan umur > 15 tahun (Stasiun C). Hasil menunjukkan 16 logam berat yaitu As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, dan V terdeteksi di danau tersebut. Logam As, Co, Cu, Ga, Mn, Ni, Pb, Th, dan Zn di Stasiun C memiliki konsentrasi lebih tinggi dibandingkan Stasiun A dan Stasiun B. Logam Cr, Fe, Hf, Sn, Ta, Te, dan V di Stasiun A dan Stasiun B memiliki konsentrasi lebih tinggi dibandingkan Stasiun C. Pola korelasi positif, negatif, dan dinamis diperoleh dari distribusi logam berat terhadap kronosekuensi waktu. Konsentrasi Ta dan V menunjukkan korelasi positif karena konsentrasinya meningkat, sedangkan konsentrasi As, Cu, Ga, Mn, dan Zn menunjukkan korelasi negatif karena konsentrasinya menurun seiring kronosekuensi waktu. Korelasi dinamis terlihat pada konsentrasi Co, Ni, Pb, Sn, dan Th yang menurun dari Stasiun A ke Stasiun B dan kemudian meningkat di Stasiun C, serta konsentrasi Cr, Fe, Hf, dan Te meningkat dari Stasiun A ke Stasiun B dan kemudian menurun di Stasiun C.

Kata kunci: pola distribusi, logam berat, kronosekuensi, danau pascatambang timah

INTRODUCTION

Tin mining activity is an anthropogenic activity that has contributed to the ecological problems. It has been known as the major sources of heavy metal contamination such as heavy metals in ex-tin mining ponds. Ex-tin mining ponds as one of water resource were contaminated by heavy metals that can be accumulated in food chains and secondary activities of the water [1, 2-3]. Some of the

heavy metals such as Pb, Zn, Mn, Fe, Cr, Cu, Ni, Sn, As, and Cd were detected in site of ex-tin mining [4-5].

The each of waters had their pattern of physical and chemical characteristics that determined by climatic, geomorphological, and geochemical conditions [6]. These conditions were influenced by ecological succession in time chronosequence [7] and the ecological changes of ex-tin mining ponds have occurred for a long time [8]. The implication of ecological changes as long as time chronosequence was interactions and reactions of chemical complexion between organic compounds, environmental factors, and inorganic compounds, include presence of heavy metals [9].

The researches of heavy metals in ex-tin mining ecosystems have been done, however, not for correspondence of heavy metals distributions and ex-tin mining ponds in time chronosequence. While an understanding this correspondence was crucial to predict ecosystems conditions and to detect environment change. The understanding of heavy metals distribution in time chronosequence of ex-tin mining ponds becomes an important part of aquatic ecosystem management as water sources for secondary activities and habitable for organisms.

EXPERIMENTAL SECTION

Materials

The study areas were located in Bangka Regency, Bangka Belitung Archipelago Province of Indonesia. The considered study sites cover three ponds of ex-tin mining with chronosequence ranging in time. The study areas were encoded as Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years).

The coordinate of Station A were 01°59' S in points 36,0"; 36,2"; 36,4"; 36,5"; 36,6" and 106°06' E in points 36,5"; 36,9"; 37,3"; 37,4"; 37,5". The coordinate of Station B were 01°59' S in points 41,3"; 41,4"; 41,5"; 42,4"; 42,5" and 106°06' E in points 39,2"; 39,5"; 41,4"; 42,7"; 43,1". The coordinate of Station C were 01°55' S in points 40,9"; 58,9"; 59,1"; 59,2"; 59,5" and 106°06' E in points 19,5"; 19,7"; 19,9"; 22,4"; 29,2".

In the each of research stations points, water sampling was done < 4m in depth (station code A.1; B.1; and C.1) and composite sampling were done to water and sediment > 4m in depth (station code A.2; B.2; and C.2).

Instrumentation

The water samples and composite samples were analyzed pH value by pH meter (PH-009(I)-A) Sun Care. The speciation and concentration of heavy metals analyzed by X-Ray Fluorescence (Rigaku) with three light spreader metals of copper (Cu), molybdenum (Mo), and aluminum (Al). Also, the pH and moisture of soil around the ponds were measured by pH-moisture meter. The

correspondence between heavy metals distribution and the time chronosequence of ex-tin mining ponds was analyzed by PAST (*Palaentological Statistics*) 3.

RESULTS AND DISCUSSION

pH Value of Water and Soil

Tin mining activity had generated the ex-tin mining ponds as one of the water resources. However, water quality in the ponds had serious ecological problem. The ex-mining activity produced acid mine drainage (AMD) that can caused the long-term impairment of water quality and biodiversity. The pH value of ex-tin mining water and soil pH value indicated that the acid could be found in Station A (pond in age < 1 year) and Station B (pond in age 5-10 years) and the neutral value could be found in Station C (pond in age > 15 years). In the each of research stations points, water samples < 4m in depth indicated more acid than the composite samples (water and sediment > 4m in depth) (Figure 1).




Figure 1. Soil and water pH value in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B(pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The acid mine drainage (AMD) was produced when sulfide-bearing material is exposed to oxygen and water [10]. In acid mine drainage, presence and mobility of elements such as Al, As, Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Zn, and also ionic activity and sulphide minerals oxidation caused pH value decreasing [11-12]. Furthermore, the low pH condition implicated in oxidation state, toxicity, and mobility of elements [9].

In long time chronosequence more than 15 years in the Station C caused changes water quality of ex-tin mining ponds were more stable than Station A and Station B. The indicator for it was pH value increased to neutral conditions (7,0-7,4) and other parameters that indicated water quality was more habitable for organisms' life such as fishes. The pH value and other parameters changes, indirectly, caused ecological productivity increasing. The chronosequence in long time caused changes in nutrient availability and resource supply for ecosystem productivity [13].

The pH value was found to play the most important role in the determination of metal speciation, solubility from mineral surfaces, movement, and bioavailability of metals. Furthermore, acid mine drainage had a major contribution in heavy metal contamination [14]. There was a negative correlation between soil pH and distribution heavy metal where pH value decreasing caused increasing in heavy metal desorption [15].

The Pattern of Heavy Metals Distribution

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C) (Table 1).

Table 1. Distribution of heavy metals in time chronosequence in water of ex-tin mining ponds

No	Heavy Metals	Heavy Metals Concentration (ppm)					
		In Station A		In Station B		In Station C	
		Station A.1	Station A.2	Station B.1	Station B.2	Station C.1	Station C.2
1	As	nd	2,155	nd	3,215	2,4	1,095
2	Ga	nd	4,275	nd	4,47	9,76	nd
3	V	nd	0,84	nd	nd	nd	0,67
4	Te	nd	nd	102	nd	nd	5,8
5	Cr	nd	4,735	nd	4,945	nd	0,67
6	Zn	nd	6,05	nd	8,1	8,17	3,905
7	Sn	47	82,8	50,3	51,4	61,8	57,15
8	Mn	nd	13,05	nd	13,6	30,8	14,85
9	Fe	21,4	1481,3	148	2344	1590	849,05
10	Co	nd	5,05	nd	nd	6,64	nd
11	Ni	nd	4,09	nd	2,875	3,48	1,905
12	Cu	3,08	6,875	3,19	7,775	8,94	3,81
13	Pb	nd	6,7	nd	6	8,21	4,62
14	Th	nd	4,605	nd	4,38	13,5	7,05
15	Hf	6,71	3,21	8,21	3,32	nd	9,38
16	Ta	4,94	2,615	4,16	2,995	nd	2,805

*) *nd (not detected). The Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).*

There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. The average of As concentration were 1,7475 ppm (Station C) > 1,6075 ppm (Station B) > 1,0775 ppm (Station A). The concentration of Co were 3,32 ppm (Station C) > 2,525 ppm (Station A) > not detected (Station B). The concentration of Cu were 6,375 ppm (Station C) > 5,4825 ppm (Station B) > 4,9775 ppm (Station A). The concentration of Ga were 4,88 ppm (Station C) > 2,235 ppm (Station B) > 2,1375 ppm (Station A). The concentration of Mn were 22,825 ppm (Station C) > 6,8 ppm (Station B) > 6,525 ppm (Station A). The concentration of Ni were 2,6925 ppm (Station C) > 2,045 ppm (Station A) > 1,4375 ppm (Station B). The concentration of Pb were 6,415 ppm (Station C) > 3,35 ppm (Station A) > 3,0 ppm (Station B). The concentration of Th were 10,275 ppm (Station C) > 2,3025 ppm (Station A) > 2,19

ppm (Station B). The concentration of Zn were 6,0375 ppm (Station C) > 4,05 ppm (Station B) > 3,025 ppm (Station A).

In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V. The average of Cr concentration were 2,4725 ppm (Station B) > 2,3675 ppm (Station A) > 0,335 ppm (Station C). The concentration of Fe were 1246 ppm (Station B) > 1219,53 ppm (Station C) > 751,35 ppm (Station A). The concentration of Hf were 5,765 ppm (Station B) > 4,96 ppm (Station A) > 4,69 ppm (Station C). The concentration of Sn were 64,9 ppm (Station A) > 59,475 ppm (Station C) > 50,85 ppm (Station B). The concentration of Ta were 3,7775 ppm (Station A) > 3,5775 ppm (Station B) > 1,4025 ppm (Station C). The concentration of Te were 51 ppm (Station B) > 2,9 ppm (Station C) > not detected (Station A). The concentration of V were 0,42 ppm (Station A) > 0,335 ppm (Station C) > not detected (Station B).

Their concentration of heavy metals in time chronosequence in ex-tin mining ponds showed three patterns distribution that a positive correlation, a negative correlation, and a dynamic correlation. The positive correlation indicated that concentration of heavy metals decreased along chronosequence in time in ex-tin mining ponds. The negative correlation indicated that concentration of heavy metals increased along chronosequence in time in ex-tin mining ponds. The dynamic correlation indicated that concentration of heavy metals on decreased or increased along chronosequence in time in ex-tin mining ponds.

The elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. The elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increasing along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5-10 years and then increased concentration in ponds with age > 15 years. The elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5-10 years and then decreased concentration in ponds with age > 15 years (Figure 2).




Figure 2. Distribution of heavy metals in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The chronosequence in long time caused changes in heavy metals concentration were decreased as the primary succession progression [16]. The low pH value implicated to dissolved oxygen (DO) [17] and low DO in acid mine drainage implicated to biological oxygen demand (BOD) by microorganisms for their growth and other ecological factor changes such as nitrogen and phosphate [7, 18-19]. Nitrogen (ammonia, nitrit, and nitrate) and phosphate were important for sustenance of various life forms, primary succession, and led to changes in the biogeochemical cycles of nutrients in ponds [20, 21-22], indirectly implicated to dissolved and suspended solid [23-24] and ecosystem eutrophic levels [25, 26-31]. Therefore, pH value changes, availability of elements included heavy metals, environmental characteristics, and eutrophication had a positive inter-relationships in their correspondence.

These conditions, however, had a negative correlation to heavy metals residue decreasing in Station C. The heavy metals in Station C were dominated As, Zn, Mn, Co, Ni, Cu, Pb, and Th were higher than Station A and Station B, although these metals and include Ga, Cr, Sn, Fe, Hf, and Ta were detected in all of the research station. The results indicated that chronosequence in time of ponds caused ecological changes and had positive correlations, but not for distribution of heavy metals.

The Correspondence Analysis of Heavy Metals Distribution

The correspondence analysis also showed that there was not a positive correlation between in time chronosequence in water of ex-tin mining ponds and heavy metals distribution (Figure 3). The ecological changes in chronosequence succession of ex-tin mining ponds had not significantly affect on speciation and concentration of heavy metals. Thus, there was the other factor that had affected the distribution of heavy metals.




Figure 3. Correspondence analysis between distribution of heavy metals and in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The soil and sediment of water contributed to the distribution of heavy metals [32]. In general, the mobility and availability of heavy metals were controlled by adsorption and desorption characteristics of soils, include cation exchange capacity and the contents of clay minerals [15]. The clay and mud were found to be dominant in the sediment of Station A and Station B, however, sand was found to be dominant in Station C.

The clay and mud had a positive correlation and sand had a negative correlation to a presence, affinity, and minerals structure [33-34]. The clay and mud had a larger surface area, higher cation exchange capability (CEC), and particles of clay in sediments were often closely associated with organic matter and also exist in the form of organic-mineral complexes. These particles adsorbed the active metals from the water phases, accumulated, and carried them to the bottom sediments diagenetically [35-36]. Therefore, distribution of heavy metals in Station A and Station B were lower than Station C.

CONCLUSION

In time chronosequence of ex-tin mining ponds had affected ecological dynamic such as pH value change and heavy metals distribution. The pH value of ex-tin mining ponds moved from acid to neutral condition, however, it had been taken for a long time of the chronosequence. The consequence of low pH value can implicated to microorganisms' growth, nutrients, and environment factors in primary succession. Further, it implicated to heavy metals interaction, whereasthe heavy metals residue should be in low concentration in long time chronosequence of ex-tin mining ponds.

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C). There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V.

There were three patterns of heavy metals distribution in time chronosequence in ex-tin mining ponds that a positive correlation, a negative correlation, and a dynamic correlation. The chronosequence in time had affects on heavy metals distribution. Elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. Elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increased along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5-10 years and than increased concentration in ponds with age > 15 years. Elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5-10 years and than decreased concentration in ponds with age > 15 years.

The chronosequence in time can be a factor that contributied to heavy metals distribution. However, the sediment factor had a significant effect to distribution of heavy metals in the ex-tin mining ponds, where clay and mud were more effective than sand in accumulation, adsorption, and interaction with heavy metals and carried them diagenetically to bottom sediments.

REFERENCES

1. De Lima e Silva, AA., Ribeiro de Carvalho, MA., L de Souza, SA., Dias, PMT., Da Silva Filho, RG., De Meirelles Saramago, CS., De Melo Bento, CA., Hofer, E., 2012, Heavy metal tolerance (Cr, Ag and Hg) in bacteria isolated from sewage, *Brazilian Journal of Microbiology*, 43(4), 1620-1631.
2. Guan, Y., Shao, C., Ju, M., 2014, Heavy metal contamination assessment and partition for industrial and mining gathering areas, *International Journal of Environmental Research and Public Health*, 11(7), 7286-7303.

Commented [xx1]: Check the references and use abbreviation for the journal name

3. Kurniawan, A., 2017, Chronosequence effect of post tin mining ponds to metals residu and microecosystem change, *Omni-Akuatika*, 13(1), 60-65.
4. Ashraf, MA., Maah, MJ., Yusoff, I., 2012, Speciation of heavy metals in the sediments of former tin mining catchment, *Iranian Journal of Science and Technology*, 36(A2), 163-180.
5. Daniel, VN., Chudusu, ES., Chup, JA., Pius, ND., 2014, Variations of heavy metals in agricultural soils irrigated with tin water in Heipang District of Barkin Ladi, Plateau State, Nigeria, *International Journal of Science and Technology*, 3(5), 255-263.
6. Ashraf, MA., Maah, MJ., Yusoff, I., 2012. Morphology, geology and water quality assessment of former tin mining catchment, *The Scientific World Journal*, 2012, 1-15.
7. Urbanov´a, M., Kopecký, J., Val´as´kov´a, V., S´agov´a-Marec´kov´a, M., Elhottov´a, D., Kyselkov´a, M., Moe¨nne-Loccoz, Y., Baldrian, P., 2011, Development of bacterial community during spontaneous succession on spoil heaps after brown coal mining, *FEMS Microbiology Ecology*, 78(1), 59-69.
8. Kurniawan, A., 2016, Microorganism communities response of ecological changes in post tin mining ponds, *Journal of Microbiology and Virology*, 6(1), 17-26.
9. Violante, A., Cozzolino, V., Perelomov, L., Caporale, AG., Pigna, M., 2010, Mobility and bioavailability of heavy metals and metalloids in soil environments, *Journal of Soil Science and Plant Nutrition*, 10(3), 268-292.
10. Akcil, A., and Koldas, S., 2006, Acid mine drainage (AMD): causes, treatment and case studies, *Journal of Cleaner Production*, 14(12-13), 1139-1145.
11. Tan, G., Shu, W., Hallberg, KB., Li, F., Lan, C., Huang, L., 2007, Cultivation-dependent and cultivation-independent characterization of the microbial community in acid mine drainage associated with acidic Pb/Zn mine tailings at Lechang, Guangdong, China, *FEMS Microbiology Ecology*, 59(1), 118-126.
12. Campaner, VP., Luiz-Silva, W., Machado, W., 2014, Geochemistry of acid mine drainage from a coal mining area and processes controlling metal attenuation in stream waters, Southern Brazil, *Anais da Academia Brasileira de Ciências*, 86(2), 539-554.
13. Bardgett, RD., Bowman, WD., Kaufmann, R., Schmidt, SK., 2005, A temporal approach to linking aboveground and belowground ecology, *TRENDS in Ecology and Evolution*, 20(11), 634-641.
14. Ahmad, AK., and Al-Mahaqeri, SA., 2015, Human health risk assessment of heavy metals in fish species collected from catchments of former tin mining, *International Journal of Research Studies in Science, Engineering and Technology*, 2(4), 9-21.
15. Zeng, F., Ali, S., Zhang, H., Ouyang, Y., Qiu, B., Wu, F., Zhang, G., 2011, The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants, *Environmental Pollution*, 159(1), 84-91.
16. Huang, L., Tang, F., Song, Y., Wan, C., Wang, S., Liu, W., Shu, W., 2010, Biodiversity, abundance, and activity of nitrogen-fixing bacteria during primary succession on a copper mine tailings, *FEMS Microbiology Ecology*, 78(3), 439-450.
17. Abdel-Raouf, N., Al-Homaidan, AA., Ibraheem, IBM., 2012, Microalgae and wastewater treatment, *Saudi Journal of Biological Sciences* 19(3), 257-275.
18. Tscherko, D., Hammesfahr, U., Zeltner, G., Kandeler, E., Bocker, R., 2005, Plant succession and rhizosphere microbial communities in a recently deglaciated alpine terrain. *Basic and Applied Ecology*, 6(4), 367-383.
19. Fierer, N., Nemergut, D., Knight, R., Craine, JM., 2010, Changes through time: integrating microorganisms into the study of succession, *Research in Microbiology*, 161(8), 635-642.
20. Quirós, R., 2003, The relationship between nitrate and ammonia concentrations in the pelagic zone of lakes, *Limnetica*, 22(1-2), 37-50.
21. Michalski, R., and Kurzyca, I. 2006, Determination of nitrogen species (nitrate, nitrite and ammonia ions) in environmental samples by ion chromatography, *Polish Journal of Environmental Studies*, 15(1), 5-18.
22. Mihale, MJ., 2015, Nitrogen and phosphorus dynamics in the waters of the Great Ruaha River, Tanzania, *Journal of Water Resources and Ocean Science*, 4(5), 59-71.
23. Akan, JC., Audu, SI., Mohammed, Z., Ogunbuaja, VO., 2013, Assessment of heavy metals, pH, organic matter and organic carbon in roadside soils in Makurdi Metropolis, Benue State, Nigeria, *Journal of Environmental Protection*, 4(6), 618-628.

24. Kuriata-Potasznik, A., Szymczyk, S., Skwierawski, A., Glińska-Lewczuk, K., Cymes, I., 2016, Heavy metal contamination in the surface layer of bottom sediments in a flow-through lake: a case study of Lake Symsar in Northern Poland, *Water*, 8(8), 1-15.
25. Maher, W., Krikowa, F., Wruck, D., Louie, H., Nguyen, T., Huang, WY., 2002, Determination of total phosphorus and nitrogen in turbid waters by oxidation with alkaline potassium peroxydisulfate and low pressure microwave digestion, autoclave heating or the use of closed vessels in a hot water bath: comparison with kjeldahl digestion, *Analytica Chimica Acta*, 463(2), 283-293.
26. Song, Y., Hahn, HH., Hoffmann, E., 2002, Effects of solution conditions on the precipitation of phosphate for recovery a thermodynamic evaluation, *Chemosphere*, 48(10), 1029-1034.
27. Mahadevaiah., Kumar, MSY., Galil, MSA., Suresha, MS., Sathish, MA., Nagendrappa, G., 2007, A simple spectrophotometric determination of phosphate in sugarcane juices, water, and detergent samples, *E-Journal of Chemistry*, 4(4): 467-473.
28. Topcu, A., and Pulatsu, S., 2014, Phosphorus fractions and cycling in the sediment of a shallow eutrophic pond, *Journal of Agricultural Sciences*, 2014(20), 63-70.
29. Sengupta, S., Nawaz, T., Beaudry, J., 2015, Nitrogen and phosphorus recovery from wastewater, *Current Pollution Reports*, 1(3), 155-166.
30. Da Silva Cerozi, B., and Fitzsimmons, K., 2016, The effect of pH on phosphorus availability and speciation in an aquaponics nutrient solution, *Bioresource Technology*, 2016(219), 778-781.
31. Lei, Y., Song, B., Van Der Weijden, RD., Saakes, M., Buisman, CJN., 2017, Electrochemical induced calcium phosphate precipitation: importance of local pH, *Environmental Science & Technology*, 51(19): 11156-11164.
32. De Saedeleer, V., Cappuyns, V., De Cooman, W., Swennen, R., 2010, Influence of major elements on heavy metal composition of river sediments, *Geologica Belgica*, 13(3), 257-268.
33. Jiang, M., Zeng, G., Zhang, C., Ma, X., Chen, M., Zhang, J., Lu, L., Yu, Q., Hu, L., Liu, L., 2012, Assessment of heavy metal contamination in the surrounding soils and surface sediments in Xiawangang River, Qingshuitang District, *PLoS ONE*, 8(8), e71176.
34. Alamgir, M., Islam, M., Hossain, N., Kibria, MG., Rahman, MM., 2015, Assessment of heavy metal contamination in urban soils of Chittagong City, Bangladesh, *International Journal of Plant & Soil Science*, 7(6), 362-372.
35. Zhang, C., Yu, Z., Zeng, G., Jiang, M., Yang, Z., Cui, F., Zhu, M., Shen, L., Hu, L., 2014, Effects of sediment geochemical properties on heavy metal bioavailability. *Environment International*, 2014(73), 270-281.
36. Yao, Q., Wang, X., Jian, H., Chen, H., Yu, Z., 2015, Characterization of the particles size fraction associated with heavy metals in suspended sediments of the Yello River, *International Journal of Environment Research and Public Health*, 12(6), 6725-6744.

The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia

Andri Kurniawan^{a,b,*}, Oedjijono^b, Tamad^c, and Uyi Sulaeman^d

^a Department of Aquaculture, University of Bangka Belitung
Jl. Balunijuk, Merawang, Bangka Belitung Archipelago Province

^b Faculty of Biology, University of Jenderal Soedirman
Jl. dr. Suparno, Grendeng, Purwokerto, Central Java

^c Departement of Agriculture, University of Jenderal Soedirman
Jl. dr. Suparno, Grendeng, Purwokerto, Central Java

^d Department of Chemistry, University of Jenderal Soedirman
Jl. dr. Suparno, Grendeng, Purwokerto, Central Java

* Corresponding author, tel/fax : +6281351714747, email: andri-kurniawan@ubb.ac.id

ABSTRACT

The heavy metals distribution of ex-tin mining ponds were investigated. The time chronosequence was determined at the pond of age < 1 year (Station A), the pond of age 5-10 years (Station B), and the pond of age > 15 years (Station C). The results showed sixteen heavy metals of As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V could be detected in the ponds. The metals such as As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn in Station C showed higher concentration compared to the Station A and Station B. The metals such as Cr, Fe, Hf, Sn, Ta, Te, and V in Station A and Station B showed higher concentration compared to the Station C. The positive, negative, and dynamic correlation pattern could be found in distribution of heavy metal to time chronosequence. The concentration of Ta and V showed a positive correlation because their concentration decrease, whereas concentration of As, Cu, Ga, Mn, and Zn showed a negative correlation because their concentration increase along in time chronosequence. The dynamic correlation could be found that concentration of Co, Ni, Pb, Sn, and Th decrease from Station A to Station B and then increase in Station C, whereas concentration of Cr, Fe, Hf, and Te increase from Station A to Station B and then decrease in Station C.

Keywords: distribution pattern, heavy metals, chronosequence, ex-tin mining ponds

ABSTRAK

Distribusi logam berat dari danau pascatambang timah telah dikaji. Kronosekuensi waktu danau ditetapkan pada umur < 1 tahun (Stasiun A), umur 5-10 tahun (Stasiun B), dan umur > 15 tahun (Stasiun C). Hasil menunjukkan 16 logam berat yaitu As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, dan V terdeteksi di danau tersebut. Logam As, Co, Cu, Ga, Mn, Ni, Pb, Th, dan Zn di Stasiun C memiliki konsentrasi lebih tinggi dibandingkan Stasiun A dan Stasiun B. Logam Cr, Fe, Hf, Sn, Ta, Te, dan V di Stasiun A dan Stasiun B memiliki konsentrasi lebih tinggi dibandingkan Stasiun C. Pola korelasi positif, negatif, dan dinamis diperoleh dari distribusi logam berat terhadap kronosekuensi waktu. Konsentrasi Ta dan V menunjukkan korelasi positif karena konsentrasinya meningkat, sedangkan konsentrasi As, Cu, Ga, Mn, dan Zn menunjukkan korelasi negatif karena konsentrasinya menurun seiring kronosekuensi waktu. Korelasi dinamis terlihat pada konsentrasi Co, Ni, Pb, Sn, dan Th yang menurun dari Stasiun A ke Stasiun B dan kemudian meningkat di Stasiun C, serta konsentrasi Cr, Fe, Hf, dan Te meningkat dari Stasiun A ke Stasiun B dan kemudian menurun di Stasiun C.

Kata kunci: pola distribusi, logam berat, kronosekuensi, danau pascatambang timah

INTRODUCTION

Tin mining activity is an anthropogenic activity that has contributed to the ecological problems. It has been known as the major sources of heavy metal contamination such as heavy metals in ex-tin mining ponds. Ex-tin mining ponds as one of water resource were contaminated by heavy metals that can be accumulated in food chains and secondary activities of the water [1, 2-3]. Some of the heavy

metals such as Pb, Zn, Mn, Fe, Cr, Cu, Ni, Sn, As, and Cd were detected in site of ex-tin mining [4-5].

The each of waters had their pattern of physical and chemical characteristics that determined by climatic, geomorphological, and geochemical conditions [6]. These conditions were influenced by ecological succession in time chronosequence [7] and the ecological changes of ex-tin mining ponds have occurred for a long time [8]. The implication of ecological changes as long as time chronosequence was interactions and reactions of chemical complexation between organic

compounds, environmental factors, and inorganic compounds, include presence of heavy metals [9].

The researches of heavy metals in ex-tin mining ecosystems have been done, however, not for correspondence of heavy metals distributions and ex-tin mining ponds in time chronosequence. While an understanding this correspondence was crucial to predict ecosystems conditions and to detect environment change. The understanding of heavy metals distribution in time chronosequence of ex-tin mining ponds becomes an important part of aquatic ecosystem management as water sources for secondary activities and habitable for organisms.

EXPERIMENTAL SECTION

Materials

The study areas were located in Bangka Regency, Bangka Belitung Archipelago Province of Indonesia. The considered study sites cover three ponds of ex-tin mining with chronosequence ranging in time. The study areas were encoded as Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years).

The coordinate of Station A were 01°59' S in points 36,0"; 36,2"; 36,4"; 36,5"; 36,6" and 106°06' E in points 36,5"; 36,9"; 37,3"; 37,4"; 37,5". The coordinate of Station B were 01°59' S in points 41,3"; 41,4"; 41,5"; 42,4"; 42,5" and 106°06' E in points 39,2"; 39,5"; 41,4"; 42,7"; 43,1". The coordinate of Station C were 01°55' S in points 40,9"; 58,9"; 59,1"; 59,2"; 59,5" and 106°06' E in points 19,5"; 19,7"; 19,9"; 22,4"; 29,2".

In the each of research stations points, water sampling was done < 4m in depth (station code A.1; B.1; and C.1) and composite sampling were done to water and sediment > 4m in depth (station code A.2; B.2; and C.2).

Instrumentation

The water samples and composite samples were analyzed pH value by pH meter (PH-009(I)-A) Sun Care. The speciation and concentration of heavy metals analyzed by X-Ray Fluorescence (Rigaku) with three light spreader metals of copper (Cu), molybdenum (Mo), and aluminum (Al). Also, the pH and moisture of soil around the ponds were measured by pH-moisture meter. The correspondence between heavy metals distribution and the time chronosequence of ex-tin mining ponds was analyzed by PAST (*Palaentological Statistics*) 3.

RESULTS AND DISCUSSION

pH Value of Water and Soil

Tin mining activity had generated the ex-tin mining ponds as one of the water resources. However, water quality in the ponds had serious ecological problem. The ex-mining activity produced

acid mine drainage (AMD) that can caused the long-term impairment of water quality and biodiversity. The pH value of ex-tin mining water and soil pH value indicated that the acid could be found in Station A (pond in age < 1 year) and Station B (pond in age 5-10 years) and the neutral value could be found in Station C (pond in age > 15 years). In the each of research stations points, water samples < 4m in depth indicated more acid than the composite samples (water and sediment > 4m in depth) (Figure 1).




Figure 1. Soil and water pH value in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B(pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The acid mine drainage (AMD) was produced when sulfide-bearing material is exposed to oxygen and water [10]. In acid mine drainage, presence and mobility of elements such as Al, As, Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Zn, and also ionic activity and sulphide minerals oxidation caused pH value decreasing [11-12]. Furthermore, the low pH condition implicated in oxidation state, toxicity, and mobility of elements [9].

In long time chronosequence more than 15 years in the Station C caused changes water quality of ex-tin mining ponds were more stable than Station A and Station B. The indicator for it was pH value increased to neutral conditions (7,0-7,4) and other parameters that indicated water quality was more habitable for organisms' life such as fishes. The pH value and other parameters changes, indirectly, caused ecological productivity increasing. The chronosequence in long time caused changes in nutrient availability and resource supply for ecosystem productivity [13].

The pH value was found to play the most important role in the determination of metal speciation, solubility from mineral surfaces, movement, and bioavailability of metals. Furthermore, acid mine drainage had a major contribution in heavy metal contamination [14].

There was a negative correlation between soil pH and distribution heavy metal where pH value decreasing caused increasing in heavy metal desorption [15].

The Pattern of Heavy Metals Distribution

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr,

Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C) (Table 1).

Table 1. Distribution of heavy metals in time chronosequence in water of ex-tin mining ponds

No	Heavy Metals	Heavy Metals Concentration (ppm)					
		In Station A		In Station B		In Station C	
		Station A.1	Station A.2	Station B.1	Station B.2	Station C.1	Station C.2
1	As	nd	2,155	nd	3,215	2,4	1,095
2	Ga	nd	4,275	nd	4,47	9,76	nd
3	V	nd	0,84	nd	nd	nd	0,67
4	Te	nd	nd	102	nd	nd	5,8
5	Cr	nd	4,735	nd	4,945	nd	0,67
6	Zn	nd	6,05	nd	8,1	8,17	3,905
7	Sn	47	82,8	50,3	51,4	61,8	57,15
8	Mn	nd	13,05	nd	13,6	30,8	14,85
9	Fe	21,4	1481,3	148	2344	1590	849,05
10	Co	nd	5,05	nd	nd	6,64	nd
11	Ni	nd	4,09	nd	2,875	3,48	1,905
12	Cu	3,08	6,875	3,19	7,775	8,94	3,81
13	Pb	nd	6,7	nd	6	8,21	4,62
14	Th	nd	4,605	nd	4,38	13,5	7,05
15	Hf	6,71	3,21	8,21	3,32	nd	9,38
16	Ta	4,94	2,615	4,16	2,995	nd	2,805

**) nd (not detected). The Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).*

There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. The average of As concentration were 1,7475 ppm (Station C) > 1,6075 ppm (Station B) > 1,0775 ppm (Station A). The concentration of Co were 3,32 ppm (Station C) > 2,525 ppm (Station A) > not detected (Station B). The concentration of Cu were 6,375 ppm (Station C) > 5,4825 ppm (Station B) > 4,9775 ppm (Station A). The concentration of Ga were 4,88 ppm (Station C) > 2,235 ppm (Station B) > 2,1375 ppm (Station A). The concentration of Mn were 22,825 ppm (Station C) > 6,8 ppm (Station B) > 6,525 ppm (Station A). The concentration of Ni were 2,6925 ppm (Station C) > 2,045 ppm (Station A) > 1,4375 ppm (Station B). The concentration of Pb were 6,415 ppm (Station C) > 3,35 ppm (Station A) > 3,0 ppm (Station B). The concentration of Th were 10,275 ppm (Station C) > 2,3025 ppm (Station A) > 2,19 ppm (Station B). The concentration of Zn were 6,0375 ppm (Station C) > 4,05 ppm (Station B) > 3,025 ppm (Station A).

In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V. The average of Cr concentration were 2,4725 ppm (Station B) > 2,3675 ppm (Station A) > 0,335 ppm (Station C). The concentration of Fe were 1246 ppm (Station B) > 1219,53 ppm (Station C) > 751,35 ppm (Station A). The concentration of Hf were 5,765 ppm (Station B) > 4,96 ppm (Station A) > 4,69 ppm (Station C). The concentration of Sn were 64,9 ppm (Station A) > 59,475 ppm (Station C) > 50,85 ppm (Station B). The concentration of Ta were 3,7775 ppm (Station A) > 3,5775 ppm (Station B) > 1,4025 ppm (Station C). The concentration of Te were 51 ppm (Station B) > 2,9 ppm (Station C) > not detected (Station A). The concentration of V were 0,42 ppm (Station A) > 0,335 ppm (Station C) > not detected (Station B).

Their concentration of heavy metals in time chronosequence in ex-tin mining ponds showed three patterns distribution that a positive correlation, a negative correlation, and a dynamic correlation. The positive correlation indicated that concentration

of heavy metals decreased along chronosequence in time in ex-tin mining ponds. The negative correlation indicated that concentration of heavy metals increased along chronosequence in time in ex-tin mining ponds. The dynamic correlation indicated that concentration of heavy metals on decreased or increased along chronosequence in time in ex-tin mining ponds.

The elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. The

elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increasing along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5-10 years and then increased concentration in ponds with age > 15 years. The elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5-10 years and then decreased concentration in ponds with age > 15 years (Figure 2).




Figure 2. Distribution of heavy metals in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The chronosequence in long time caused changes in heavy metals concentration were decreased as the primary succession progression [16]. The low pH value implicated to dissolved oxygen (DO) [17] and low DO in acid mine drainage implicated to biological oxygen demand (BOD) by microorganisms for their growth and other ecological factor changes such as nitrogen and phosphate [7, 18-19]. Nitrogen (ammonia, nitrit, and nitrate) and phosphate were important for sustenance of various life forms, primary succession, and led to changes in the biogeochemical cycles of nutrients in ponds [20, 21-22], indirectly implicated to dissolved and suspended solid [23-24] and ecosystem eutrophic levels [25, 26-31]. Therefore, pH value changes, availability of elements included heavy metals, environmental characteristics, and

eutrophication had a positive inter-relationships in their correspondence.

These conditions, however, had a negative correlation to heavy metals residue decreasing in Station C. The heavy metals in Station C were dominated As, Zn, Mn, Co, Ni, Cu, Pb, and Th were higher than Station A and Station B, although these metals and include Ga, Cr, Sn, Fe, Hf, and Ta were detected in all of the research station. The results indicated that chronosequence in time of ponds caused ecological changes and had positive correlations, but not for distribution of heavy metals.

The Correspondence Analysis of Heavy Metals Distribution

The correspondence analysis also showed that there was not a positive correlation between in

time chronosequence in water of ex-tin mining ponds and heavy metals distribution (Figure 3). The ecological changes in chronosequence succession of ex-tin mining ponds had not significantly affect on

speciation and concentration of heavy metals. Thus, there was the other factor that had affected the distribution of heavy metals.




Figure 3. Correspondence analysis between distribution of heavy metals and in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5-10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4m in depth).

The soil and sediment of water contributed to the distribution of heavy metals [32]. In general, the mobility and availability of heavy metals were controlled by adsorption and desorption characteristics of soils, include cation exchange capacity and the contents of clay minerals [15]. The clay and mud were found to be dominant in the sediment of Station A and Station B, however, sand was found to be dominant in Station C.

The clay and mud had a positive correlation and sand had a negative correlation to a presence, affinity, and minerals structure [33-34]. The clay and mud had a larger surface area, higher cation exchange capability (CEC), and particles of clay in sediments were often closely associated with organic matter and also exist in the form of organic-mineral complexes. These particles adsorbed the active metals from the water phases, accumulated, and carried them to the bottom sediments diagenetically [35-36]. Therefore, distribution of heavy metals in Station A and Station B were lower than Station C.

CONCLUSION

In time chronosequence of ex-tin mining ponds had affected ecological dynamic such as pH value change and heavy metals distribution. The pH value of ex-tin mining ponds moved from acid to neutral condition, however, it had been taken for a long time of the chronosequence. The consequence of low pH value can implicated to microorganisms' growth, nutrients, and environment factors in primary

succession. Further, it implicated to heavy metals interaction, whereasthe heavy metals residue should be in low concentration in long time chronosequence of ex-tin mining ponds.

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C). There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V.

There were three patterns of heavy metals distribution in time chronosequence in ex-tin mining ponds that a positive correlation, a negative correlation, and a dynamic correlation. The chronosequence in time had affects on heavy metals distribution. Elements of Ta and V had a positive correlation in time chronosequence in ex-tin mining ponds with their concentration on decreasing. Elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increased along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5-10 years and than increased

concentration in ponds with age > 15 years. Elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5-10 years and then decreased concentration in ponds with age > 15 years.

The chronosequence in time can be a factor that contributed to heavy metals distribution. However, the sediment factor had a significant effect to distribution of heavy metals in the ex-tin mining ponds, where clay and mud were more effective than sand in accumulation, adsorption, and interaction with heavy metals and carried them diagenetically to bottom sediments.

REFERENCES

- [1] De Lima e Silva, AA., Ribeiro de Carvalho, MA., L de Souza, SA., Dias, PMT., Da Silva Filho, RG., De Meirelles Saramago, CS., De Melo Bento, CA., Hofer, E., 2012, Heavy metal tolerance (Cr, Ag and Hg) in bacteria isolated from sewage, *Brazilian Journal of Microbiology*, 43(4), 1620-1631.
- [2] Guan, Y., Shao, C., Ju, M., 2014, Heavy metal contamination assessment and partition for industrial and mining gathering areas, *International Journal of Environmental Research and Public Health*, 11(7), 7286-7303.
- [3] Kurniawan, A., 2017, Chronosequence effect of post tin mining ponds to metals residu and microecosystem change, *Omni-Akuatika*, 13(1), 60-65.
- [4] Ashraf, MA., Maah, MJ., Yusoff, I., 2012, Speciation of heavy metals in the sediments of former tin mining catchment, *Iranian Journal of Science and Technology*, 36(A2), 163-180.
- [5] Daniel, VN., Chudusu, ES., Chup, JA., Pius, ND., 2014, Variations of heavy metals in agricultural soils irrigated with tin water in Heipang District of Barkin Ladi, Plateau State, Nigeria, *International Journal of Science and Technology*, 3(5), 255-263.
- [6] Ashraf, MA., Maah, MJ., Yusoff, I., 2012. Morphology, geology and water quality assessment of former tin mining catchment, *The Scientific World Journal*, 2012, 1-15.
- [7] Urbanov´a, M., Kopecký, J., Val´asˇkov´a, V., S´agov´a-Marecˇkov´a, M., Elhottov´a, D., Kyselkov´a, M., Moe¨nne-Loccoz, Y., Baldrian, P., 2011, Development of bacterial community during spontaneous succession on spoil heaps after brown coal mining, *FEMS Microbiology Ecology*, 78(1), 59-69.
- [8] Kurniawan, A., 2016, Microorganism communities response of ecological changes in post tin mining ponds, *Journal of Microbiology and Virology*, 6(1), 17-26.
- [9] Violante, A., Cozzolino, V., Perelomov, L., Caporale, AG., Pigna, M., 2010, Mobility and bioavailability of heavy metals and metalloids in soil environments, *Journal of Soil Science and Plant Nutrition*, 10(3), 268-292.
- [10] Akcil, A., and Koldas, S., 2006, Acid mine drainage (AMD): causes, treatment and case studies, *Journal of Cleaner Production*, 14(12-13), 1139-1145.
- [11] Tan, G., Shu, W., Hallberg, KB., Li, F., Lan, C., Huang, L., 2007, Cultivation-dependent and cultivation-independent characterization of the microbial community in acid mine drainage associated with acidic Pb/Zn mine tailings at Lechang, Guangdong, China, *FEMS Microbiology Ecology*, 59(1), 118-126.
- [12] Campaner, VP., Luiz-Silva, W., Machado, W., 2014, Geochemistry of acid mine drainage from a coal mining area and processes controlling metal attenuation in stream waters, Southern Brazil, *Anais da Academia Brasileira de Ciências*, 86(2), 539-554.
- [13] Bardgett, RD., Bowman, WD., Kaufmann, R., Schmidt, SK., 2005, A temporal approach to linking aboveground and belowground ecology, *TRENDS in Ecology and Evolution*, 20(11), 634-641.
- [14] Ahmad, AK., and Al-Mahaqeri, SA., 2015, Human health risk assessment of heavy metals in fish species collected from catchments of former tin mining, *International Journal of Research Studies in Science, Engineering and Technology*, 2(4), 9-21.
- [15] Zeng, F., Ali, S., Zhang, H., Ouyang, Y., Qiu, B., Wu, F., Zhang, G., 2011, The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants, *Environmental Pollution*, 159(1), 84-91.
- [16] Huang, L., Tang, F., Song, Y., Wan, C., Wang, S., Liu, W., Shu, W., 2010, Biodiversity, abundance, and activity of nitrogen-fixing bacteria during primary succession on a copper mine tailings, *FEMS Microbiology Ecology*, 78(3), 439-450.
- [17] Abdel-Raouf, N., Al-Homaidan, AA., Ibraheem, IBM., 2012, Microalgae and wastewater treatment, *Saudi Journal of Biological Sciences* 19(3), 257-275.
- [18] Tschërko, D., Hammesfahr, U., Zeltner, G., Kandeler, E., Bocker, R., 2005, Plant succession and rhizosphere microbial communities in a recently deglaciated alpine terrain. *Basic and Applied Ecology*, 6(4), 367-383.
- [19] Fierer, N., Nemergut, D., Knight, R., Craine, JM., 2010, Changes through time: integrating microorganisms into the study of succession, *Research in Microbiology*, 161(8), 635-642.
- [20] Quirós, R., 2003, The relationship between nitrate and ammonia concentrations in the pelagic zone of lakes, *Limnetica*, 22(1-2), 37-50.

- [21] Michalski, R., and Kurzyca, I. 2006, Determination of nitrogen species (nitrate, nitrite and ammonia ions) in environmental samples by ion chromatography, *Polish Journal of Environmental Studies*, 15(1), 5-18.
- [22] Mihale, MJ., 2015, Nitrogen and phosphorus dynamics in the waters of the Great Ruaha River, Tanzania, *Journal of Water Resources and Ocean Science*, 4(5), 59-71.
- [23] Akan, JC., Audu, SI., Mohammed, Z., Ogugbuaja, VO., 2013, Assessment of heavy metals, pH, organic matter and organic carbon in roadside soils in Makurdi Metropolis, Benue State, Nigeria, *Journal of Environmental Protection*, 4(6), 618-628.
- [24] Kuriata-Potasznik, A., Szymczyk, S., Skwierawski, A., Glińska-Lewczuk, K., Cymes, I., 2016, Heavy metal contamination in the surface layer of bottom sediments in a flow-through lake: a case study of Lake Symsar in Northern Poland, *Water*, 8(8), 1-15.
- [25] Maher, W., Krikowa, F., Wruck, D., Louie, H., Nguyen, T., Huang, WY., 2002, Determination of total phosphorus and nitrogen in turbid waters by oxidation with alkaline potassium peroxodisulfate and low pressure microwave digestion, autoclave heating or the use of closed vessels in a hot water bath: comparison with kjeldahl digestion, *Analytica Chimica Acta*, 463(2), 283-293.
- [26] Song, Y., Hahn, HH., Hoffmann, E., 2002, Effects of solution conditions on the precipitation of phosphate for recovery a thermodynamic evaluation, *Chemosphere*, 48(10), 1029-1034.
- [27] Mahadevaiah., Kumar, MSY., Galil, MSA., Suresha, MS., Sathish, MA., Nagendrappa, G., 2007, A simple spectrophotometric determination of phosphate in sugarcane juices, water, and detergent samples, *E-Journal of Chemistry*, 4(4): 467-473.
- [28] Topcu, A., and Pulatsu, S., 2014, Phosphorus fractions and cycling in the sediment of a shallow eutrophic pond, *Journal of Agricultural Sciences*, 2014(20), 63-70.
- [29] Sengupta, S., Nawaz, T., Beaudry, J., 2015, Nitrogen and phosphorus recovery from wastewater, *Current Pollution Reports*, 1(3), 155-166.
- [30] Da Silva Cerozi, B., and Fitzsimmons, K., 2016, The effect of pH on phosphorus availability and speciation in an aquaponics nutrient solution, *Bioresource Technology*, 2016(219), 778-781.
- [31] Lei, Y., Song, B., Van Der Weijden, RD., Saakes, M., Buisman, CJN., 2017, Electrochemical induced calcium phosphate precipitation: importance of local pH, *Environmental Science & Technology*, 51(19): 11156-11164.
- [32] De Saedeleer, V., Cappuyns, V., De Cooman, W., Swennen, R., 2010, Influence of major elements on heavy metal composition of river sediments, *Geologica Belgica*, 13(3), 257-268.
- [33] Jiang, M., Zeng, G., Zhang, C., Ma, X., Chen, M., Zhang, J., Lu, L., Yu, Q., Hu, L., Liu, L., 2012, Assessment of heavy metal contamination in the surrounding soils and surface sediments in Xiawangang River, Qingshuitang District, *PLoS ONE*, 8(8), e71176.
- [34] Alamgir, M., Islam, M., Hossain, N., Kibria, MG., Rahman, MM., 2015, Assessment of heavy metal contamination in urban soils of Chittagong City, Bangladesh, *International Journal of Plant & Soil Science*, 7(6), 362-372.
- [35] Zhang, C., Yu, Z., Zeng, G., Jiang, M., Yang, Z., Cui, F., Zhu, M., Shen, L., Hu, L., 2014, Effects of sediment geochemical properties on heavy metal bioavailability. *Environment International*, 2014(73), 270-281.
- [36] Yao, Q., Wang, X., Jian, H., Chen, H., Yu, Z., 2015, Characterization of the particles size fraction associated with heavy metals in suspended sediments of the Yello River, *International Journal of Environment Research and Public Health*, 12(6), 6725-6744.

SHORT COMMUNICATION**The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia****Andri Kurniawan^{1,2,*}, Oedjijono², Tamad³, and Uyi Sulaeman⁴**¹Department of Aquaculture, University of Bangka Belitung,
Jl. Balunijuk, Merawang, Bangka, Bangka Belitung 33172, Indonesia²Faculty of Biology, Jenderal Soedirman University,
Jl. Dr. Suparno no. 63, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia³Department of Agriculture, Jenderal Soedirman University,
Jl. Dr. Suparno no. 61, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia⁴Department of Chemistry, Faculty of Science and Technology, Jenderal Soedirman University,
Jl. Dr. Suparno no. 61, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia

Received March 1, 2018; Accepted July 17, 2018

ABSTRACT

The heavy metals distribution of ex-tin mining ponds were investigated. The time chronosequence was determined at the pond of age < 1 year (Station A), the pond of age 5–10 years (Station B), and the pond of age > 15 years (Station C). The results showed sixteen heavy metals of As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V could be detected in the ponds. The metals such as As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn in Station C showed higher concentration compared to the Station A and Station B. The metals such as Cr, Fe, Hf, Sn, Ta, Te, and V in Station A and Station B showed higher concentration compared to the Station C. The positive, negative, and dynamic correlation pattern could be found in distribution of heavy metal to time chronosequence. The concentration of Ta and V showed a positive correlation because their concentration decrease, whereas concentration of As, Cu, Ga, Mn, and Zn showed a negative correlation because their concentration increase along in time chronosequence. The dynamic correlation could be found that concentration of Co, Ni, Pb, Sn, and Th decrease from Station A to Station B and then increase in Station C, whereas concentration of Cr, Fe, Hf, and Te increase from Station A to Station B and then decrease in Station C.

Keywords: distribution pattern; heavy metals; chronosequence; ex-tin mining ponds**ABSTRAK**

Distribusi logam berat dari danau pascatambang timah telah dikaji. Kronosekuensi waktu danau ditetapkan pada umur < 1 tahun (Stasiun A), umur 5–10 tahun (Stasiun B), dan umur > 15 tahun (Stasiun C). Hasil menunjukkan 16 logam berat yaitu As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, dan V terdeteksi di danau tersebut. Logam As, Co, Cu, Ga, Mn, Ni, Pb, Th, dan Zn di Stasiun C memiliki konsentrasi lebih tinggi dibandingkan Stasiun A dan Stasiun B. Logam Cr, Fe, Hf, Sn, Ta, Te, dan V di Stasiun A dan Stasiun B memiliki konsentrasi lebih tinggi dibandingkan Stasiun C. Pola korelasi positif, negatif, dan dinamis diperoleh dari distribusi logam berat terhadap kronosekuensi waktu. Konsentrasi Ta dan V menunjukkan korelasi positif karena konsentrasinya meningkat, sedangkan konsentrasi As, Cu, Ga, Mn, dan Zn menunjukkan korelasi negatif karena konsentrasinya menurun seiring kronosekuensi waktu. Korelasi dinamis terlihat pada konsentrasi Co, Ni, Pb, Sn, dan Th yang menurun dari Stasiun A ke Stasiun B dan kemudian meningkat di Stasiun C, serta konsentrasi Cr, Fe, Hf, dan Te meningkat dari Stasiun A ke Stasiun B dan kemudian menurun di Stasiun C.

Kata Kunci: pola distribusi; logam berat; kronosekuensi; danau pascatambang timah**INTRODUCTION**

Tin mining activity is an anthropogenic activity that has contributed to the ecological problems. It has been

known as the major sources of heavy metal contamination such as heavy metals in ex-tin mining ponds. Ex-tin mining ponds as one of water resource were contaminated by heavy metals that can be accumulated in food chains and secondary activities of

* Corresponding author. Tel : +62-81351714747
Email address : andri-kurniawan@ubb.ac.id

the water [1-3]. Some of the heavy metals such as Pb, Zn, Mn, Fe, Cr, Cu, Ni, Sn, As, and Cd were detected in site of ex-tin mining [4-5].

The each of waters had their pattern of physical and chemical characteristics that determined by climatic, geomorphological, and geochemical conditions [6]. These conditions were influenced by ecological succession in time chronosequence [7] and the ecological changes of ex-tin mining ponds have occurred for a long time [8]. The implication of ecological changes as long as time chronosequence was interactions and reactions of chemical complexation between organic compounds, environmental factors, and inorganic compounds, include presence of heavy metals [9].

The researches of heavy metals in ex-tin mining ecosystems have been done, however, not for correspondence of heavy metals distributions and ex-tin mining ponds in time chronosequence. While an understanding this correspondence was crucial to predict ecosystems conditions and to detect environment change. The understanding of heavy metals distribution in time chronosequence of ex-tin mining ponds becomes an important part of aquatic ecosystem management as water sources for secondary activities and habitable for organisms.

EXPERIMENTAL SECTION

Materials

The study areas were located in Bangka Regency, Bangka Belitung Archipelago Province of Indonesia. The considered study sites cover three ponds of ex-tin mining with chronosequence ranging in time. The study areas were encoded as Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years).

The coordinate of Station A were 01°59' S in points 36,0"; 36,2"; 36,4"; 36,5"; 36,6" and 106°06' E in points 36,5"; 36,9"; 37,3"; 37,4"; 37,5". The coordinate of Station B were 01°59' S in points 41,3"; 41,4"; 41,5"; 42,4"; 42,5" and 106°06' E in points 39,2"; 39,5"; 41,4"; 42,7"; 43,1". The coordinate of Station C were 01°55' S in points 40,9"; 58,9"; 59,1"; 59,2"; 59,5" and 106°06' E in points 19,5"; 19,7"; 19,9"; 22,4"; 29,2".

In the each of research stations points, water sampling was done < 4 m in depth (station code A.1; B.1; and C.1) and composite sampling were done to water and sediment > 4 m in depth (station code A.2; B.2; and C.2).

Instrumentation

The water samples and composite samples were analyzed pH value by pH meter (PH-009(I)-A) Sun Care. The speciation and concentration of heavy metals

analyzed by X-Ray Fluorescence (Rigaku) with three light spreader metals of copper (Cu), molybdenum (Mo), and aluminum (Al). Also, the pH and moisture of soil around the ponds were measured by pH-moisture meter. The correspondence between heavy metals distribution and the time chronosequence of ex-tin mining ponds was analyzed by PAST (*Paleontological Statistics*) 3.

RESULT AND DISCUSSION

pH Value of Water and Soil

Tin mining activity had generated the ex-tin mining ponds as one of the water resources. However, water quality in the ponds had serious ecological problem. The ex-mining activity produced acid mine drainage (AMD) that can caused the long-term impairment of water quality and biodiversity. The pH value of ex-tin mining water and soil pH value indicated that the acid could be found in Station A (pond in age < 1 year) and Station B (pond in age 5–10 years) and the neutral value could be found in Station C (pond in age > 15 years). In the each of research stations points, water samples < 4 m in depth indicated more acid than the composite samples (water and sediment > 4 m in depth) (Fig. 1).

The acid mine drainage (AMD) was produced when sulfide-bearing material is exposed to oxygen and water [10]. In acid mine drainage, presence and mobility of elements such as Al, As, Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Zn, and also ionic activity and sulphide minerals oxidation caused pH value decreasing [11-12]. Furthermore, the low pH condition implicated in




Fig 1. Soil and water pH value in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth).

Table 1. Distribution of heavy metals in time chronosequence in water of ex-tin mining ponds

No	Heavy Metals	Heavy Metals Concentration (ppm)					
		In Station A		In Station B		In Station C	
		Station A.1	Station A.2	Station B.1	Station B.2	Station C.1	Station C.2
1	As	nd	2.155	nd	3.215	2.4	1.095
2	Ga	nd	4.275	nd	4.47	9.76	nd
3	V	nd	0.84	nd	nd	nd	0.67
4	Te	nd	nd	102	nd	nd	5.8
5	Cr	nd	4.735	nd	4.945	nd	0.67
6	Zn	nd	6.05	nd	8.1	8.17	3.905
7	Sn	47	82.8	50.3	51.4	61.8	57.15
8	Mn	nd	13.05	nd	13.6	30.8	14.85
9	Fe	21.4	1481.3	148	2344	1590	849.05
10	Co	nd	5.05	nd	nd	6.64	nd
11	Ni	nd	4.09	nd	2.875	3.48	1.905
12	Cu	3.08	6.875	3.19	7.775	8.94	3.81
13	Pb	nd	6.7	nd	6	8.21	4.62
14	Th	nd	4.605	nd	4.38	13.5	7.05
15	Hf	6.71	3.21	8.21	3.32	nd	9.38
16	Ta	4.94	2.615	4.16	2.995	nd	2.805

*) nd (not detected). The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth).

oxidation state, toxicity, and mobility of elements [9].

In long time chronosequence more than 15 years in the Station C caused changes water quality of ex-tin mining ponds were more stable than Station A and Station B. The indicator for it was pH value increased to neutral conditions (7.0-7.4) and other parameters that indicated water quality was more habitable for organisms' life such as fishes. The pH value and other parameters changes, indirectly, caused ecological productivity increasing. The chronosequence in long time caused changes in nutrient availability and resource supply for ecosystem productivity [13].

The pH value was found to play the most important role in the determination of metal speciation, solubility from mineral surfaces, movement, and bioavailability of metals. Furthermore, acid mine drainage had a major contribution in heavy metal contamination [14]. There was a negative correlation between soil pH and distribution heavy metal where pH value decreasing caused increasing in heavy metal desorption [15].

The Pattern of Heavy Metals Distribution

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C) (Table 1).

There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. The

average of As concentration were 1.7475 ppm (Station C) > 1.6075 ppm (Station B) > 1.0775 ppm (Station A). The concentration of Co were 3.32 ppm (Station C) > 2.525 ppm (Station A) > not detected (Station B). The concentration of Cu were 6.375 ppm (Station C) > 5.4825 ppm (Station B) > 4.9775 ppm (Station A). The concentration of Ga were 4.88 ppm (Station C) > 2.235 ppm (Station B) > 2.1375 ppm (Station A). The concentration of Mn were 22.825 ppm (Station C) > 6.8 ppm (Station B) > 6.525 ppm (Station A). The concentration of Ni were 2.6925 ppm (Station C) > 2.045 ppm (Station A) > 1.4375 ppm (Station B). The concentration of Pb were 6.415 ppm (Station C) > 3.35 ppm (Station A) > 3.0 ppm (Station B). The concentration of Th were 10.275 ppm (Station C) > 2.3025 ppm (Station A) > 2.19 ppm (Station B). The concentration of Zn were 6.0375 ppm (Station C) > 4.05 ppm (Station B) > 3.025 ppm (Station A).

In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V. The average of Cr concentration were 2.4725 ppm (Station B) > 2.3675 ppm (Station A) > 0.335 ppm (Station C). The concentration of Fe were 1246 ppm (Station B) > 1219.53 ppm (Station C) > 751.35 ppm (Station A). The concentration of Hf were 5.765 ppm (Station B) > 4.96 ppm (Station A) > 4.69 ppm (Station C). The concentration of Sn were 64.9 ppm (Station A) > 59.475 ppm (Station C) > 50.85 ppm (Station B). The concentration of Ta were 3.7775 ppm (Station A) > 3.5775 ppm (Station B) > 1.4025 ppm (Station C). The concentration of Te were 51 ppm (Station B) > 2.9 ppm (Station C) > not detected (Station A). The concentration




Fig 2. Distribution of heavy metals in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth).

of V were 0.42 ppm (Station A) > 0.335 ppm (Station C) > not detected (Station B).

Their concentration of heavy metals in time chronosequence in ex-tin mining ponds showed three patterns distribution that a positive correlation, a negative correlation, and a dynamic correlation. The positive correlation indicated that concentration of heavy metals decreased along chronosequence in time in ex-tin mining ponds. The negative correlation indicated that concentration of heavy metals increased along chronosequence in time in ex-tin mining ponds. The dynamic correlation indicated that concentration of heavy metals on decreased or increased along chronosequence in time in ex-tin mining ponds.

The elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. The elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increasing along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then increased concentration in ponds with age > 15 years. The elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5–10

years and then decreased concentration in ponds with age > 15 years (Fig. 2).

The chronosequence in long time caused changes in heavy metals concentration were decreased as the primary succession progression [16]. The low pH value implicated to dissolved oxygen (DO) [17] and low DO in acid mine drainage implicated to biological oxygen demand (BOD) by microorganisms for their growth and other ecological factor changes such as nitrogen and phosphate [7,18-19]. Nitrogen (ammonia, nitrite, and nitrate) and phosphate were important for sustenance of various life forms, primary succession, and led to changes in the biogeochemical cycles of nutrients in ponds [20-22], indirectly implicated to dissolved and suspended solid [23-24] and ecosystem eutrophic levels [25-31]. Therefore, pH value changes, availability of elements included heavy metals, environmental characteristics, and eutrophication had a positive inter-relationships in their correspondence.

These conditions, however, had a negative correlation to heavy metals residue decreasing in Station C. The heavy metals in Station C were dominated As, Zn, Mn, Co, Ni, Cu, Pb, and Th were higher than Station A and Station B, although these metals and include Ga, Cr, Sn, Fe, Hf, and Ta were




Fig 3. Correspondence analysis between distribution of heavy metals and in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth).

detected in all of the research station. The results indicated that chronosequence in time of ponds caused ecological changes and had positive correlations, but not for distribution of heavy metals.

The Correspondence Analysis of Heavy Metals Distribution

The correspondence analysis also showed that there was not a positive correlation between in time chronosequence in water of ex-tin mining ponds and heavy metals distribution (Fig. 3). The ecological changes in chronosequence succession of ex-tin mining ponds had not significantly affect on speciation and concentration of heavy metals. Thus, there was the other factor that had affected the distribution of heavy metals.

The soil and sediment of water contributed to the distribution of heavy metals [32]. In general, the mobility and availability of heavy metals were controlled by adsorption and desorption characteristics of soils, include cation exchange capacity and the contents of clay minerals [15]. The clay and mud were found to be dominant in the sediment of Station A and Station B, however, sand was found to be dominant in Station C.

The clay and mud had a positive correlation and sand had a negative correlation to a presence, affinity, and minerals structure [33-34]. The clay and mud had a larger surface area, higher cation exchange capability (CEC), and particles of clay in sediments were often closely associated with organic matter and also exist in the form of organic-mineral complexes. These particles adsorbed the active metals from the water phases, accumulated, and carried them to the bottom sediments diagenetically [35-36]. Therefore, distribution of heavy

metals in Station A and Station B were lower than Station C.

CONCLUSION

In time chronosequence of ex-tin mining ponds had affected ecological dynamic such as pH value change and heavy metals distribution. The pH value of ex-tin mining ponds moved from acid to neutral condition, however, it had been taken for a long time of the chronosequence. The consequence of low pH value can implicated to microorganisms' growth, nutrients, and environment factors in primary succession. Further, it implicated to heavy metals interaction, whereas the heavy metals residue should be in low concentration in long time chronosequence of ex-tin mining ponds.

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C). There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V.

There were three patterns of heavy metals distribution in time chronosequence in ex-tin mining ponds that a positive correlation, a negative correlation, and a dynamic correlation. The chronosequence in time had affects on heavy metals distribution. Elements of Ta and V had a positive correlation in time chronosequence

in ex-mining ponds with their concentration on decreasing. Elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increased along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then increased concentration in ponds with age > 15 years. Elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then decreased concentration in ponds with age > 15 years.

The chronosequence in time can be a factor that contributed to heavy metals distribution. However, the sediment factor had a significant effect to distribution of heavy metals in the ex-tin mining ponds, where clay and mud were more effective than sand in accumulation, adsorption, and interaction with heavy metals and carried them diagenetically to bottom sediments.

REFERENCES

- [1] de Silva, A.A.L., de Carvalho, M.A.R., de Souza, S.A.L., Dias, P.M.T., Filho, R.G.S., Saramago, C.S.M., Bento, C.A.M., and Hofer, E., 2012, Heavy metal tolerance (Cr, Ag and Hg) in bacteria isolated from sewage, *Braz. J. Microbiol.*, 43 (4), 1620–1631.
- [2] Guan, Y., Shao, C., and Ju, M., 2014, Heavy metal contamination assessment and partition for industrial and mining gathering areas, *Int. J. Environ. Res. Public Health*, 11 (7), 7286–7303.
- [3] Kurniawan, A., 2017, Chronosequence effect of post tin mining ponds to metals residu and microecosystem change, *Omni-Akuatika*, 13 (1), 60–65.
- [4] Ashraf, M.A., Maah, M.J., and Yusoff, I., 2012, Speciation of heavy metals in the sediments of former tin mining catchment, *Iran. J. Sci. Technol. Trans. A Sci.*, 36 (2), 163–180.
- [5] Daniel, V.N., Chudusu, E.S., Chup, J.A., and Pius, N.D., 2014, Variations of heavy metals in agricultural soils irrigated with tin water in Heipang District of Barkin Ladi, Plateau State, Nigeria, *Int. J. Sci. Technol.*, 3 (5), 255–263.
- [6] Ashraf, M.A., Maah, M.J., and Yusoff, I., 2012, Morphology, geology and water quality assessment of former tin mining catchment, *Sci. World J.*, 2012, 369206.
- [7] Urbanová, M., Kopecký, J., Valášková, V., Ságová-Marečková, M., Elhottová, D., Kyselková, M., Moëne-Loccoz, Y., and Baldrian, P., 2011, Development of bacterial community during spontaneous succession on spoil heaps after brown coal mining, *FEMS Microbiol. Ecol.*, 78 (1), 59–69.
- [8] Kurniawan, A., 2016, Microorganism communities response of ecological changes in post tin mining ponds, *RRJoMV*, 6 (1), 17–26.
- [9] Violante, A., Cozzolino, V., Perelomov, L., Caporale, A.G., and Pigna, M., 2010, Mobility and bioavailability of heavy metals and metalloids in soil environments, *J. Soil Sci. Plant Nutr.*, 10 (3), 268–292.
- [10] Akcil, A., and Koldas, S., 2006, Acid mine drainage (AMD): causes, treatment and case studies, *J. Cleaner Prod.*, 14 (12-13), 1139–1145.
- [11] Tan, G.L., Shu, W.S., Hallberg, K.B., Li, F., Lan, C.Y., and Huang, L.N., 2007, Cultivation-dependent and cultivation-independent characterization of the microbial community in acid mine drainage associated with acidic Pb/Zn mine tailings at Lechang, Guangdong, China, *FEMS Microbiol. Ecol.*, 59 (1), 118–126.
- [12] Campaner, V.P., Luiz-Silva, W., and Machado, W., 2014, Geochemistry of acid mine drainage from a coal mining area and processes controlling metal attenuation in stream waters, Southern Brazil, *An. Acad. Bras. Ciênc.*, 86 (2), 539–554.
- [13] Bardgett, R.D., Bowman, W.D., Kaufmann, R., and Schmidt, S.K., 2005, A temporal approach to linking aboveground and belowground ecology, *Trends Ecol. Evol.*, 20 (11), 634–641.
- [14] Ahmad, A.K., and Al-Mahaqeri, S.A., 2015, Human health risk assessment of heavy metals in fish species collected from catchments of former tin mining, *Int. J. Res. Stud. Sci. Eng. Technol.*, 2 (4), 9–21.
- [15] Zeng, F., Ali, S., Zhang, H., Ouyang, Y., Qiu, B., Wu, F., and Zhang, G., 2011, The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants, *Environ. Pollut.*, 159 (1), 84–91.
- [16] Huang, L.N., Tang, F.Z., Song, Y.S., Wan, C.Y., Wang, S.L., Liu, W.Q., and Shu, W.S., 2010, Biodiversity, abundance, and activity of nitrogen-fixing bacteria during primary succession on a copper mine tailings, *FEMS Microbiol. Ecol.*, 78 (3), 439–450.
- [17] Abdel-Raouf, N., Al-Homaidan, A.A., and Ibraheem, I.B.M., 2012, Microalgae and wastewater treatment, *Saudi J. Biol. Sci.*, 19 (3), 257–275.
- [18] Tschirko, D., Hammesfahr, U., Zeltner, G., Kandeler, E., and Böcker, R., 2005, Plant succession and rhizosphere microbial communities in a recently deglaciated alpine terrain, *Basic Appl. Ecol.*, 6 (4), 367–383.
- [19] Fierer, N., Nemergut, D., Knight, R., and Craine, J.M., 2010, Changes through time: Integrating

- microorganisms into the study of succession, *Res. Microbiol.*, 161 (8), 635–642.
- [20] Quirós, R., 2003, The relationship between nitrate and ammonia concentrations in the pelagic zone of lakes, *Limnetica*, 22 (1-2), 37–50.
- [21] Michalski, R., and Kurzyca, I., 2006, Determination of nitrogen species (nitrate, nitrite and ammonia ions) in environmental samples by ion chromatography, *Pol. J. Environ. Stud.*, 15 (1), 5–18.
- [22] Mihale, M.J., 2015, Nitrogen and phosphorus dynamics in the waters of the Great Ruaha River, Tanzania, *J. Water Resour. Ocean Sci.*, 4 (5), 59–71.
- [23] Akan, J.C., Audu, S.I., Mohammed, Z., and Ogugbuaja, V.O., 2013, Assessment of heavy metals, pH, organic matter and organic carbon in roadside soils in Makurdi Metropolis, Benue State, Nigeria, *J. Environ. Prot.*, 4 (6), 618–628.
- [24] Kuriata-Potasznik, A., Szymczyk, S., Skwierawski, A., Glińska-Lewczuk, K., and Cymes, I., 2016, Heavy metal contamination in the surface layer of bottom sediments in a flow-through lake: A case study of Lake Symsar in Northern Poland, *Water*, 8 (8), 358.
- [25] Maher, W., Krikowa, F., Wruck, D., Louie, H., Nguyen, T., and Huang, W.Y., 2002, Determination of total phosphorus and nitrogen in turbid waters by oxidation with alkaline potassium peroxodisulfate and low pressure microwave digestion, autoclave heating or the use of closed vessels in a hot water bath: Comparison with Kjeldahl digestion, *Anal. Chim. Acta*, 463 (2), 283–293.
- [26] Song, Y., Hahn, H.H., and Hoffmann, E., 2002, Effects of solution conditions on the precipitation of phosphate for recovery a thermodynamic evaluation, *Chemosphere*, 48 (10), 1029–1034.
- [27] Kumar, M.M.S.Y., Galil, M.S.A., Suresha, M.S., Sathish, M.A., and Nagendrappa, G., 2007, A simple spectrophotometric determination of phosphate in sugarcane juices, water, and detergent samples, *E-J. Chem.*, 4 (4), 467–473.
- [28] Topçu, A., and Pulatsü, S., 2014, Phosphorus fractions and cycling in the sediment of a shallow eutrophic pond, *J. Agric. Sci.*, 20 (1), 63–70.
- [29] Sengupta, S., Nawaz, T., and Beaudry, J., 2015, Nitrogen and phosphorus recovery from wastewater, *Curr. Pollut. Rep.*, 1 (3), 155–166.
- [30] Cerozi, B.S., and Fitzsimmons, K., 2016, The effect of pH on phosphorus availability and speciation in an aquaponics nutrient solution, *Bioresour. Technol.*, 219, 778–781.
- [31] Lei, Y., Song, B., van der Weijden, R.D., Saakes, M., and Buisman, C.J.N., 2017, Electrochemical induced calcium phosphate precipitation: importance of local pH, *Environ. Sci. Technol.*, 51 (19), 11156–11164.
- [32] De Saedeleer, V., Cappuyns, V., De Cooman, W., and Swennen, R., 2010, Influence of major elements on heavy metal composition of river sediments, *Geol. Belg.*, 13 (3), 257–268.
- [33] Jiang, M., Zeng, G., Zhang, C., Ma, X., Chen, M., Zhang, J., Lu, L., Yu, Q., Hu, L., and Liu, L., 2012, Assessment of heavy metal contamination in the surrounding soils and surface sediments in Xiawangang River, Qingshuitang District, *PLoS One*, 8 (8), e71176.
- [34] Alamgir, M., Islam, M., Hossain, N., Kibria, M.G., and Rahman, M.M., 2015, Assessment of heavy metal contamination in urban soils of Chittagong City, Bangladesh, *Int. J. Plant Soil Sci.*, 7 (6), 362–372.
- [35] Zhang, C., Yu, Z., Zeng, G., Jiang, M., Yang, Z., Cui, F., Zhu, M., Shen, L., and Hu, L., 2014, Effects of sediment geochemical properties on heavy metal bioavailability, *Environ. Int.*, 73, 270–281.
- [36] Yao, Q., Wang, X., Jian, H., Chen, H., and Yu, Z., 2015, Characterization of the particles size fraction associated with heavy metals in suspended sediments of the Yellow River, *Int. J. Environ. Res. Public Health*, 12 (6), 6725–6744.

Short Communication:**The Pattern of Heavy Metals Distribution in Time Chronosequence of Ex-Tin Mining Ponds in Bangka Regency, Indonesia**Andri Kurniawan^{1,2,*}, Oedjijono², Tamad³, and Uyi Sulaeman⁴¹Department of Aquaculture, University of Bangka Belitung,
Jl. Balunijuk, Merawang, Bangka, Bangka Belitung 33172, Indonesia²Faculty of Biology, Jenderal Soedirman University,
Jl. Dr. Suparno no. 63, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia³Department of Agriculture, Jenderal Soedirman University,
Jl. Dr. Suparno no. 61, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia⁴Department of Chemistry, Faculty of Science and Technology, Jenderal Soedirman University,
Jl. Dr. Suparno no. 61, Karang Bawang, Grendeng, Purwokerto 53122, Central Java, Indonesia*** Corresponding author:**

tel: +62-81351714747

email: andri-kurniawan@ubb.ac.id

Received: March 1, 2018

Accepted: July 17, 2018

DOI: 10.22146/ijc.33613

Abstract: The heavy metals distribution of ex-tin mining ponds were investigated. The time chronosequence was determined at the pond of age < 1 year (Station A), the pond of age 5–10 years (Station B), and the pond of age > 15 years (Station C). The results showed sixteen heavy metals of As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V could be detected in the ponds. The metals such as As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn in Station C showed higher concentration compared to the Station A and Station B. The metals such as Cr, Fe, Hf, Sn, Ta, Te, and V in Station A and Station B showed higher concentration compared to the Station C. The positive, negative, and dynamic correlation pattern could be found in distribution of heavy metal to time chronosequence. The concentration of Ta and V showed a positive correlation because their concentration decrease, whereas concentration of As, Cu, Ga, Mn, and Zn showed a negative correlation because their concentration increase along in time chronosequence. The dynamic correlation could be found that concentration of Co, Ni, Pb, Sn, and Th decrease from Station A to Station B and then increase in Station C, whereas concentration of Cr, Fe, Hf, and Te increase from Station A to Station B and then decrease in Station C.

Keywords: distribution pattern; heavy metals; chronosequence; ex-tin mining ponds

■ INTRODUCTION

Tin mining activity is an anthropogenic activity that has contributed to the ecological problems. It has been known as the major sources of heavy metal contamination such as heavy metals in ex-tin mining ponds. Ex-tin mining ponds as one of water resource were contaminated by heavy metals that can be accumulated in food chains and secondary activities of the water [1-3]. Some of the heavy metals such as Pb, Zn, Mn, Fe, Cr, Cu, Ni, Sn, As, and Cd were detected in site of ex-tin mining [4-5].

The each of waters had their pattern of physical and chemical characteristics that determined by climatic, geomorphological, and geochemical conditions [6]. These conditions were influenced by ecological succession in time chronosequence [7] and the ecological changes of ex-tin mining ponds have occurred for a long time [8]. The implication of ecological changes as long as time chronosequence was interactions and reactions of chemical complexation between organic compounds, environmental factors, and inorganic compounds, include presence of heavy metals [9].

The researches of heavy metals in ex-tin mining ecosystems have been done, however, not for correspondence of heavy metals distributions and ex-tin mining ponds in time chronosequence. While an understanding this correspondence was crucial to predict ecosystems conditions and to detect environment change. The understanding of heavy metals distribution in time chronosequence of ex-tin mining ponds becomes an important part of aquatic ecosystem management as water sources for secondary activities and habitable for organisms.

■ EXPERIMENTAL SECTION

Materials

The study areas were located in Bangka Regency, Bangka Belitung Archipelago Province of Indonesia. The considered study sites cover three ponds of ex-tin mining with chronosequence ranging in time. The study areas were encoded as Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years).

The coordinate of Station A were 01°59' S in points 36.0"; 36.2"; 36.4"; 36.5"; 36.6" and 106°06' E in points 36.5"; 36.9"; 37.3"; 37.4"; 37.5". The coordinate of Station B were 01°59' S in points 41.3"; 41.4"; 41.5"; 42.4"; 42.5" and 106°06' E in points 39.2"; 39.5"; 41.4"; 42.7"; 43.1". The coordinate of Station C were 01°55' S in points 40.9"; 58.9"; 59.1"; 59.2"; 59.5" and 106°06' E in points 19.5"; 19.7"; 19.9"; 22.4"; 29.2".

In the each of research stations points, water sampling was done < 4 m in depth (station code A.1; B.1; and C.1) and composite sampling were done to water and sediment > 4 m in depth (station code A.2; B.2; and C.2).

Instrumentation

The water samples and composite samples were analyzed pH value by pH meter (PH-009(I)-A) Sun Care. The speciation and concentration of heavy metals analyzed by X-Ray Fluorescence (Rigaku) with three light spreader metals of copper (Cu), molybdenum (Mo), and aluminum (Al). Also, the pH and moisture of soil around the ponds were measured by pH-moisture meter. The correspondence between heavy metals distribution and

the time chronosequence of ex-tin mining ponds was analyzed by PAST (*Palaentological Statistics*) 3.

■ RESULTS AND DISCUSSION

pH Value of Water and Soil

Tin mining activity had generated the ex-tin mining ponds as one of the water resources. However, water quality in the ponds had serious ecological problem. The ex-mining activity produced acid mine drainage (AMD) that can caused the long-term impairment of water quality and biodiversity. The pH value of ex-tin mining water and soil pH value indicated that the acid could be found in Station A (pond in age < 1 year) and Station B (pond in age 5–10 years) and the neutral value could be found in Station C (pond in age > 15 years). In the each of research stations points, water samples < 4m in depth indicated more acid than the composite samples (water and sediment > 4 m in depth) (Fig. 1).

The acid mine drainage (AMD) was produced when sulfide-bearing material is exposed to oxygen and water [10]. In acid mine drainage, presence and mobility of elements such as Al, As, Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Zn, and also ionic activity and sulphide




Fig 1. Soil and water pH value in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth)

minerals oxidation caused pH value decreasing [11-12]. Furthermore, the low pH condition implicated in oxidation state, toxicity, and mobility of elements [9].

In long time chronosequence more than 15 years in the Station C caused changes water quality of ex-tin mining ponds were more stable than Station A and Station B. The indicator for it was pH value increased to neutral conditions (7.0-7.4) and other parameters that indicated water quality was more habitable for organisms' life such as fishes. The pH value and other parameters changes, indirectly, caused ecological productivity increasing. The chronosequence in long time caused changes in nutrient availability and resource supply for ecosystem productivity [13].

The pH value was found to play the most important role in the determination of metal speciation, solubility from mineral surfaces, movement, and bioavailability of metals. Furthermore, acid mine drainage had a major contribution in heavy metal contamination [14]. There was a negative correlation between soil pH and

distribution heavy metal where pH value decreasing caused increasing in heavy metal desorption [15].

The Pattern of Heavy Metals Distribution

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and neutral condition (in Station C) (Table 1).

There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. The average of As concentration were 1.7475 ppm (Station C) > 1.6075 ppm (Station B) > 1.0775 ppm (Station A). The concentration of Co were 3.32 ppm (Station C) > 2.525 ppm (Station A) > not detected (Station B). The concentration of Cu were 6.375 ppm (Station C) > 5.4825 ppm (Station B) > 4.9775 ppm (Station A).

Table 1. Distribution of heavy metals in time chronosequence in water of ex-tin mining ponds

No	Heavy Metals	Heavy Metals Concentration (ppm)					
		In Station A		In Station B		In Station C	
		Station A.1	Station A.2	Station B.1	Station B.2	Station C.1	Station C.2
1	As	nd	2.155	nd	3.215	2.4	1.095
2	Ga	nd	4.275	nd	4.47	9.76	nd
3	V	nd	0.84	nd	nd	nd	0.67
4	Te	nd	nd	102	nd	nd	5.8
5	Cr	nd	4.735	nd	4.945	nd	0.67
6	Zn	nd	6.05	nd	8.1	8.17	3.905
7	Sn	47	82.8	50.3	51.4	61.8	57.15
8	Mn	nd	13.05	nd	13.6	30.8	14.85
9	Fe	21.4	1481.3	148	2344	1590	849.05
10	Co	nd	5.05	nd	nd	6.64	nd
11	Ni	nd	4.09	nd	2.875	3.48	1.905
12	Cu	3.08	6.875	3.19	7.775	8.94	3.81
13	Pb	nd	6.7	nd	6	8.21	4.62
14	Th	nd	4.605	nd	4.38	13.5	7.05
15	Hf	6.71	3.21	8.21	3.32	nd	9.38
16	Ta	4.94	2.615	4.16	2.995	nd	2.805

*) nd (not detected). The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth)

The concentration of Ga were 4.88 ppm (Station C) > 2.235 ppm (Station B) > 2.1375 ppm (Station A). The concentration of Mn were 22.825 ppm (Station C) > 6.8 ppm (Station B) > 6.525 ppm (Station A). The concentration of Ni were 2.6925 ppm (Station C) > 2.045 ppm (Station A) > 1.4375 ppm (Station B). The concentration of Pb were 6.415 ppm (Station C) > 3.35 ppm (Station A) > 3.0 ppm (Station B). The concentration of Th were 10.275 ppm (Station C) > 2.3025 ppm (Station A) > 2.19 ppm (Station B). The concentration of Zn were 6.0375 ppm (Station C) > 4.05 ppm (Station B) > 3.025 ppm (Station A).

In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V. The average of Cr concentration were 2.4725 ppm (Station B) > 2.3675 ppm (Station A) > 0.335 ppm (Station C). The concentration of Fe were 1246 ppm (Station B) > 1219.53 ppm (Station C) > 751.35 ppm (Station A). The concentration of Hf were 5.765 ppm (Station B) > 4.96 ppm

(Station A) > 4.69 ppm (Station C). The concentration of Sn were 64.9 ppm (Station A) > 59.475 ppm (Station C) > 50.85 ppm (Station B). The concentration of Ta were 3.7775 ppm (Station A) > 3.5775 ppm (Station B) > 1.4025 ppm (Station C). The concentration of Te were 51 ppm (Station B) > 2.9 ppm (Station C) > not detected (Station A). The concentration of V were 0.42 ppm (Station A) > 0.335 ppm (Station C) > not detected (Station B).

Their concentration of heavy metals in time chronosequence in ex-tin mining ponds showed three patterns distribution that a positive correlation, a negative correlation, and a dynamic correlation. The positive correlation indicated that concentration of heavy metals decreased along chronosequence in time in ex-tin mining ponds. The negative correlation indicated that concentration of heavy metals increased along chronosequence in time in ex-tin mining ponds. The dynamic correlation indicated that concentration of




Fig 2. Distribution of heavy metals in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth)

heavy metals on decreased or increased along chronosequence in time in ex-tin mining ponds.

The elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. The elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increasing along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then increased concentration in ponds with age > 15 years. The elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then decreased concentration in ponds with age > 15 years (Fig. 2).

The chronosequence in long time caused changes in heavy metals concentration were decreased as the primary succession progression [16]. The low pH value implicated to dissolved oxygen (DO) [17] and low DO in acid mine drainage implicated to biological oxygen demand (BOD) by microorganisms for their growth and other ecological factor changes such as nitrogen and phosphate [7,18-19]. Nitrogen (ammonia, nitrite, and nitrate) and phosphate were important for sustenance of various life forms,

primary succession, and led to changes in the biogeochemical cycles of nutrients in ponds [20-22], indirectly implicated to dissolved and suspended solid [23-24] and ecosystem eutrophic levels [25-31]. Therefore, pH value changes, availability of elements included heavy metals, environmental characteristics, and eutrophication had a positive inter-relationships in their correspondence.

These conditions, however, had a negative correlation to heavy metals residue decreasing in Station C. The heavy metals in Station C were dominated As, Zn, Mn, Co, Ni, Cu, Pb, and Th were higher than Station A and Station B, although these metals and include Ga, Cr, Sn, Fe, Hf, and Ta were detected in all of the research station. The results indicated that chronosequence in time of ponds caused ecological changes and had positive correlations, but not for distribution of heavy metals.

The Correspondence Analysis of Heavy Metals Distribution

The correspondence analysis also showed that there was not a positive correlation between in time chronosequence in water of ex-tin mining ponds and




Fig 3. Correspondence analysis between distribution of heavy metals and in time chronosequence of ex-tin mining ponds in Bangka Regency of Indonesia. The Station A (pond in age < 1 year), Station B (pond in age 5–10 years), and Station C (pond in age > 15 years). The station code (A.1, B.1, and C.1) showed water samples < 4 m in depth and the station code (A.2, B.2, and C.2) showed composite samples (water and sediment > 4 m in depth)

heavy metals distribution (Fig. 3). The ecological changes in chronosequence succession of ex-tin mining ponds had not significantly affect on speciation and concentration of heavy metals. Thus, there was the other factor that had affected the distribution of heavy metals.

The soil and sediment of water contributed to the distribution of heavy metals [32]. In general, the mobility and availability of heavy metals were controlled by adsorption and desorption characteristics of soils, include cation exchange capacity and the contents of clay minerals [15]. The clay and mud were found to be dominant in the sediment of Station A and Station B, however, sand was found to be dominant in Station C.

The clay and mud had a positive correlation and sand had a negative correlation to a presence, affinity, and minerals structure [33-34]. The clay and mud had a larger surface area, higher cation exchange capability (CEC), and particles of clay in sediments were often closely associated with organic matter and also exist in the form of organic-mineral complexes. These particles adsorbed the active metals from the water phases, accumulated, and carried them to the bottom sediments diagenetically [35-36]. Therefore, distribution of heavy metals in Station A and Station B were lower than Station C.

■ CONCLUSION

In time chronosequence of ex-tin mining ponds had affected ecological dynamic such as pH value change and heavy metals distribution. The pH value of ex-tin mining ponds moved from acid to neutral condition, however, it had been taken for a long time of the chronosequence. The consequence of low pH value can implicated to microorganisms' growth, nutrients, and environment factors in primary succession. Further, it implicated to heavy metals interaction, whereas the heavy metals residue should be in low concentration in long time chronosequence of ex-tin mining ponds.

The heavy metals that detected in ex-tin mining ponds were 16 element that As, Co, Cu, Cr, Fe, Ga, Hf, Sn, Ta, Te, Th, Mn, Ni, Pb, Zn, and V. All of these heavy metals were distributed in almost of research stations. Distribution of heavy metals can be grouped in low pH value or acid condition (in Station A and Station B) and

neutral condition (in Station C). There were nine heavy metals with higher concentration were found in Station C than Station A and Station B that As, Co, Cu, Ga, Mn, Ni, Pb, Th, and Zn. In addition, there were seven heavy metals with higher concentration in Station A or Station B than Station C that Cr, Fe, Hf, Sn, Ta, Te, and V.

There were three patterns of heavy metals distribution in time chronosequence in ex-tin mining ponds that a positive correlation, a negative correlation, and a dynamic correlation. The chronosequence in time had affects on heavy metals distribution. Elements of Ta and V had a positive correlation in time chronosequence in ex-mining ponds with their concentration on decreasing. Elements of As, Cu, Ga, Mn, and Zn had a negative correlation with their concentration on increased along in time chronosequence. In addition, Co, Ni, Pb, Sn, and Th had decreased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then increased concentration in ponds with age > 15 years. Elements of Cr, Fe, Hf, and Te had increased concentration from ponds with age < 1 year to ponds with age between 5–10 years and then decreased concentration in ponds with age > 15 years.

The chronosequence in time can be a factor that contributed to heavy metals distribution. However, the sediment factor had a significant effect to distribution of heavy metals in the ex-tin mining ponds, where clay and mud were more effective than sand in accumulation, adsorption, and interaction with heavy metals and carried them diagenetically to bottom sediments.

■ ACKNOWLEDGMENTS

I would like to thank Ministry of Research, Technology and Higher Education of the Republic of Indonesia for scholarship of Pendidikan Pascasarjana Dalam Negeri (BPPDN) and research grant of Program Disertasi Doktor (PDD) for the funding with Project No: 120.G/UN50.3.1/PP/2018.

■ REFERENCES

- [1] de Silva, A.A.L., de Carvalho, M.A.R., de Souza, S.A.L., Dias, P.M.T., Filho, R.G.S., Saramago, C.S.M., Bento, C.A.M., and Hofer, E., 2012, Heavy

- metal tolerance (Cr, Ag and Hg) in bacteria isolated from sewage, *Braz. J. Microbiol.*, 43 (4), 1620–1631.
- [2] Guan, Y., Shao, C., and Ju, M., 2014, Heavy metal contamination assessment and partition for industrial and mining gathering areas, *Int. J. Environ. Res. Public Health*, 11 (7), 7286–7303.
- [3] Kurniawan, A., 2017, Chronosequence effect of post tin mining ponds to metals residu and micro ecosystem change, *Omni-Akuatika*, 13 (1), 60–65.
- [4] Ashraf, M.A., Maah, M.J., and Yusoff, I., 2012, Speciation of heavy metals in the sediments of former tin mining catchment, *Iran. J. Sci. Technol. Trans. A Sci.*, 36 (2), 163–180.
- [5] Daniel, V.N., Chudusu, E.S., Chup, J.A., and Pius, N.D., 2014, Variations of heavy metals in agricultural soils irrigated with tin water in Heipang District of Barkin Ladi, Plateau State, Nigeria, *Int. J. Sci. Technol.*, 3 (5), 255–263.
- [6] Ashraf, M.A., Maah, M.J., and Yusoff, I., 2012, Morphology, geology and water quality assessment of former tin mining catchment, *Sci. World J.*, 2012, 369206.
- [7] Urbanová, M., Kopecký, J., Valášková, V., Ságová-Marečková, M., Elhottová, D., Kyselková, M., Moënné-Loccoz, Y., and Baldrian, P., 2011, Development of bacterial community during spontaneous succession on spoil heaps after brown coal mining, *FEMS Microbiol. Ecol.*, 78 (1), 59–69.
- [8] Kurniawan, A., 2016, Microorganism communities response of ecological changes in post tin mining ponds, *RRJoMV*, 6 (1), 17–26.
- [9] Violante, A., Cozzolino, V., Perelomov, L., Caporale, A.G., and Pigna, M., 2010, Mobility and bioavailability of heavy metals and metalloids in soil environments, *J. Soil Sci. Plant Nutr.*, 10 (3), 268–292.
- [10] Akcil, A., and Koldas, S., 2006, Acid mine drainage (AMD): causes, treatment and case studies, *J. Cleaner Prod.*, 14 (12-13), 1139–1145.
- [11] Tan, G.L., Shu, W.S., Hallberg, K.B., Li, F., Lan, C.Y., and Huang, L.N., 2007, Cultivation-dependent and cultivation-independent characterization of the microbial community in acid mine drainage associated with acidic Pb/Zn mine tailings at Lechang, Guangdong, China, *FEMS Microbiol. Ecol.*, 59 (1), 118–126.
- [12] Campaner, V.P., Luiz-Silva, W., and Machado, W., 2014, Geochemistry of acid mine drainage from a coal mining area and processes controlling metal attenuation in stream waters, Southern Brazil, *An. Acad. Bras. Ciênc.*, 86 (2), 539–554.
- [13] Bardgett, R.D., Bowman, W.D., Kaufmann, R., and Schmidt, S.K., 2005, A temporal approach to linking aboveground and belowground ecology, *Trends Ecol. Evol.*, 20 (11), 634–641.
- [14] Ahmad, A.K., and Al-Mahaqeri, S.A., 2015, Human health risk assessment of heavy metals in fish species collected from catchments of former tin mining, *Int. J. Res. Stud. Sci. Eng. Technol.*, 2 (4), 9–21.
- [15] Zeng, F., Ali, S., Zhang, H., Ouyang, Y., Qiu, B., Wu, F., and Zhang, G., 2011, The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants, *Environ. Pollut.*, 159 (1), 84–91.
- [16] Huang, L.N., Tang, F.Z., Song, Y.S., Wan, C.Y., Wang, S.L., Liu, W.Q., and Shu, W.S., 2010, Biodiversity, abundance, and activity of nitrogen-fixing bacteria during primary succession on a copper mine tailings, *FEMS Microbiol. Ecol.*, 78 (3), 439–450.
- [17] Abdel-Raouf, N., Al-Homaidan, A.A., and Ibraheem, I.B.M., 2012, Microalgae and wastewater treatment, *Saudi J. Biol. Sci.*, 19 (3), 257–275.
- [18] Tschërko, D., Hammesfahr, U., Zeltner, G., Kandeler, E., and Böcker, R., 2005, Plant succession and rhizosphere microbial communities in a recently deglaciated alpine terrain, *Basic Appl. Ecol.*, 6 (4), 367–383.
- [19] Fierer, N., Nemergut, D., Knight, R., and Craine, J.M., 2010, Changes through time: Integrating microorganisms into the study of succession, *Res. Microbiol.*, 161 (8), 635–642.
- [20] Quirós, R., 2003, The relationship between nitrate and ammonia concentrations in the pelagic zone of lakes, *Limnetica*, 22 (1-2), 37–50.
- [21] Michalski, R., and Kurzyca, I., 2006, Determination of nitrogen species (nitrate, nitrite and ammonia

- ions) in environmental samples by ion chromatography, *Pol. J. Environ. Stud.*, 15 (1), 5–18.
- [22] Mihale, M.J., 2015, Nitrogen and phosphorus dynamics in the waters of the Great Ruaha River, Tanzania, *J. Water Resour. Ocean Sci.*, 4 (5), 59–71.
- [23] Akan, J.C., Audu, S.I., Mohammed, Z., and Ogugbuaja, V.O., 2013, Assessment of heavy metals, pH, organic matter and organic carbon in roadside soils in Makurdi Metropolis, Benue State, Nigeria, *J. Environ. Prot.*, 4 (6), 618–628.
- [24] Kuriata-Potasznik, A., Szymczyk, S., Skwierawski, A., Glińska-Lewczuk, K., and Cymes, I., 2016, Heavy metal contamination in the surface layer of bottom sediments in a flow-through lake: A case study of Lake Symsar in Northern Poland, *Water*, 8 (8), 358.
- [25] Maher, W., Krikowa, F., Wruck, D., Louie, H., Nguyen, T., and Huang, W.Y., 2002, Determination of total phosphorus and nitrogen in turbid waters by oxidation with alkaline potassium peroxodisulfate and low pressure microwave digestion, autoclave heating or the use of closed vessels in a hot water bath: Comparison with Kjeldahl digestion, *Anal. Chim. Acta*, 463 (2), 283–293.
- [26] Song, Y., Hahn, H.H., and Hoffmann, E., 2002, Effects of solution conditions on the precipitation of phosphate for recovery a thermodynamic evaluation, *Chemosphere*, 48 (10), 1029–1034.
- [27] Kumar, M.M.S.Y., Galil, M.S.A., Suresha, M.S., Sathish, M.A., and Nagendrappa, G., 2007, A simple spectrophotometric determination of phosphate in sugarcane juices, water, and detergent samples, *E-J. Chem.*, 4 (4), 467–473.
- [28] Topçu, A., and Pulatsü, S., 2014, Phosphorus fractions and cycling in the sediment of a shallow eutrophic pond, *J. Agric. Sci.*, 20 (1), 63–70.
- [29] Sengupta, S., Nawaz, T., and Beaudry, J., 2015, Nitrogen and phosphorus recovery from wastewater, *Curr. Pollut. Rep.*, 1 (3), 155–166.
- [30] Cerozi, B.S., and Fitzsimmons, K., 2016, The effect of pH on phosphorus availability and speciation in an aquaponics nutrient solution, *Bioresour. Technol.*, 219, 778–781.
- [31] Lei, Y., Song, B., van der Weijden, R.D., Saakes, M., and Buisman, C.J.N., 2017, Electrochemical induced calcium phosphate precipitation: importance of local pH, *Environ. Sci. Technol.*, 51 (19), 11156–11164.
- [32] De Saedeleer, V., Cappuyns, V., De Cooman, W., and Swennen, R., 2010, Influence of major elements on heavy metal composition of river sediments, *Geol. Belg.*, 13 (3), 257–268.
- [33] Jiang, M., Zeng, G., Zhang, C., Ma, X., Chen, M., Zhang, J., Lu, L., Yu, Q., Hu, L., and Liu, L., 2012, Assessment of heavy metal contamination in the surrounding soils and surface sediments in Xiawangang River, Qingshuitang District, *PLoS One*, 8 (8), e71176.
- [34] Alamgir, M., Islam, M., Hossain, N., Kibria, M.G., and Rahman, M.M., 2015, Assessment of heavy metal contamination in urban soils of Chittagong City, Bangladesh, *Int. J. Plant Soil Sci.*, 7 (6), 362–372.
- [35] Zhang, C., Yu, Z., Zeng, G., Jiang, M., Yang, Z., Cui, F., Zhu, M., Shen, L., and Hu, L., 2014, Effects of sediment geochemical properties on heavy metal bioavailability, *Environ. Int.*, 73, 270–281.
- [36] Yao, Q., Wang, X., Jian, H., Chen, H., and Yu, Z., 2015, Characterization of the particles size fraction associated with heavy metals in suspended sediments of the Yellow River, *Int. J. Environ. Res. Public Health*, 12 (6), 6725–6744.