

EEC

by Ardiansyah Kurniawan



Submission date: 11-Feb-2023 09:31AM (UTC+0700)

Submission ID: 2011353176

File name: EEC_27_2_2021_Ardiansyah_Kurniawan.pdf (172.26K)

Word count: 1328

Character count: 7408

EEC

by Ardiansyah Kurniawan

Submission date: 11-Feb-2023 09:31AM (UTC+0700)

Submission ID: 2011353176

File name: EEC_27_2_2021_Ardiansyah_Kurniawan.pdf (172.26K)

Word count: 1328

Character count: 7408

¹ A New Record of *Osteochilus spilurus* Bleeker, 1851 (Teleostei, Cyprinidae) in Bangka and Belitung Island, Indonesia

Ardiansyah Kurniawan^{1,2*}, Anik M. Hariati³, Andi Kurniawan³ and Dewa G.R. Wiadnya³

²
¹Post-Graduate Program, Faculty of Fisheries, and Marine Science, Universitas Brawijaya,
Malang, Indonesia.

²Department of Aquaculture, Universitas Bangka Belitung, Bangka, Indonesia

³Faculty of Fisheries, and Marine Science, Universitas Brawijaya, Malang, Indonesia

⁵
(Received 15 September, 2020; accepted 11 November, 2020)

ABSTRACT

¹
Osteochilus spilurus is native freshwater fish in Sundaland. Previous records of this species in Indonesia indicate a dispersed gap between The Sumatera and the Kalimantan Islands, namely Bangka Belitung Archipelago. A new record in Bangka and the Belitung Islands completes its distribution in Southeast Asia. Fishes with *O. spilurus* morphological characters have been found in five separate regencies.

Keywords: Sundaland, Fish distribution, Southeast Asia

Introduction

Land interactions allow freshwater fish to be dispersed naturally in Indonesia during the Pleistocene. Freshwater fish zoogeography in Southeast Asia reveals that the Sumatra-Kalimantan-Java Islands area has the same native species as part of the Sundaland (Zakaria, 1994). Of these, *Osteochilus spilurus* is present in sluggish freshwater rivers (Kottelat *et al.*, 1993; Haryono and Wahyudewantoro, 2020). The position of lesser concern on it (Huckstorf, 2012) induces a limited distribution of human factors.

In Bangka, Belitung, and Java Islands, there is no record of distribution until 2019 (GBIF, 2019). The last statement on the eastern island of Belitung varies from previous studies on economic benefits, capturing and eating this fish (Kurniawan *et al.*, 2020), and the rise of artificial habitat (Kurniawan *et al.*, 2019). Its natural distribution on the Bangka and

Belitung Islands is important to complete information on the distribution in Southeast Asia and to help its efforts to domesticate aquaculture commodities.

Materials and Methods

In this study, five separate regencies were selected for sampling sites in Bangka and Belitung Islands, Indonesia. Two locations are on Belitung Island, namely Lenggang River (2°55'09.5 "S 108°06'35.4" E) in East Belitung Regency and Air Tering River (3°02'38 "S, 107°40'13 "E) in Belitung regency. Another location is on Bangka Island, a small river of Bencah Village (2°46'03" S, 106°23'48" E) in South Bangka Regency, Lebak River (2°4'38" S, 105°51'53 "E) in Bangka regency, and Lelabi river (1°50'43.7" S, 105°44'57.6" E) in West Bangka Regency. Sampling was conducted during March 2019 until July 2020. This fish does not require special permits and

collected from natural catches using fish traps for wide rivers and scoops for small rivers. Specimens were observed by Weber and de Beaufort (1916) key morphological characteristics of *Osteochilus spilurus* to identify the species.

Results

We found fishes from the five sampling sites in the Bangka and Belitung Islands have been described as *Osteochilus spilurus*. They have a specific morphological character of Weber and de Beaufort (1916) describe, follows: compressed body, snout pointed, dorsal height gradually rising before dorsal fin, prominent with numerous small pores, long maxillary barbels, 28-30 scales lateral line, dorsal fin rays 12-13 and start from on the 7th or 8th scale of lateral line, caudal fin emarginate shape, and a large black blotch on the caudal peduncle. Twenty individuals of each sampling site are labeled and immersed in a 5% formalin solution and then placed in Brawijaya Ichthyologicum Depository, Malang. Standard lengths of them ranging from 61.04 to 22.44 mm.



Fig. 1. Specimen of *Osteochilus spilurus*.




Fig. 1. Distribution of *O. spilurus* in Sundaland (Southeast Asia). Black triangles are previous records before 2019 (GBIF, 2019). Black squares are the recent record in Bangka and Belitung Islands.

Discussion

O. spilurus presence on Bangka Island and Belitung Island is an essential supplementary record in Southeast Asia. Currently, Bangka Belitung Archipelago between Bangka Strait and Karimata Strait are not barriers to the spread of this species. These records solve the Sundaland distribution gap. Figure 2 indicates the more complete distribution of this species in Southeast Asia.

Geographically, the islands of Bangka and Belitung are included in the part of Sundaland during Pleistocene period. We estimate that rivers that were related spread spilurus in the past, including to Bangka and Belitung Islands. Historical events affect how ichthyofauna is distributed (Berendzen *et al.*, 2003). Ancient river is a historical barrier for ichthyofauna migration (Waters *et al.*, 2001; Mayden, 1988). Bangka and Belitung islands are related to the modern Indragiri, Hari, and Musi rivers on the island of Sumatra, Kapuas river in West Kalimantan on the ancient northern Sundanese rivers (Voris, 2000; Hutama *et al.*, 2016). However, it is also possible to become the upstream portion of the East Sunda Purba river that connects the southern part of the island of Sumatra, Java Island, and the south of the island of Kalimantan with the estuary in the Java Sea (Kusuma *et al.*, 2016). The lack of economic usage made the study of *O. spilurus* more precise in its distribution.

Acknowledgement

Our sincere gratitude goes out to LPDP (Indonesia

Endowment Fund for Education) for providing my study scholarship and supporting this research, Ms. Weni, Ms. Nicko, and Suheri for kind assistance in the field.

References

- Berendzen, P. B. Simons, A. M. and Wood, R. M. 2003. Phylogeography of the northern hogsucker, *Hypentelium nigricans* (Teleostei: Cypriniformes): genetic evidence for the existence of the ancient Teays River. *Journal of Biogeography*. 30(8): 1139-1152.
- GBIF. 2019. *Osteochilus spilurus* (Bleeker, 1851). GBIF Backbone Taxonomy. Checklist dataset <https://doi.org/10.15468/390mei> accessed via GBIF.org on 2020-09-15.
- Haryono, and Wahyudewantoro, G. 2020. The freshwater fishes and species status of peatland areas in Central Kalimantan, Indonesia. *Eco. Env. & Cons.* 26 (June Suppl. Issue). S14-S19
- Huckstorf, V. 2012. *Osteochilus spilurus*. The IUCN Red List of Threatened Species 2012: e.T181063A1694304. DOI: 10.2305/ iucn . u k . 2 0 1 2 - 1 . r l t s . t 1 8 1 0 6 3 a 1 6 9 4 3 0 4 . e n
- Hutama, A. A. Hadiaty, R.K. Hubert, N. 2016. Biogeography Of Indonesian Freshwater Fishes: Current Progress. *Treubia*. 43: 17-30
- Kottelat, M. Whitten, A.J. Kartikasari S.N. and Wirjoatmodjo. S. 1993. *Freshwater Fishes of Western Indonesia and Sulawesi*. Periplus Editions, Hong Kong.
- Kurniawan, A. Azhari, M. and Prasetyono, E. 2019. Domestication of *Osteochilus spilurus*: survival and growth in recirculated water. *IOP Conference Series: Earth and Environmental Science* 230 in International Conference on Green Agro-industry and Bioeconomy, East Java Indonesia. September 18-20, 2018. DOI:10.1088/1755-1315/230/1/012116
- Kurniawan, A., Hariati, A. M., Darmawan, A. and Wiadnya, D.G.R. 2020. Morphometric comparison of *Osteochilus spilurus* (Bleeker 1851) from Bangka and Belitung Island, Indonesia. In IOP Conference Series: Earth and Environmental Science 493 (1) in International Conference on Sustainable Aquatic Resources 27-28 August 2019, Malang, Indonesia. DOI: 10.1088/1755-1315/493/1/012024.
- Kusuma, W. E., Ratmuangkhwang, S. and Kumazawa, Y. 2016. Molecular phylogeny and historical biogeography of the Indonesian freshwater fish *Rasbora lateristriata* species complex (Actinopterygii: Cyprinidae): cryptic species and west-to-east divergences. *Molecular Phylogenetics and Evolution*. 105: 212-223.
- Mayden, R. L. 1988. Vicariance biogeography, parsimony, and evolution in North American freshwater fishes. *Systematic Biology*. 37(4): 329-355.
- Voris, H.K. 2000. Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. *Journal of Biogeography*. 27: 1153-1167.
- Waters, J. M. Craw, D. Youngson, J. H. and Wallis, G. P. 2001. Genes meet geology: fish phylogeographic pattern reflects ancient, rather than modern, drainage connections. *Evolution*. 55(9): 1844-1851.
- Weber, M. W. C. and de Beaufort, L. F. 1916. *The Fishes of the Indo-Australian Archipelago*. Volume 3. EJ Brill Limited.
- Zakaria, I. M. 1994. Zoogeography and biodiversity of the freshwater fishes of Southeast Asia. In: *Ecology and conservation of Southeast Asian marine and freshwater environments including wetlands*. 41-48. Springer, Dordrecht.

ORIGINALITY REPORT

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

5%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	isa.niscair.res.in Internet Source	8%
2	A Kurniawan, A M Hariati, A Kurniawan, A Darmawan, D G R Wiadnya. "Morphometric comparison of Osteochilus spilurus (Bleeker 1851) from Bangka and Belitung Island, Indonesia", IOP Conference Series: Earth and Environmental Science, 2020 Publication	3%
3	Submitted to Sriwijaya University Student Paper	1%
4	biodiversitas.mipa.uns.ac.id Internet Source	1%
5	link.springer.com Internet Source	1%

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On