



The contamination of heavy metals (Pb and Cu) at tin sea mining field and its impact to marine tourism and fisheries

by Sudirman Adibrata

Submission date: 12-Mar-2021 08:45AM (UTC+0700)

Submission ID: 1530756932

File name: 37219-112472-1-SM.doc (172.73K)

Word count: 4113

Character count: 22048

The contamination of heavy metals (Pb and Cu) at tin sea mining field and its impact to marine tourism and fisheries

Sudirman Adibrata^{1*}, Muh Yusuf², Irvani³, Mu'alimah Hudatwi⁴, Aditya Pamungkas⁴, Maulana Firdaus⁵

¹Aquatic Resources Management Department of Bangka Belitung University

²Oceanographic Department of Diponegoro University

³Mining Engineering Department of Bangka Belitung University

⁴Marine Science Department of Bangka Belitung University

⁵Department of Agribusiness Management, Graduate School of Agriculture, Tokyo University of Agriculture, Japan

*email: sadibrata@gmail.com

Abstract. This study aims to determine the extent of heavy metals (Pb and Cu) distribution adjacent sea mining field and its impact on marine tourism and fisheries. The purposive sampling method was used to identify the sampling locations from the 13 closest locations to the farthest from marine mining sources. The closest and farthest Pb and Cu values from the tin mining activities were 0.16 mg/l; 0.03 mg/l and 0.02 mg/l; <0.003 mg/l. The Hydro-oceanographic conditions from the highest to the lowest water currents of 0.03 m/ s - 0.001 m/ s. This research represented the east monsoon, which showed that the heavy metal distributions' dynamics are not too distance-reaching due to weak water currents, so that the dilution and sedimentation rate is slow. It is stated that the waters are polluted where the highest contamination value is indicated by the location closest to the source of marine mining and further weakened at the location farthest from the source of pollutants. This condition has significantly threatened marine tourism and fishery activities by reducing economic benefits so it is necessary to regulate tin mining activities and sites for the common purposes and prevent conflicts.

Key Words: distributions, environmental impact, heavy metal, mining, Pb and Cu

INTRODUCTION

Tin mining (Sn) activity is a local economic activity that is one of Bangka Belitung Island's primary sectors. Bangka Belitung island is an island located in the western Indonesia, known as a marine tourism site and most of the population in coastal areas depends on fisheries activities (Valeriani *et al.*, 2015; Bidayani *et al.*, 2019). In recent decades, tin mining activities continuously cause a decrease in water quality and marine biodiversity, and this could affect people's economic activities, including marine tourism and fisheries (Nurtjahya *et al.*, 2017; Yulianti 2019). Tin mining activities in the land to the coastal and marine areas are thought to cause ecological damage to both areas. Furthermore, the water's function will be reduced if the beach and the sea waste more and more. The exploitation of tin mining in coastal and ocean areas naturally leaves waste such as tailings runoff, oil, damage to seabed land structure, and sea noise. Wastes, including heavy metals in the waters, flow from the surrounding land to the sea. It will change the quality of seawater, thereby reducing the carrying capacity of these waters. Heavy metals in waters related to marine biota according to Ministry of Environmental Decree of 2004 No. 51 such as Mercury (Hg), Chromium (Cr), Arsenic (As), Cadmium (Cd), Copper (Cu), Lead (Pb), Zinc (Zn), and Nickel (Ni). The common heavy metals polluting waters in urban areas are Pb, Zn, Cu, Cd, and Co (Bibby and Webster-Brown, 2006; Cirik *et al.*, 2012). According to Thomann and Muller (1987), The United States for the Environmental Protection Agency (USEPA) has established a list of pollutants that are mostly wasted into the environment including Copper (Cu) and Lead (Pb). The increase in the content of heavy metal elements is one of the problems and water quality limitations.

The geochemical fraction of heavy metals are essential in biological processes in ecosystems. This fraction component is intended to be absorbed by creatures that live on life (Wang *et al.*, 2015), which can be accumulated in the body of marine biota directly through the food chain. These heavy metals also meet the safe limits of what represents an essential aquatic source for human food (Puttiwongrak *et al.*, 2019). The distribution of Pb and Zn dissolved may vary according to season and tidal conditions. The presence and effect of metals in the system are determined by several main factors such as heavy metals and the composition of soluble fractions and heavy metal particulates in the waters (Hamad *et al.*, 2012). Further, Najamuddin *et al* (2016) stated that the concentrations were influenced by season, with the Pb and Zn's sources derived mainly from rivers and natural sources. Therefore, waters that contain metal pollutants can spread in all directions and it is necessary to know their value.

The dissolved metal concentrations of Pb, Cd, Cu, and Zn in the Bangka Regency especially Kelabat Bay, are generally relatively low and safe for biota life. The concentration of Pb, Cu, and Zn in sediments is twice as high as the outer bay and is not influenced by factors season (Arifin, 2011). Previous research of the same type on Bangka Island needs to be done elsewhere to compare the impacts of mining activities. In terms of the importance of the problem for ecosystems and aquatic resources, it is very essential to know the dynamics of heavy metal pollutants' distribution due to mining activities at sea. This study aims to determine the dynamics of the distribution of heavy metals (Pb and Cu) waters at tin sea mining field and its impact for marine tourism and fisheries activities.

MATERIALS AND METHODS

Sampling Location

The study was conducted in April 2019 with the location in the waters around the coast of Tanah Merah Central Bangka Regency, the geographical location at 02°12'50" S and 106°13'00" E (Figure 1).




Figure 1. Sampling location

Methods

Simple random sampling method from the 13 closest locations to the farthest from marine mining sources with Pb and Cu parameters. Equipment for measuring hydro-oceanography and water quality, namely current meters, DO-meters, hand refractometer, salinometers, Secchi disks, ± 5 GT vessels equipped with GPS map 585 Garmin (measuring time, coordinates, temperature, and water depth), sample bottles to analyze content Lead (Pb) and copper (Cu) in waters.

Data analysis

The samples were taken then analyzed by the laboratory in Baristand Palembang. Data analysis with Indonesian Standard (SNI 6989.6.2009 for Cu and SNI 6989.8.2009 for Pb) for quality control (Baristand Laboratory, 2019). The results of the Pb and Cu analysis are then compared with the seawater quality standard for marine tourism and marine biota according to the Ministry of Environmental Decree of 2004 No. 51.

RESULTS

The results of the analysis are presented in Table 1.

Table 1. Heavy metals content from each station.

No	Station Survey	Pb (mg/l)	Cu (mg/l)	Quality Standards*)		Annotation	
				Pb	Cu	Pb	Cu
22 April 2019							
1	001	0,12	0,03	0,005a;0,008b	0,050a;0,008b	>ab	>ab
2	002	0,12	0,02			>ab	>ab
3	003	0,05	<0,003			>ab	<ab

4	004	0,11	<0,003	>ab	<ab
5	005	0,13	<0,003	>ab	<ab
6	006	0,14	<0,003	>ab	<ab
7	007	0,08	<0,003	>ab	<ab
8	008	0,07	<0,003	>ab	<ab
9	009	0,10	<0,003	>ab	<ab
10	010	0,16	<0,003	>ab	<ab
11	011	0,14	<0,003	>ab	<ab
12	012	0,16	<0,003	>ab	<ab
13	013	0,02	<0,003	>ab	<ab

Note: * Quality Standards based on Ministry of Environment Decree of 2004 No. 51 Appendix II and III (a = for marine tourism and b = for marine biota).

Information on the content of heavy metal elements (Pb and Cu) in the sediment at five stations (9-13) around the Tanah Merah coast, namely Pb with a range of 2.19 - 15.09 ppm and Cu with a range of 1.35 - 2.56 ppm (Ahmad, 2013).

Table 2. Hydro-oceanographic conditions on Tanah Merah Beach

No	Station Survey	Temperature (°C)	Salinity (‰)	Water current (m/s)	Water clarity (%)
1	001	30.4	29	0.15	56.0
2	002	30.6	28	0.23	35.6
3	003	30.2	26	0.23	35.9
4	004	30.2	27	0.23	28.9
5	005	29.0	28	0.15	42.6
6	006	30.2	27	0.18	37.0
7	007	30.7	26	0.09	35.1
8	008	30.3	30	0.16	65.8
9	009	29.6	29	0.09	43.4
10	010	29.7	30	0.08	81.4
11	011	29.7	30	0.12	51.4
12	012	29.9	29	0.07	40.3
13	013	29.2	29	0.12	52.4

Note: * Weak current speed (<0.25 m/s), normal (0.25-0.3 m/s), strong (> 0.3 m/s) (Adibrata *et al.*, 2013)

DISCUSSIONS

The utilization of small islands in Indonesia is prioritized for conservation, education and training, research and development, marine culture, tourism, fisheries and marine business and sustainable fishing industry, organic agriculture, animal husbandry, and/or national defense and security (Law of 2007 No. 27 juncto Law of 2014 No. 1). Coastal research and development can be carried out in marine areas where there is already another activity at that location, for example mining activities in the sea. Where is Bangka Island, which has abundant tin ore resources, has no zoning regulations. Therefore, the area from the coastline up to 12 miles to the sea has no legality of permanent land use. This is what supports the need to control regional governance so that the pace of development can be controlled and not degrade the environment, especially coastal areas. Offshore mining can reduce water quality and change the seabed, which can cause changes in biodiversity, including damage to coral reefs as fish habitat. Onshore mining activities can reduce biodiversity and cause flooding and infrastructure damage (Nurtjahya *et al.*, 2017). Of course, mining activities in the sea will reduce the quality of marine biodiversity.

Marine mining activities on the Tanah Merah coast (Figure 1) show community tin mines and the presence of suction boats. Current patterns strongly support the dynamics of the distribution of heavy metal pollutant content (Pb and Cu). The presence of heavy metals Pb, Cd, and Cu is strongly influenced by the sediment's organic matter content and grain size. Heavy metal content will increase with increasing organic matter in sediments (Maslukah, 2013). Relative abundance of trace metals in sediments was in the order of Zn > Cu > Ni > Cr > Co > Pb > Cd. Zn, Cu, and Ni (Ahmed *et al.*, 2017). Further, the heavy metal concentrations of the sediments were found to decrease in the sequence Fe>Ti>Mn>Zn>V>Cu>Cr>Ni (Syakti *et al.*, 2015). Where there is the lowest Pb data at stations 3, 7, 8, and 13 with respective values of 0.05; 0.08; 0.07; and 0.02 mg/l. At other stations, there is a Pb value of more than or equal to 0.10 mg/l. The Pb value at all stations is stated above the seawater quality standard for marine tourism and marine life. The amount of Pb content in the sample exceeds the National Standard threshold for the Pb content in seawater for marine biota of 0.008 mg/l and causes Pb pollution around the mining area. The disposal of waste water from mining activities

continues into the waters without the filtering process causes the pollution of Pb in the mining environment. Although the Cu value is not as high as the Pb value at the same station, one of the waters' heavy metal elements has provided evidence that the waters have experienced environmental pollution. Environmental pollution is the entry or inclusion of living things, substances, energy, and other components into the environment by human activities so that they exceed the established environmental quality standards (Law of 2009 No. 32). The evidence indicates that marine mining activities cannot be conducted side by side with activities for marine tourism and fisheries.

Marine tourism activities that have developed around the Tanah Merah coast waters are on the island of Ketawai with swimming, snorkeling, diving, panoramic views of the small island, and turtle landing tours. Besides, mangrove tours around the mouth of the Kurau river. This marine tourism activity is closely related to good water quality conditions where the most influential hydro-oceanographic parameters include turbidity or clarity (brightness) of seawater, chemical content, temperature, salinity, and depth. Turbidity from seawater can affect coral reefs growth, where the biota is one of the most attractive marine tourism objects. The more turbid waters, the growth of coral reefs can be hampered, and the more siltation, siltation can occur and coral polyps can be closed so that the chance of bleaching and damage to coral reefs will increase. The movement of water affects coral growth because the movement of water for aquatic organisms is related to oxygen supply (Osingan *et al.*, 2011) and food (Sebens *et al.*, 1998). Also, the process of hydrodynamics (currents and turbulence) can reduce the level of attack on coral-eating biota (Lenihan *et al.*, 2015). Although corals have a mechanism for cleaning up sediment deposits, excessive sediment deposition will affect coral growth (Roy and Smith, 1971). The content of heavy metals (Pb and Cu) in the waters can cause bioaccumulation and biomagnification in marine biota including in especially filter-feeding bivalves (Ip *et al.*, 2005) and coral reefs (Setiawan, 2013). In addition to coral reefs, heavy metals can accumulate in mangroves. *Avicennia marina* is a type of mangrove that absorbs the heaviest metals with a Pb content of 24.2 ppm and Cu of 71.2 ppm (Setiawan, 2013).

Fishery activities that have developed around the coast of Tanah Merah are fishermen who catch various types of fish and land around the West Kurau river's mouth. Also, there is fish farming around Semujur Island, including grouper farming. In that situation change, conservation of biodiversity and degradation of fish habitat assumes the highest priority gradually (Mohan *et al.*, 2016). This fishery activity is closely related to the condition of good water quality, where the most influential hydro-oceanographic parameters include turbidity or clarity (brightness) of seawater, chemical element content, temperature, salinity, and depth. Turbid seawater can affect the distance of fish caught by fishermen. The more turbid waters in a wide area, the chances of catching fish are increasingly complicated and further its reach, which can increase fishermen's fuel costs. Aquaculture requires good water quality conditions, including clear, and does not contain heavy metal elements in the waters. The content of heavy metals (Pb and Cu) in the waters can lead to bioaccumulation and biomagnification in marine biota, including grouper farming. Economically, this condition has impacted decreasing income for fishermen and grouper farmers (Nurtjahya *et al.*, 2019; Ramadona *et al.*, 2020). Research by Sulista *et al.* (2019) states that income from fishing activities is higher (IDR 100,000 / day) compared to income from tin mining (IDR. 150,000/3 months). The value of heavy metals (Pb) in grouper meat of 6,500 mg / l indicates that this value is far above the quality standard threshold value of 0.050 mg / l with a difference of 6,450 mg / l above the quality standard, it is suspected that grouper may have experienced biomagnification from the food he ate (Adibrata *et al.*, 2013), found heavy metal concentrations (Pb and Cu) in fish and shellfish (Arifin, 2011). Besides, the presence of fish parasites such as ectoparasites attached to the fish skins of floating net cage systems (KJA) is thought to be related to the hydro-oceanographic dynamics such as current conditions and water temperatures (Adibrata *et al.*, 2020). The weaker water currents and the lower the water temperature can increase ectoparasites against fish in floating cage, this is certainly very detrimental to grouper fish farmers.

The distribution of heavy metals in the sea under conditions of weak water currents is homogeneous for the east monsoon (April - October). Comparative research is needed to determine the condition of heavy metals distribution in the western season, where the flow of waters is greater (Arifin, 2011) states that seasonal factors do not influence the distribution of heavy metals. Tidal water plays a significant role where this research is carried out on the tide's condition to the tide so that tidal currents strongly influence the water flow. Waters at high tide indicate that the mass transfer of water from the edge tends towards the middle with a certain angle. This mass of water certainly carries the burden of heavy metal pollutants that have been stirred due to mining at sea.

Furthermore, Arifin (2011) states that dissolved heavy metals are generally relatively low with the following ranges, Pb (1.0 - 26.0 $\mu\text{g L}^{-1}$) and Cu (1-2.0 $\mu\text{g L}^{-1}$). Levels of heavy metals Pb, Cd, Cu, Hg, and As the surface of Natuna waters are so small that they are still below the quality standard and concluded that they are still not polluted (Sagala *et al.*, 2014). Ahmad (2013) states that the heavy metals in sediments based on the Pollution Load Index and Geoaccumulation Index are still lower than the permitted threshold value for sediments. Some literature shows it dissolved heavy metals in the water column are affected and scattered according to water flow patterns. The dynamics of the distribution of heavy metals are not too far-reaching because of weak water currents so that the rate of dilution and sedimentation rate is slow, and stated that the waters had been polluted where the highest contamination value is indicated by the location closest to the source of marine mining and is weakening at the location farthest from the source polluter.

In summary, this study provided the closest and farthest Pb and Cu values from the community tin mining sources were 0.16 mg/l; 0.03 mg/l and 0.02 mg/l; <0.003 mg/l. Hydro-oceanographic conditions in the form of the highest to lowest water currents of 0.03 m/s - 0.001 m/s indicate that heavy metals in the waters are slow to be diluted. However, the

Pb value is above the water quality threshold according to seawater quality standards for tourism marine and marine biota and Cu values that are already above and some that are still below seawater quality standards applicable in Indonesia. Quality Standards based on Ministry of Environment Decree of 2004 No. 51 Appendix II for marine tourism and III for marine biota. This condition has threatened the sustainability of marine tourism and fishery activities, especially in areas located near tin mining sites. The research result shows that to represent the east monsoon, the dynamics of heavy metals' distribution are not too far-reaching due to weak water currents, the dilution and sedimentation rate is slow. It is stated that the waters are polluted where the highest contamination value is indicated by the location closest to the source of marine mining and further weakened at the location farthest from the source of pollutants. The results of this study have complemented the literature, the evidence provided confirms that it is necessary to regulate activities and site of tin mining as a win-win solution so that industrial interests and local community activities can continue to exist side by side and avoid conflicts in the use of open waters areas.

ACKNOWLEDGMENTS

This research was funded by the University Level Lecturer Research Program (PDTU) of the University of Bangka Belitung (UBB), with Contract Number 189.I/UN.50.3.1/PP/2019. Thank you to UBB so that this research can be published and become one of the references for similar research, especially in Bangka Belitung.

REFERENCES

- Adibrata, S., Kamal, M.M., & Yulianda, F. 2013. Daya dukung lingkungan untuk budidaya kerapu (Famili Serranidae) di perairan Pulau Pongok Kabupaten Bangka Selatan. *Jurnal pesisir dan pulau-pulau kecil* 2(1):43-58.
- Adibrata, S., Yusuf, M., & Manullang, C. 2020. Ectoparasite Prevalences of Grouper Fish (*Epinephelus fuscogutatus* x *Epinephelus polyphekadion*) Cultured in Floating Net Cages. *Indonesian Journal of Marine Sciences* 25(1):23-30.
- Ahmad, F. 2013. Distribusi dan prediksi tingkat pencemaran logam berat (Pb, Cd, Cu, Zn, dan Ni) dalam sedimen di perairan Pulau Bangka menggunakan Indeks Beban Pencemaran dan Indeks Geoakumulasi. *J. Ilmu dan Teknologi Kelautan Tropis* 5(1):170-181.
- Ahmed, M.M., Doumenq, P., Awaleh, M.O., Syakti, A.D., Asia, L., & Chiron, S. 2017. Levels and sources of heavy metals and PAHs in sediment of Djibouti-city (Republic of Djibouti). *Marine Pollution Bulletin* 1-7.
- Arifin, Z. 2011. Konsentrasi logam berat di air, sedimen, dan biota di Teluk Kelabat, Pulau Bangka. *J. Ilmu dan Teknologi Kelautan Tropis* 3(1):104-114.
- Baristand Laboratory. 2019. Test Report No. 391/BPPI/Baristand-Palembang-SERT.1/Up-0377 s/d Up-0389/V/2019. Palembang Institute For Industrial Research and Standarization. Baristand Industry Palembang Laboratory.
- Bibby, R.L., & Webster-Brown, J.G. 2006. Trace metal adsorption onto urban stream suspended particulate matter (Auckland region, New Zealand). *Appl. Geochem* 21:1135-1151.
- Bidayani, E., Kurniawan., Anggeraini, L., & Aisyah, S. 2019. Utilization Conflict Analysis of Fisheries Resources with Tin Mining and Marine Tourism in East Coast of Bangka Island. In Proceedings of the 13th International Interdisciplinary Studies Seminar, IISS.
- Cirik, Y., Bekci, Z.M., Buyukates, Y., Ilknur, A., & Merdivan, M. 2012. Heavy metals uptake from aqueous solutions using marine algae (*Colpomenia sinuosa*): kinetics and isotherms. *Chem and Ecol* 28(5):469-480.
- Hamad, S.H., Schauer, J.J., Shafer, M.M., Al-Raheem, A.E., & Satar, H. 2012. The distribution between the dissolved and the particulate forms of 49 metals across the Tigris River, Baghdad, Iraq. *The Scientific World J.* 2012:1-13.
- Ip, C.C.M., Li, X.D., Zhang, G., Wong, C.S.C., & Zhang, W.L. 2005. Heavy metal and Pb isotopic compositions of aquatic organisms in the Pearl River Estuary, South China. *Environmental Pollution* 138:494-504. PMID: 15970366 <https://doi.org/10.1016/j.envpol.2005.04.016>
- [Law of 2007 No. 27 juncto Law of 2014 No. 1]. Pengelolaan Wilayah Pesisir dan Pulau-pulau Kecil. Lembaran Negara RI Tahun 2007 No. 84 dan Lembaran Negara RI Tahun 2014 No. 2. Jakarta.
- [Law of 2009 No.32]. Perlindungan dan Pengelolaan Lingkungan Hidup. Lembaran Negara RI Tahun 2009 No. 140. Jakarta.
- Lenihan, H.S., Hench, J.L., Holbrook, S.J., Schmitt, R.J., & Potoski, M. 2015. Hydrodynamics influence coral performance through simultaneous direct and indirect effects. *Ecology* 96(6):1540-1549.
- Maslukah, L. 2013. Hubungan antara Konsentrasi Logam Berat Pb, Cd, Cu, Zn dengan Bahan Organik dan Ukuran Butir dalam Sedimen di Estuari Banjir Kanal Barat, Semarang. *Buletin Oseanografi Marina* 2:55-62.
- [Ministry of Environmental Decree of 2004 No. 51]. Keputusan Menteri Negara Lingkungan Hidup. 2004. Baku Mutu Air Laut untuk Biota Laut. Jakarta.
- Mohan., Divya, S., Arumugam, S., Jesslin, G., Edwin, T.P., & Soranam, R. 2016. Analysis of Water Quality Parameters and Heavy Metals Concentration from the Selected sites of Tamiraparani River, Tirunelveli District, Tamilnadu, India. *Int. Res. J. Environment Sci.* 5(7):47-54, E-ISSN 2319-1414.

- Najamuddin., Prartono, T., Sanusi, H.S., & Nurjaya, I.W. 2016. Seasonal distribution and geochemical fractionation of heavy metals from surface sediment in a tropical estuary of Jeneberang River, Indonesia. *Mar. Pollut. Bull.* 111:456-462.
- Nurtjahya, E., Jennifer, F., Umroh., & Fournita, A. 2017. The Impact of Tin Mining in Bangka Belitung and its Reclamation Studies. *MATEC Web Conferences* 101.04010. DOI: 10.1051/mateconf/201710104010.
- Osingan, R., Schutter, M., Griffioen, B., Wijffels, R.H., Verreth, J.A.J., Shafir, S., Henard, S., Taruffi, M., Gili, C., & Lavorano, S. 2011. The biology and economics of coral growth. *Marine Biotechnology (New-York)* 13:658–671.
- Puttiwongrak, A., Suteerasak, T., Mai, P.K., Hashimoto, K., Gonzalez, J.C., Rattanakom, R., & Prueksakorn, K. 2019. Application of multi-monitoring methods to investigate the contamination levels and dispersion of Pb and Zn from tin mining in coastal sediments at Saphan Hin, Phuket, Thailand. *Journal of Cleaner Production* 218:108-117.
- Ramadona, T., Rengi, P., Warningsih, T., Septya, F., Ngesti, S.M., & Purnomo, A. 2020. Does Capture Fisheries In The Vicinity of Tin Mining Area Sustainable?: A case study in Bangka Island, Indonesia. In *IOP Conference Series: Earth and Environmental Science* (Vol. 430, No. 1, p. 012015). IOP Publishing
- Roy, K.J., & Smith, S.V. 1971. Sedimentation and coral reef development in turbid water: Fanning Lagoon. *Pacific Science* 25:234-248.
- Sagala, S.L., Bramawanto, R., Kuswardani, A.R.T.D., & Pranowo, W.S. 2014. Distribusi logam berat di perairan Natuna. *J. Ilmu dan Teknologi Kelautan Tropis* 6(2):297-310.
- Sebens, K.P., Grace, S.P., Helmuth, B., Maney Jr., E.J., & Miles, J.S. 1998. Water flow and prey capture by three scleractinian corals, *Madracis mirabilis*, *Montastrea cavernosa*, and *Porites porites*, in a field enclosure. *Marine Biology* 131(2):347-360.
- Setiawan, H. 2013. Accumulation and distribution of heavy metals in mangrove vegetation in coastal waters of South Sulawesi. *J. Ilmu Kehutanan* 7(1):12-24.
- Sulista, S., Ibrahim, I., & Pratama, S. 2019. Accommodation, Resistance and Divided Community: Study of the Dynamics of Offshore Tin Mining Conflict Between the Fishermen of the Coastal Area and Companies in Bangka Island. *PEOPLE: International Journal of Social Sciences* 5(3):275-296.
- Syakti, A.D., Demelas, C., Hidayati, N.V., Rakasiwi, G., Vassalo, L., Kumar, N., Prudent, P., & Doumenq, P. 2015. Heavy metal concentrations in natural and human-impacted sediments of Segara Anakan Lagoon, Indonesia. *Environ Monit Assess* 187:4079:1-15.
- Thomann, R.V., & Muller, J.A. 1987. *Principles of surface water quality modeling and control*. New York (US): Harper & Row.
- Valeriani, D., & Wardhani, R.S. 2015. Alternative tourism in the belitung island with the blue tourism concepts as the effort to achieve the quality of life, quality of opportunity and quality of experience. *Mediterranean Journal of Social Sciences* 6(5 S5), 46.
- Wang, Z., Wang, Y., Chen, L., Yan, C., Yan, Y., & Chi, Q. 2015. Assessment of metal contamination in coastal sediments of the Maluan Bay (China) using geochemical indices and multivariate statistical approaches. *Mar. Pollut. Bull.* 99: 43–53.
- Yulianti, A. 2019. Potentials of Leading Sectors In Bangka Belitung Island Province On 2013- 2017. *Jurnal Ekonomi Pembangunan* 17(1):32-39.

The contamination of heavy metals (Pb and Cu) at tin sea mining field and its impact to marine tourism and fisheries

ORIGINALITY REPORT

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

8%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	worldwidescience.org Internet Source	1%
2	ejournal.undip.ac.id Internet Source	1%
3	www.science.gov Internet Source	1%
4	lenny-nevitasary-fst17.web.unair.ac.id Internet Source	1%
5	Avirut Puttiwongrak, Thongchai Suteerasak, Phuong Khanh Mai, Kiyota Hashimoto et al. "Application of multi-monitoring methods to investigate the contamination levels and dispersion of Pb and Zn from tin mining in coastal sediments at Saphan Hin, Phuket, Thailand", Journal of Cleaner Production, 2019 Publication	1%
6	Submitted to Taibah University Student Paper	1%

7	Najamuddin, Tri Prartono, Harpasis S. Sanusi, I Wayan Nurjaya. "Seasonal distribution and geochemical fractionation of heavy metals from surface sediment in a tropical estuary of Jeneberang River, Indonesia", Marine Pollution Bulletin, 2016 Publication	1%
8	www.barrie.ca Internet Source	1%
9	li01.tci-thaijo.org Internet Source	1%
10	iserd.net Internet Source	1%
11	ojs.omniakuatika.net Internet Source	1%
12	www.scilit.net Internet Source	<1%
13	id.scribd.com Internet Source	<1%
14	Ghada F. El-Said, Doaa H. Youssef. "Ecotoxicological impact assessment of some heavy metals and their distribution in some fractions of mangrove sediments from Red Sea, Egypt", Environmental Monitoring and Assessment, 2012 Publication	<1%

15

Andri Taufick Rizaluddin, Yusup Setiawan.
"POTENCY TO RECYCLE HYDRAPULPER
REJECT WASTE OF PAPER MILL INTO
COMPOSITE PARTICLEBOARD", Jurnal Riset
Teknologi Pencegahan Pencemaran Industri,
2019

Publication

<1%

16

www.scribd.com

Internet Source

<1%

17

L Kurnianti, H Setiani, S H Ramadianti, R F Sari.
"Seashore Community Engagement Program for
A Clean Seawater at Banyu Biru Village", IOP
Conference Series: Earth and Environmental
Science, 2020

Publication

<1%

18

es.scribd.com

Internet Source

<1%

19

Hans Thodsen, Jes Jessen Rasmussen, Brian
Kronvang, Hans Estrup Andersen, Anders
Nielsen, Søren Erik Larsen. "Suspended matter
and associated contaminants in Danish
streams: a national analysis", Journal of Soils
and Sediments, 2019

Publication

<1%

20

ejournal2.undip.ac.id

Internet Source

<1%

21 rjls.ub.ac.id
Internet Source

<1%

22 www.freepatentsonline.com
Internet Source

<1%

23 Rachel L. Wattimena, Debby A.J. Selanno,
Semuel F. Tuhumury, Juliana W. Tuahatu.
"Analysis of Heavy Metal Content (Pb) on
Waters and Fish at The Floating Cages BPPP
Ambon", E3S Web of Conferences, 2018
Publication

<1%

24 Sadaf Nazneen, Shubhra Singh, N.Janardhana
Raju. "Heavy metal fractionation in core
sediments and potential biological risk
assessment from Chilika lagoon, Odisha state,
India", Quaternary International, 2018
Publication

<1%

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On

The contamination of heavy metals (Pb and Cu) at tin sea mining field and its impact to marine tourism and fisheries

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
